

CH 5

for earth gravity

center of mass = center of gravity

Centroid = center of Area

$$\bar{x} = \frac{\int x dm}{m} \quad / \quad \bar{y} = \frac{\int y dm}{m}$$


$$\bar{x} = \frac{\int x dA}{A} \quad / \quad \bar{y} = \frac{\int y dA}{A} \quad \text{For 2d}$$

$A\bar{x} = Q_y$ = First moment of Area with respect to y

$$A\bar{x} = Q_y = \int x dA$$

هذا يعني أن Q_y هي العزم الأول لـ A حول المحور y
 حيث $\bar{x}$ هو المركز الجذبوي لـ A في المحور x
 Moment of force Q_y = Moment of Area A about y


$Q_x = A\bar{y} = \int y dA$ = first moment of Area with respect to x



$$Q_y = A\bar{x} = \int x dA$$

$$Q_x = A\bar{y} = \int y dA$$

For thin elements



$$A = \text{cross section} \times L$$

$$L = \text{length}$$

$$dl = \text{small length}$$

$$\rho = \text{density}$$

$$M_x = A \rho g \bar{y} = \int y \times A \times \rho \times g \times dl$$

$$L \bar{y} = \int y dl$$


$$\bar{y} = \frac{\int y dl}{L}$$

$$\bar{x} = \frac{\int x dl}{L}$$

$$\bar{x} L = \text{first moment of length with respect to } (y) \text{ axis} = \int x dl$$

the same for $\bar{x}$ axis

Centroid by integration



$$\bar{x}A = \int x_{el} dA$$

$$\bar{x}A = \int x * y dx$$

$$\bar{x}A = \int_{x_1}^{x_2} x * f(x) dx$$

$$\bar{x} = \frac{\int_{x_1}^{x_2} x f(x) dx}{\int dA}$$


$$\bar{y}A = \int y_{el} dA$$

$$\bar{y}A = \int \frac{y}{2} * y * dx$$

$$\bar{y} \int dA = \int \frac{(f(x))^2}{2} dx$$


⚠ مهم جداً انه نحتاج عندنا شرحاً
عمودية على المحور فإني الشتر فيه
نبي نشتغل للارتفاع على $y/2$ مثلاً

⚠ ممكن انه شرحاً عمودية او موازية
للمحور !! مش مهم



$$\bar{x}A = \int_{y_1}^{y_2} \frac{(x_2 + x)}{2} * (x_2 - x) dy$$

$$\bar{y}A = \int y (x_2 - x) dy$$



Polar coordinates

$$r = \frac{2}{3} \quad \theta$$

$$x_{el} = \frac{2}{3} r \cos \theta$$

$$y_{el} = \frac{2}{3} r \sin \theta$$

$$dA = \frac{1}{2} r^2 \sin d\theta$$

$$\bar{X}A = \int x_{el} dA$$

$$= \int \frac{2}{3} r \cos \theta \times \frac{1}{2} r^2 d\theta$$

but $\sin d\theta = d\theta$

$$\bar{X} = \frac{\int \frac{1}{3} r^3 \cos \theta d\theta}{A}$$


$$yA = \int y_{el} dA$$

$$= \int \frac{2}{3} r \sin \theta \times \frac{1}{2} r^2 d\theta$$

$$\bar{Y} = \frac{\int \frac{1}{3} r^3 \sin \theta d\theta}{A}$$

فقط
حساب
تفاضل و
تكامل

sample 2



Find $\bar{x}, \bar{y}$ of the centroid

Find Q_x / Q_y

1 $k = ?$

at $x = a \rightarrow y = b$

$b = a^2 k$

$\boxed{\frac{b}{a^2} = k}$

2 $A = \int dA = \int_0^a y dx = \int_0^a kx^2 dx = \frac{kx^3}{3} \Big|_0^a = \frac{ka^2 \cdot a}{3}$
 $= \frac{ab}{3}$

3 $\bar{x} = \frac{\int x dA}{A} = \frac{\int x \cdot kx^2 dx}{A} = \frac{\int kx^3 dx}{A}$
 $= \frac{kx^4}{4} \Big|_0^a = \frac{ka^2 \cdot a^2}{4A} = \frac{a^2 \cdot 3}{4 \cdot b}$



$\boxed{\bar{x} = \frac{3}{4} a}$

4 $\bar{y} = \frac{\int y dA}{A} = \frac{3 \int \frac{kx^2}{2} \cdot kx^2 dx}{ab} = \frac{3 \int \frac{k^2 x^4}{2} dx}{ab}$
 $= \frac{k^2 x^5}{10(ab)} \Big|_0^a = \frac{k^2 a^5}{10 \cdot b}$
 $= \frac{b \cdot 3}{10} = \boxed{\frac{3b}{10} = \bar{y}}$

$$Q_x = A y = \frac{A y^2}{2}$$


$$Q_y = A x = \frac{b a^2}{4}$$

Horizontal
or
Vertical



in vertical it's hard
because we'll need
2 integrals

Symmetry




$$Q_y = 0$$

because the centroid
is on y axis because
of symmetry



so the distance between
centroid and axis = 0

$$\bar{x}A = 0 * A = 0$$



the same for $Q_{BB'} = \overline{PP'A} = 0$


اذا وجدنا * خطي تماثل متعامدين
انما ال centroid هو مركز
التقاطع



نقطة تماثل متعامدين


هذه النقطة هي ال centroid

Composite plates and Areas




$$\bar{X}A = \bar{X}_1 A_1 + A_2 \bar{X}_2 + \bar{X}_3 A_3$$


OR




$$= \frac{\bar{X}_{sq} A_{sq} - A_3 \bar{X}_3 - A_2 \bar{X}_2}{A - A_3 - A_2}$$



$$\frac{2A \times 2 - A \times 1}{A}$$



sample
determine the first moment
with respect to X & Y
and centroid



$$A = \overset{A_3}{3600} + \overset{A_2}{9600} + \overset{A_1}{5655} - 5026.5$$

$$= 13828 \quad \checkmark$$

$$\bar{x}A = 180 \times 3600 + 9600 \times 60 + 60 \times 5655 - 5026 \times 60$$

$$\boxed{\bar{x}A = Q_y = 757740}$$

$$Q_x = \bar{x}y = 3600 \times -20 + 9600 \times 40 + 5655 \times 105.5 - 80 \times 5026$$


$$\boxed{Q_x = 506522.5}$$

$$\boxed{\bar{x} = 54.8}$$

$$\boxed{\bar{y} = 36.6}$$

تأكد من
صحة
الحل دوماً

Areas and Volumes of Revolution in general




$$A = 2\pi \bar{y} L$$



$$\bar{y} = \frac{2r}{\pi} =$$

$$2\pi * \frac{2r}{\pi} * \frac{1}{2} 2\pi r$$

$$= 4\pi r^2$$



more general




$$\int 2\pi y dl$$

$$= 2\pi \int y dl \leftarrow \text{first way}$$

$$= 2\pi * \bar{y} * l$$


Volumes / $(2\pi \bar{y} A)$



$$= \text{sphere} = 2\pi \bar{y} A$$

نقطه المركز
نقطه مركز ثقل
Centroid

example



the outside pulley is 0.8 m and the cross section of its rim is shown

Knowing that the pulley is made of steel and that the density of steel is $\rho = 7.85 \times 10^3 \text{ kg/m}^3$

determine the mass and weight of the rim

first find Volume = $A \times 2\pi \bar{y}$

$$A = 1200 + 2000 = 3200 \text{ mm}^2 = 3.2 \times 10^{-3} \text{ m}^2$$

$$\bar{y} = \frac{A_1 y_1 + A_2 y_2 + A_3 y_3}{A} = \frac{1200 \times 365 + 2000 \times 300}{3.2 \times 10^3}$$

$$= 380.625 \text{ mm} = 0.380625 \text{ m}$$

$$V = 2\pi \bar{y} A = 7.653 \times 10^{-3} \text{ m}^3$$

$$= 60 \text{ kg} \quad \text{mass}$$


$$= 588.7 \text{ kg} \quad \text{weight}$$

distributed force

$$N/m \times \text{length} = \text{force}$$

$$\text{or } \int \text{load} \cdot dx$$


to find moment or reactions
we need to find a single force
at the centroid of the shape
of distributed load



$$\text{load} = \int dw$$

$$dw = F dx$$


$$\text{load} = \int F dx = \text{Area} \\ = \int dA = A$$




$$A \cdot \bar{x} = \int x dA$$

= moment
of load
at B

Sample Problem



Find reactions




$$\sum F_x = 0$$

$$B_x = 0 \checkmark$$

$$M_A = F\bar{x} - B_y L$$

$$\text{or } A_x - B_x \times 6$$

Finding the $\bar{x}$



$$6 \times 3 \times 1500 + 4 \times 3600 \times 3 = A\bar{x}$$



$$63000 - 6B_y = 0$$

$$B_y = 10500 \uparrow \text{N}$$


$$A_y + B_y - 18000 = 0$$

$$A_y = 7500 \uparrow \text{N}$$

what if



انتي انا
عند الـ
للـ




Home work 4

ch5

✓

Prob 4



A_1 is symmetric around two perpendicular axes


Axis 1: $y = 75 \text{ mm}$

Axis 2: $x = 60$

so $C_1 = (60, 75)$

$$A_1 = 2 * 60 * 75 = 9 * 10^3 \text{ mm}^2$$

A_2 is a triangle



we have 2 perpendicular bases for triangle A_2

so we can apply the rule $\bar{y} = \frac{h}{3}$

once in x direction and another in y direction

$$C_2 = (20, 25) / A_2 = 2250$$


$$\bar{X} = \frac{X_1 A_1 + X_2 A_2}{A} = \frac{60 * 9 * 10^3 + 20 * 2250}{9 * 10^3 + 2250}$$

$$\bar{X} = 52$$

$$\bar{Y} = \frac{Y_1 A_1 + Y_2 A_2}{\Sigma A} = \frac{75 * 9 * 10^3 + 25 * 2250}{9 * 10^3 + 2250}$$


$$\bar{Y} = 65$$

$$C = (52, 65)$$



$A_3 = A_2 + A_1 = \text{square}$
with side r

9



$$x_3 = \frac{A_1 x_1 + A_2 x_2}{A_1 + A_2}$$

$$(A_1 + A_2) x_3 = A_1 x_1 + A_2 x_2$$

$$A_1 x_1 = (A_1 + A_2) x_3 - A_2 x_2$$

$$\therefore x_1 = \frac{A_3 x_3 - A_2 x_2}{A_3 - A_2}$$

$$\bar{x}_1 = \frac{r^2 * \frac{r}{2} - (r - \frac{4}{3\pi}) r \frac{\pi r^2}{4}}{r^2 - \frac{r^2 \pi}{4}}$$


$$\bar{x}_1 = 16.753 \text{ mm}$$

the shape is symmetric around axis $(y=x)$
then the point is on that axis then

$$\bar{y} = \bar{x} = 16.753 \text{ mm}$$

$$C = (16.75, 16.75)$$

P₁₃



$$A_2 = A_3 - A_1$$

$$= \frac{4 \times 8}{2} - \frac{4 \times 8}{3} = \frac{4 \times 8}{6} = 5.333 \text{ m}^2$$

$$x_2 = \frac{x_3 A_3 - x_1 A_1}{A_3 - A_1} = \frac{\frac{8}{3} \times 16 - 3 \times 0.8 \times \frac{4 \times 8}{3}}{\frac{4 \times 8}{6}}$$

$$x_2 = 3.2$$


$$y_2 = \frac{y_3 A_3 - y_1 A_1}{A_3 - A_1} = \frac{\frac{8}{3} \times 16 - 3 \times \frac{4 \times 8}{3}}{\frac{4 \times 8}{6}}$$

$$y_2 = 2$$

$$C = (3.2, 2)$$

Problem 30

P 30



$$M_1 + M_2 = 0$$

$$|M_1| - |M_2| = 0$$

when $|M_1| = |M_2|$

$$F_1 x_1 = F_2 x_2$$

$$M_1 \cdot x_1 = M_2 \cdot x_2$$


$$\lambda L_1 x_1 = \lambda L_2 x_2$$

$$L_1 x_1 = L_2 x_2$$

Portion ABC

Both BC And BA

have the same $\bar{x} = x_1$



$$L(AB) + L(BC) = 80 + \sqrt{80^2 + 60^2}$$

$$L_1 = 180$$

$$x_1 = 40 \text{ for the symmetry}$$


$$x_2 = \frac{L_2}{2} \text{ because of symmetry}$$

$$180 \cdot 40 = \frac{L_2^2}{2}$$

$$\sqrt{180 \cdot 40 \cdot 2} = L_2$$

$$L_2 = 120 \text{ mm}$$

P 35



$$\bar{x} = \frac{\int x dA}{A}$$

$$A = \int dA = \int_0^a (mx - kx^2) dx = \frac{mx^2}{2} - \frac{kx^3}{3} \Big|_0^a$$

$$= \frac{m}{2} x^2 \Big|_0^a - \frac{k}{3} x^3 \Big|_0^a = \left(\frac{ma^2}{2} - \frac{ka^3}{3} \right)$$

$$\int x dA = \int_0^a x(mx - kx^2) dx$$

$$= \int_0^a (mx^2 - kx^3) dx = \left(\frac{mx^3}{3} - \frac{kx^4}{4} \right) \Big|_0^a$$

$$= \left(\frac{ma^3}{3} - \frac{ka^4}{4} \right)$$

$$\bar{x} = \frac{\frac{4ma^3 - 3ka^4}{2} \cdot \frac{3ma^2 - 2ka^3}{x}}{A}$$

$$\bar{x} = \frac{a^2 (4m - 3ka)}{2a^2 (3m - 2ka)}$$

at Point (B) $y_1 = y_2 \rightarrow (mx = kx^2) \Big|_a$

$$m = kb$$

$$m = \frac{h}{a}$$


$$k = \frac{h}{a^2}$$

$$k = \frac{m}{a}$$

$$\frac{a}{2} \cdot \frac{4 \times \frac{h}{a} - 3 \times \frac{h}{a^2} a}{3 \times \frac{h}{a} - 2 \times \frac{h}{a^2} a} = \frac{a}{2} \cdot \frac{\frac{h}{a}}{\frac{h}{a}}$$

$$\bar{x} = \frac{a}{2}$$

$$\bar{y} = \frac{\int y dA}{A}$$



$$\int y da = \int_0^a \frac{mx + kx^2}{2} \times (mx + kx^2) dx$$

$$= \int \frac{(mx)^2 + k^2 x^4}{2} dx$$


$$= \frac{1}{2} \int m^2 x^2 + k^2 x^4 dx = \left(\frac{m^2 x^3}{3} + \frac{k^2 x^5}{5} \right) \Big|_0^a$$

$$= \frac{\frac{m^2 a^3}{3} + \frac{k^2 a^5}{5}}{\frac{ma^2}{2} + \frac{ka^3}{3}} = \frac{\frac{h^2 a^3}{6} + \frac{h^2 a^5}{10}}{\frac{ha^2}{2} + \frac{ha^3}{3}}$$

$$= \frac{\frac{4}{15} h^2 a^3}{\frac{5ha^2}{6} + \frac{3ha^3}{6}} = \boxed{\frac{4h}{10} = \bar{y}}$$

$$C = \left(\frac{a}{2}, \frac{4h}{10} \right)$$

P55



$$V = 2 \times \text{Volume of cylinder} + 2 \times \text{Volume of ends}$$

$$= (2A \times L) + 2(A \times \pi R)$$

$$= 2A(L + \pi R)$$

$$= 2 \times 3^2 \times \pi (30 + \pi \times 10)$$

$$= 3472.9 \text{ mm}^3$$

$$\approx 3.473 \times 10^{-6} \text{ m}^3$$


$$\text{Surface area} = 2 \times \text{surface area of cylinder} + 2 \times \text{surface of ends}$$

$$= 2(2\pi r L) + 2(2\pi R^2)$$

$$= 2\pi r(L + \pi R)$$

$$= 2315.3 \text{ mm}^2$$

P69



$$F_1 =$$

$$\frac{1}{2} \times 12 \times 50 + 50 \times 20$$

$$= 1300 \text{ lb}$$

$$F_1 \bar{x} = F_1 x_1 + F_2 x_2$$

$$300 \times \frac{2}{3} \times 12 + 1000 \times 22$$

$$1300$$



$$= 18.76 \text{ in}$$

$$F_{Ay} = 1300 - 400 = 900 \text{ lb} \uparrow \quad [A_x = 0]$$

$$M_A = 22000 + 2400 - 400 \times 38$$

$$M_A = +9200 \text{ lb} \cdot \text{in} \text{ counter clockwise}$$

Prob 71



$$A_y + B_y = +375 \uparrow$$

$$\sum M_A = 0 = 600 * 1.15 - 975 * 1.4 + B_y * 2.5$$

$$0 = -675 + B_y * 2.5$$

$$B_y = +270 \text{ N} \uparrow$$

$$A_y = +105 \text{ N} \uparrow$$

Prob 76


$$F_1 = 12 * 100 * \frac{1}{2} = 600 \text{ Ib}$$

$$F_2 = 300 * 12 = 3600 \text{ Ib}$$

$$M_c = F_1 * \frac{2}{3} * 12 + 6 * F_2 - B_y * 7$$

$$7B_y = 4800 + 21600$$

$$B_y = +3771.4 \text{ Ib} \uparrow$$



$$\sum F_y = 0 \rightarrow F_1 + F_2 + B + C = 0$$

$$-4200 + 3771.4 + C = 0$$

$$C = -428.6 \text{ Ib} \downarrow$$