

2.3 + 2.4

The Precise Definitions of Limits

29


Def: Example $f(x) = 2x + 1$ $\epsilon = 2 \Rightarrow \delta = 1$

Let $f(x)$ be defined on an open interval about x_0 , except possibly at x_0 itself. We say $f(x)$ approaches the limit L as x approaches x_0 , and we write:

$$\lim_{x \rightarrow x_0} f(x) = L$$

if for every number $\epsilon > 0$, there exists a corresponding number $\delta > 0$ such that for all

$$0 < |x - x_0| < \delta \Rightarrow |f(x) - L| < \epsilon$$



Example: show that $\lim_{x \rightarrow 1} (5x - 3) = 2$

Let $\epsilon > 0$, we must find $\delta > 0$ s.t for all x if $|x - x_0| < \delta$ then $|f(x) - L| < \epsilon$

$x_0 = 1$, $L = 2$, $f(x) = 5x - 3$

That is we need to find $\delta > 0$ s.t for all x if $|x - 1| < \delta$ then


$$|(5x - 3) - 2| < \epsilon$$

$$\Rightarrow |5x - 5| < \epsilon \Rightarrow 5|x - 1| < \epsilon \Rightarrow |x - 1| < \frac{\epsilon}{5}$$

Thus, we can take $0 < \delta \leq \frac{\epsilon}{5}$ because if

STUDENTS-HUB.COM

if $|x - 1| < \delta = \frac{\epsilon}{5}$, then $|(5x - 3) - 2| = |5x - 5| = 5|x - 1| < 5 \cdot \frac{\epsilon}{5} = \epsilon$



To find the maximum δ^* in which $0 < \delta \leq \delta^*$, we may use $f(x_0 - \delta^*) = L - \epsilon$ or $f(x_0 + \delta^*) = L + \epsilon$

In the previous example:

$$f(x_0 - \delta^*) = L - \epsilon \Rightarrow f(1 - \delta^*) = 2 - \epsilon \Rightarrow$$

$$5(1 - \delta^*) - 3 = 2 - \epsilon \Rightarrow 5 - 5\delta^* - 3 = 2 - \epsilon \Rightarrow 2 - 5\delta^* = 2 - \epsilon$$

$$\Rightarrow \delta^* = \frac{\epsilon}{5}$$


Uploaded By: Malak Obaid

Example: show that $\lim_{x \rightarrow x_0} x = x_0$

$f(x) = x$ and $L = x_0$


(30)

Let $\epsilon > 0$, we need to find $\delta > 0$ such that for all x
if $|x - x_0| < \delta$ then $|x - x_0| < \epsilon$
Take $0 < \delta \leq \epsilon$. This proves that $\lim_{x \rightarrow x_0} x = x_0$



Example: Prove that $\lim_{x \rightarrow x_0} k = k$

Let $\epsilon > 0$, we need to find $\delta > 0$ s.t for all x
if $|x - x_0| < \delta$ then $|k - k| < \epsilon$.



$0 < \epsilon$ This is true for any $\delta > 0$.

This proves that $\lim_{x \rightarrow x_0} k = k$

How to find δ for a given f, L, x_0 and ϵ :

Two steps to find $\delta > 0$ s.t for all x

if $|x - x_0| < \delta$ then $|f(x) - L| < \epsilon$

1) Solve the inequality $|f(x) - L| < \epsilon$ to find an open interval (a, b) about x_0 on which the inequality holds for all $x \neq x_0$.

2) Find $\delta > 0$ that places the open interval $(x_0 - \delta, x_0 + \delta)$ centered at x_0 inside the interval (a, b) . The inequality $|f(x) - L| < \epsilon$ will hold for all $x \neq x_0$ in this δ -interval.

STUDENTS-HUB.COM

Uploaded By: Malak Obaid

Example: Prove that $\lim_{x \rightarrow 2} f(x) = 4$ where $f(x) = \begin{cases} x^2, & x \neq 2 \\ 1, & x = 2 \end{cases}$

Let $\epsilon > 0$, we need to find $\delta > 0$ s.t for all x

if $|x - 2| < \delta$ then $|f(x) - 4| < \epsilon$

Step 1: ^{solve} $|f(x) - 4| < \epsilon$ to find an open interval about $x_0 = 2$
in which the inequality $|f(x) - 4| < \epsilon$ holds for all $x \neq x_0$


$\Rightarrow$ For $x \neq x_0 = 2$ we have $f(x) = x^2$

$$\Rightarrow |x^2 - 4| < \epsilon \Rightarrow$$

$$-\epsilon < x^2 - 4 < \epsilon \Rightarrow 4 - \epsilon < x^2 < 4 + \epsilon$$

$$\Rightarrow \sqrt{4 - \epsilon} < |x| < \sqrt{4 + \epsilon} \quad (\text{Assume } \epsilon < 4)$$

$$\sqrt{4 - \epsilon} < x < \sqrt{4 + \epsilon}$$



The inequality $|f(x) - 4| < \epsilon$ holds for all $x \neq 2$ in the open interval $(\sqrt{4 - \epsilon}, \sqrt{4 + \epsilon})$.

step 2 Find $\delta > 0$ that places the centered interval $(2 - \delta, 2 + \delta)$ inside the interval $(\sqrt{4 - \epsilon}, \sqrt{4 + \epsilon})$

Take $\delta = \min \{2 - \sqrt{4 - \epsilon}, \sqrt{4 + \epsilon} - 2\}$.

Thus, the inequality $|x - 2| < \delta$ will automatically place x between $\sqrt{4 - \epsilon}$ and $\sqrt{4 + \epsilon}$ to make $|f(x) - 4| < \epsilon$.

Example: Given that $\lim_{x \rightarrow c} f(x) = L$ and $\lim_{x \rightarrow c} g(x) = M$. } we now can prove Theorems
Prove that $\lim_{x \rightarrow c} (f(x) + g(x)) = L + M$.

Let $\epsilon > 0$, we need to find $\delta > 0$ s.t for all x
if $|x - c| < \delta$ then $|f(x) + g(x) - (L + M)| < \epsilon$.

$\Rightarrow$ since $\lim_{x \rightarrow c} f(x) = L$, $\exists \delta_1 > 0$ s.t for all x if $|x - c| < \delta_1$, then $|f(x) - L| < \frac{\epsilon}{2}$

STUDENTS-HUB.com

Uploaded By: Malak Obaid

$\Rightarrow$ since $\lim_{x \rightarrow c} g(x) = M$, $\exists \delta_2 > 0$ s.t for all x if $|x - c| < \delta_2$, then $|g(x) - M| < \frac{\epsilon}{2}$

step 1 $|f(x) + g(x) - (L + M)| = |(f(x) - L) + (g(x) - M)|$
 $\leq |f(x) - L| + |g(x) - M|$ Triangle Inequality
 $\leq \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon$

step 2 Take $\delta = \min \{\delta_1, \delta_2\}$, so that
 if $|x - c| < \delta$, then $|x - c| < \delta_1$, thus $|f(x) - L| < \frac{\epsilon}{2}$.
 if $|x - c| < \delta_2$ then $|x - c| < \delta$, thus $|g(x) - M| < \frac{\epsilon}{2}$. Thus