
Chapter TEN Photosynthesis



Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

Overview: The Process That Feeds the Biosphere

 Photosynthesis is the process that converts solar energy into chemical energy

 Directly or indirectly, photosynthesis nourishes almost the entire living world Autotrophs sustain themselves without eating anything derived from other organisms

 Autotrophs are the *producers* of the biosphere, producing organic molecules from CO₂ and other inorganic molecules

 Almost all plants are photo autotrophs, using the energy of sunlight to make organic molecules from H₂O and CO₂ Photosynthesis occurs in plants, algae, certain other protists, and some prokaryotes

 These organisms feed not only themselves but also most of the living world

Photoautotrophs (b) Multicellular alga (a) Plants (d) Cyanobacteria 40 μm Unicellular 10 µm protists 1 µm e Purple STUDENTS-HUB.com sulfur

 Heterotrophs obtain their organic material from other organisms

Heterotrophs are the consumers of the biosphere

 Almost all heterotrophs, including humans, depend on photoautotrophs for food and O₂

Concept 10.1: Photosynthesis converts light energy to the chemical energy of food


- Chloroplasts are structurally similar to and likely <u>evolved</u> from photosynthetic bacteria
- The structural organization of these cells allows for the chemical reactions of photosynthesis

Chloroplasts: The Sites of Photosynthesis in Plants

- Leaves are the major locations of photosynthesis
- Their green color is from chlorophyll, the green pigment within chloroplasts
- Light energy absorbed by chlorophyll drives the synthesis of organic molecules in the chloroplast
- CO₂ enters and O₂ exits the leaf through microscopic pores called stomata

- Chloroplasts are found mainly in cells of the mesophyll, the interior tissue of the leaf
- A typical mesophyll cell has 30–40 chloroplasts
- The chlorophyll is in the membranes of thylakoids (connected sacs in the chloroplast); thylakoids may be stacked in columns called grana
- Chloroplasts also contain stroma, a dense fluid

Zooming in on the location of photosynth esis in a plant.




Tracking Atoms Through Photosynthesis: *Scientific Inquiry*

 Photosynthesis can be summarized as the following equation:

$$6 CO_2 + 12 H_2O + Light energy \rightarrow C_6H_{12}O_6 + 6 O_2 + 6 H_2O_1$$

Tracking atoms through photosynthesis.



© 2011 Pearson Education, Inc.

The Splitting of Water

 Chloroplasts split H₂O into hydrogen and oxygen, incorporating the electrons of hydrogen into sugar molecules


```
- Plants: CO_2 + 2H_2O \square [CH_2O] + H_2O + O_2
```

Sulfur bacteria: CO₂ + H₂S □ [CH₂O] + H₂O +
 S₂

 CO2 is not split into C and O2!! This hypothesis was <u>cancelled</u> by van Neil in 1930s

Photosynthesis as a Redox Process

Photosynthesis is a redox process in which H₂O is oxidized and CO₂ is reduced



- Summary:
- H2O is split, & electrons are transferred <u>along with H ions</u> from H2O to CO2 reducing it to sugars!
- Endergonic reaction.

"absorbing energy in the form of work."

How this reaction relates to cellular respiration??




The Two Stages of Photosynthesis: A Preview


- Photosynthesis consists of the light reactions (the *photo* part) and Calvin cycle (the *synthesis* part)
- The light reactions (in the thylakoids):
 - Split H₂O
 - Release O₂
 - Reduce NADP⁺ to NADPH
 - Generate ATP from ADP by photophosphorylation


 The Calvin cycle (in the Stroma) forms sugar from CO₂, using ATP and NADPH


 The Calvin cycle begins with carbon fixation, incorporating CO₂ into organic molecules



An overview of photosynthesis: cooperation of the light reactions and the Calvin cycle.







Concept 10.2: The light reactions <u>convert</u> solar energy <u>to the</u> chemical energy of ATP and NADPH

Chloroplasts are <u>SOlar-powered</u>
 <u>chemical factories</u>

 Their thylakoids transform light energy into the chemical energy of ATP and NADPH

The Nature of Sunlight

- Light is a form of <u>electromagnetic</u> energy
 - = electromagnetic radiation

Light travels in rhythmic waves

Wavelength is the distance between crests of waves


Wavelength <u>determines</u> the type of electromagnetic energy

 The electromagnetic spectrum is the entire range of electromagnetic energy, or radiation

 Visible light consists of wavelengths (including those that drive photosynthesis) that produce colors we can see

 Light also behaves as though it consists of discrete particles, called photons


Fig. 10-6



Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

Photosynthetic Pigments: The Light Receptors

- Substances that absorb visible light
- Different pigments absorb different wavelengths
- Wavelengths that are not absorbed are reflected or transmitted
- Leaves appear green because <u>chlorophyll</u> reflects and <u>transmits</u> green light




© 2011 Pearson Education, Inc.

 A spectrophotometer measures a pigment's ability to absorb various wavelengths

 This machine sends light through pigments and measures the fraction of light transmitted at each wavelength An absorption spectrum is a graph plotting a pigment's light absorption versus wavelength

 The absorption spectrum of chlorophyll a suggests that Violet-blue and red light work best for photosynthesis

Fig. 10-9




 Chlorophyll a is the main photosynthetic pigment

 Accessory pigments, such as chlorophyll b, broaden the spectrum used for photosynthesis

 Accessory pigments called carotenoids absorb excessive light that would damage chlorophyll


Fig. 10-10




Excitation of Chlorophyll by Light

- When a pigment absorbs light, it goes from <u>a ground state to an excited state</u>, which is unstable
- When excited electrons fall back to the ground state, photons are given off, an afterglow called <u>fluorescence</u>

 If illuminated, an isolated solution of chlorophyll will fluoresce, giving off light and heat






(a) Excitation of isolated chlorophyll molecule ducation, Inc., publishing as Pearson Benjamin Cummings.

(b) Fluorescence

A Photosystem:


A Reaction-Center Complex Associated with Light-Harvesting Complexes

- A photosystem consists of a <u>reaction-center complex</u> (protein <u>complex</u>) surrounded by light-harvesting complexes
- The light-harvesting complexes (pigment molecules bound to proteins) funnel the energy of photons to the reaction center




 A <u>primary electron acceptor</u> in the reaction center accepts an excited electron from chlorophyll a

 Solar-powered transfer of an electron from a chlorophyll a molecule to the primary electron acceptor is the <u>first</u> <u>step of the light reactions</u>



(a) How a photosystem harvests light

(b) Structure of photosystem II



Two types of photosystems in the thylakoid membrane

 Photosystem II (PS II) functions first (the numbers reflect order of discovery) and is best at absorbing a wavelength of 680 nm

The reaction-center chlorophyll a of PS II is called P680


 Photosystem I (PS I) is best at absorbing a wavelength of <u>700 nm</u>

The reaction-center chlorophyll a of PS I is called P700

Linear Electron Flow

- During the light reactions, there are two possible routes for electron flow:
 - cyclic and
 - linear


Linear electron flow, the <u>primary</u>
 <u>pathway</u>, involves both photosystems
 and produces ATP and NADPH using
 light energy




P680⁺ is a very <u>strong oxidizing</u> agent

 H₂O is <u>Split</u> by enzymes, and the electrons are transferred from the hydrogen atoms to P680⁺, thus reducing it to P680


O₂ is released as a <u>by-product</u> of this reaction




- Each electron "falls" down an electron transport chain from the primary electron acceptor of PS II to PS I
- Energy released by the fall drives the creation of a proton gradient across the thylakoid membrane
- Diffusion of H⁺ (protons) across the membrane <u>drives ATP synthesis</u>




- In PS I (like PS II), transferred light energy excites P700, which loses an electron to an electron acceptor
- P700⁺ (P700 that is missing an electron) accepts an electron passed down from PS II via the electron transport chain



- Each electron "falls" down an electron transport chain from the primary electron acceptor of PS I to the protein ferredoxin (Fd)
- The electrons are then transferred to NADP⁺ and reduce it to NADPH
- The electrons of NADPH are available for the reactions of the Calvin cycle
- This process also removes an H⁺ from the stroma






Cyclic Electron Flow

 Cyclic electron flow <u>uses only</u> <u>photosystem I</u> and <u>produces ATP</u>, <u>but not NADPH</u>


Cyclic electron flow generates
 Surplus ATP, satisfying the higher demand in the Calvin cycle



A Comparison of Chemiosmosis in Chloroplasts and Mitochondria

Chloroplasts and mitochondria generate
 ATP by <u>Chemiosmosis</u>, but use
 different sources of energy


 Mitochondria transfer chemical energy from food to ATP; chloroplasts transform light energy into the chemical energy of ATP



STUDENTS-HUB.com

 ATP and NADPH are produced on the side facing the stroma, where the Calvin cycle takes place

 In summary, light reactions generate ATP and increase the potential energy of electrons by moving them from H₂O to NADPH



Concept 10.3: The Calvin cycle uses ATP and NADPH to convert CO₂ to sugar

• The <u>Calvin cycle</u>, like the <u>citric acid</u> <u>cycle</u>, regenerates its starting material after molecules enter and leave the cycle

 The cycle builds sugar from smaller molecules by using ATP and the reducing power of electrons carried by NADPH Carbon enters the cycle as CO₂ and leaves as a sugar named glyceraldehyde-3-phospate (G3P)

 For net synthesis of <u>1 G3P</u>, the cycle must take place <u>three times</u>, fixing 3 molecules of CO₂

- The Calvin cycle has <u>three phases</u>:
 - Carbon fixation (catalyzed by rubisco)
 - Reduction
 - Regeneration of the CO₂ acceptor (RuBP)

Figure 10.19-1

