

30 tablets

2 mg

**GLIMARYL**<sup>®</sup>  
Glimepiride



Pharmacare plc



**Patient Package Insert**

Prescription Only Medicine

Carefully read the entire leaflet before starting use

**Glimaryl<sup>®</sup> 1, 2, 3, 4 mg Tablets**

**COMPOSITION:** Each tablet contains 1, 2, 3, 4 mg of Glimepiride.

**THERAPEUTIC ACTIVITY:** Oral hypoglycemic agent.

**APPROVED INDICATIONS:** It is indicated for the treatment of non-Insulin dependent diabetes mellitus (TYPE II).

**WHEN SHOULD THE PREPARATION NOT BE USED?**

- Do not use this medicine if you are sensitive to the active ingredient, or to any other component of the drug, or to other sulfonyl urea, or to sulfonamides.
- Do not use this medicine if you have diabetes type 1 or ketoacidosis with or without coma.
- If you are pregnant or breastfeeding.

**DO NOT TAKE THIS MEDICINE WITHOUT CONSULTING A PHYSICIAN:** If you are suffering or have suffered in the past from impaired function of the heart, kidney, liver and / or vascular system. If you suffer from malnutrition, pituitary and adrenal insufficiency.

**WARNINGS:** If you are sensitive to any type of food or medicine, inform your doctor before commencing treatment with this medicine. Blood and urine glucose, should be monitored periodically. Do not drink alcohol while taking glimaryl.

A correct diet, regular & sufficient physical exercise & decrease body weight are important to achieve optimal control of blood sugar.

**Note:** This medicine contains sodium (Na).

**DRUG INTERACTIONS:** If you are taking any additional drugs or if you have just finished a course with another medicine, inform the attending physician, in order to prevent hazards or lack of efficacy arising from drug interactions, especially with medicines belonging to the following: Non-steroid anti-inflammatory drugs, salicylates medications, sulfonamides, chloramphenicol, coumarins, probenecid, Monoamineoxidase inhibitors, thiazides, thyroids products, corticosteroids, propranolol, oral miconazole and oral contraceptives.

**SIDE EFFECTS:** In addition to the desired effect of the medicine, adverse reactions may occur during the course of taking this medicine such as: hypoglycemia, gastrointestinal reactions, allergic skin reactions, blurred vision, headache, nausea, ravenous hunger, lassitude, sleepiness, dizziness, shallow respiration or bradycardia.

In the event that you experience side effects not mentioned in this leaflet, or if there is a change in your