


6.3 Arc Length

Note Title

٢٢/٠١/٢١



DEFINITION If f' is continuous on $[a, b]$, then the length (arc length) of the curve $y = f(x)$ from the point $A = (a, f(a))$ to the point $B = (b, f(b))$ is the value of the integral

$$L = \int_a^b \sqrt{1 + [f'(x)]^2} dx = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx. \quad (3)$$

Remark: A fun with continuous derivative on $[a, b]$ is called smooth, and its curve is called smooth curve on $[a, b]$.


EXAMPLE 1 Find the length of the curve (Figure 6.24)

$$y = \frac{4\sqrt{2}}{3} x^{3/2} - 1, \quad 0 \leq x \leq 1.$$

Sol. $f'(x) = \frac{4}{3} \sqrt{2} \cdot \frac{3}{2} x^{1/2} = 2\sqrt{2} \cdot \sqrt{x}$ which is continuous on $[0, 1]$. So

$$L = \int_0^1 \sqrt{1 + (f')^2} dx = \int_0^1 \sqrt{1 + 8x} dx$$

$$= \frac{1}{8} \int_1^9 \sqrt{u} du = \boxed{\frac{13}{6}}$$



$$u = 1 + 8x$$

$$du = 8 dx$$

$$\frac{du}{8} = dx$$

$$x = 0 \longrightarrow u = 1$$

$$x = 1 \longrightarrow u = 9$$

نلاحظ أننا حصلنا على حلول
المعنى للدالة المطلوبة

EXAMPLE 2

Find the length of the graph of


$$f(x) = \frac{x^3}{12} + \frac{1}{x}, \quad 1 \leq x \leq 4.$$

Sol: $f' = \frac{3x^2}{12} - \frac{1}{x^2}$ which is continuous on $[1, 4]$

Note that $1 + (f')^2 = 1 + \left(\frac{x^2}{4} - \frac{1}{x^2}\right)^2 = 1 + \frac{x^4}{16} - \frac{1}{2} + \frac{1}{x^4}$

$$= \frac{x^4}{16} + \frac{1}{2} + \frac{1}{x^4} = \left(\frac{x^2}{4} + \frac{1}{x^2}\right)^2$$

$$\begin{aligned} \therefore L &= \int_1^4 \sqrt{1 + (f')^2} \, dx = \int_1^4 \sqrt{\left(\frac{x^2}{4} + \frac{1}{x^2}\right)^2} \, dx \\ &= \int_1^4 \left(\frac{x^2}{4} + \frac{1}{x^2}\right) \, dx \quad \left(\frac{x^2}{4} + \frac{1}{x^2} > 0 \quad \forall x\right) \\ &= \left[\frac{x^3}{12} - \frac{1}{x}\right]_1^4 = \left(\frac{64}{12} - \frac{1}{4}\right) - \left(\frac{1}{12} - 1\right) = \frac{72}{12} = \boxed{6} \end{aligned}$$



المختني (لذي أودرنا حوله صوفي يا حمة) ←

ملاحظة: إذا لم تكن المسافة f متصلة على الفترة $[a, b]$ أو كان التكامل باجابه محور x صعب / فإنه قد يكون من المناسب التكامل مع طول المختني يتكامله باجابه y حسب المنهج التالي :

Formula for the Length of $x = g(y)$, $c \leq y \leq d$

If g' is continuous on $[c, d]$, the length of the curve $x = g(y)$ from $A = (g(c), c)$ to $B = (g(d), d)$ is

$$L = \int_c^d \sqrt{1 + \left(\frac{dx}{dy}\right)^2} \, dy = \int_c^d \sqrt{1 + [g'(y)]^2} \, dy. \quad (4)$$

EXAMPLE 3 Find the length of the curve $y = (x/2)^{2/3}$ from $x = 0$ to $x = 2$.

Sol: $y' = \frac{2}{3} \left(\frac{x}{2}\right)^{-\frac{1}{3}} \cdot \frac{1}{2} = \left(\frac{\sqrt[3]{2}}{3}\right) \cdot \frac{1}{\sqrt[3]{x}}$

Note that y' is not continuous at $x=0 \in [0, 2]$, so f is not smooth curve. In this case, we can't use formula (3). So, we try to use the other formula in (4) above as follows:

$$y = \left(\frac{x}{2}\right)^{\frac{2}{3}} \implies x = 2 y^{\frac{3}{2}}.$$

When $x \in [0, 2] \longrightarrow y \in [0, 1]$


($[0, 1]$ is the range of $y = \left(\frac{x}{2}\right)^{\frac{2}{3}}$ when $[0, 2]$ is the domain)

Now $\frac{dx}{dy} = 3\sqrt{y}$ which is continuous on $[0, 1]$

Therefore, using formula (4), we get,

$$L = \int_0^1 \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy = \int_0^1 \sqrt{1 + 9y} dy$$

$$= \frac{2}{27} (10\sqrt{10} - 1) \approx \boxed{2.27}$$



انظر الرسمة

ملاحظة: الرسم ليس شرطاً للحل، وإنما
يعطى للتوضيح.