Ch4 Part 3 Monday, October 25, 2021 2:20 PM

 $\underbrace{E_{XP}}_{\Lambda} \quad f(x) = \frac{x}{x^2 + 1} \quad \text{and} \quad f(x) = \frac{1 - x}{(x^2 + 1)^2} \quad f(x) = \frac{2x(x^2 - 3)}{(x^2 + 1)^3}$ Find (D D(f) = (- v, v) = IR $f(x) = \frac{x}{x^{2}+1} \implies \lim_{x \to \infty} f(x) = 0 \implies y = 0 \text{ is}$ $H \cdot Asy$ 2 Asy. $\lim_{x \to -\infty} f(x) = 0$ $\frac{1}{1+\frac{1}{x^2}} = \frac{0}{1+0} = \frac{0}{1} = 0$ $\lim_{x \to \infty} \frac{x}{x^2 + 1} = \lim_{x \to \infty} \frac{x}{x^2 + 1}$ O A57 x+1 =0 $f(x) = \frac{\chi}{\chi^2 + 1}$ V. 4.57 $|-\chi^2$ => 1-x =0 20 f (x) = 0 $\overline{(\chi^2+1)}$ $X = \pm 1$

STUDENTS-HUB.com

$$x = 1 \quad \in D(f) = 1R$$

$$x = -1 \quad \in D(f) = 1R$$

$$f(x) = \frac{x}{x^{2}+1}$$

$$(1, f(u)) = (1, -\frac{1}{2})$$

$$(-1, f(u)) = (-1, -\frac{1}{2})$$

$$(-1, f(u)) = (-1, -\frac{1}{2})$$

$$(-1, f(u)) = (-1, -\frac{1}{2})$$

$$f = 0 \quad \Rightarrow \quad 1 - x^{2} = 0 \quad \Rightarrow x = \pm 1$$

$$f = 0 \quad \Rightarrow \quad 1 - x^{2} = 0 \quad \Rightarrow x = \pm 1$$

$$f = 0 \quad \Rightarrow \quad 1 - x^{2} = 0 \quad \Rightarrow x = \pm 1$$

$$f = 0 \quad \Rightarrow \quad 1 - x^{2} = 0 \quad \Rightarrow x = \pm 1$$

$$f = 0 \quad \Rightarrow \quad 1 - x^{2} = 0 \quad \Rightarrow x = \pm 1$$

$$f = 0 \quad \Rightarrow \quad 1 - x^{2} = 0 \quad \Rightarrow x = \pm 1$$

$$f = 0 \quad \Rightarrow \quad 1 - x^{2} = 0 \quad \Rightarrow x = \pm 1$$

$$f = 0 \quad \Rightarrow \quad 1 - x^{2} = 0 \quad \Rightarrow x = \pm 1$$

$$f = 0 \quad \Rightarrow \quad 1 - x^{2} = 0 \quad \Rightarrow x = \pm 1$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 3)} = 0 \quad \Rightarrow 2 \times (x^{2} - 3) = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0 \quad \Rightarrow 2 \times (x^{2} - 3) = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0 \quad \Rightarrow 2 \times (x^{2} - 3) = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0 \quad \Rightarrow 2 \times (x^{2} - 3) = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0 \quad \Rightarrow 2 \times (x^{2} - 3) = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0 \quad \Rightarrow 2 \times (x^{2} - 3) = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0 \quad \Rightarrow 2 \times (x^{2} - 3) = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0 \quad \Rightarrow 2 \times (x^{2} - 3) = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0 \quad \Rightarrow 2 \times (x^{2} - 3) = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0 \quad \Rightarrow 2 \times (x^{2} - 3) = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0 \quad \Rightarrow 2 \times (x^{2} - 3) = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0$$

$$f = \frac{2 \times (x^{2} - 3)}{(x^{2} - 4)^{2}} = 0$$

X - - W - ··· f il concerne up on E-V3, 0] U [V3, 0) is concare down on (- 00, - V3] U [0, V3] 6 Inflections Points 2 $f = 0 \Rightarrow x_1 = 0 \in D$ f changes concarity $X_2 = \sqrt{3} \in D$ about X, X, X3 $f(x) = \frac{x}{x^2 + 1}$ ×3= - V3 ED f has forgent at X1, X1, X3 $(\sqrt{3}, f(\sqrt{3})) = (\sqrt{3}, \frac{\sqrt{3}}{4})$ $(-\sqrt{3}, f(-\sqrt{3})) = (-\sqrt{3}, -\frac{\sqrt{3}}{4})$ $(-\sqrt{3}, f(-\sqrt{3})) = (-\sqrt{3}, -\frac{\sqrt{3}}{4})$ ~ (o, f(o)) = (o, o) check at critical Bint 7) EV'S End Doins STUDENTS-HUB.com Uploaded By: Malak Obaid

$$f = - + + + + - - +$$

$$(-1, f(-1)) = (-1, -1) \qquad f \quad has \quad L. Max \quad of \quad \frac{1}{2}$$

$$(-1, f(-1)) = (-1, -1) \qquad f \quad has \quad L. Max \quad of \quad \frac{1}{2}$$

$$(1, f(-1)) = (1, \frac{1}{2}) \qquad f \quad has \quad L. Min \quad of \quad -\frac{1}{2}$$

$$(1, f(-1)) = (1, \frac{1}{2}) \qquad f \quad has \quad L. Min \quad of \quad -\frac{1}{2}$$

$$at \quad x = -1$$


$$(8) \quad Steleh \quad f(x)$$


$$(9) \quad steleh$$


$$(9) \quad stele$$

/


STUDENTS-HUB.com







$$(0, f(0) = (0, 0) \qquad x^{2} + 2x = 0 \qquad x$$



f con [a,b] =)] at least one number f diff (a,b) =) i c c (a,b) sit $\frac{f(b) - f(a)}{b - a}$ vision (f (c) f (مول العالميج ΄ ζ(•) f ((n) $\rightarrow \chi^{\dots}$ C_1 (1 Find the constant c that satisfies MUT for $f(x) = x^2$ on [1,3]Exp f = 2x $f(c) = \frac{f(3) - f(1)}{3 - 1}$ f(c) = 2 C 3 - 1 $\frac{8}{2} = 4$

STUDENTS-HUB.com

