


Elaine N. Marieb  
Katja Hoehn

A photograph of a tennis player, likely Serena Williams, in mid-air performing a backhand shot on a green tennis court. She is wearing a white and yellow athletic outfit and a white headband. Her right arm is extended forward holding a tennis racket, and her left arm is outstretched for balance. The background is a blurred green court surface.

# Human Anatomy & Physiology

SEVENTH EDITION

PowerPoint® Lecture Slides  
prepared by Vince Austin,  
Bluegrass Technical  
and Community College

CHAPTER

# 23


PART A

## The Digestive System

---

# Gastrointestinal Tract (GIT): Overview


- The alimentary canal or gastrointestinal (GI) tract digests and absorbs food
- Alimentary canal – mouth, pharynx, esophagus, stomach, small intestine, and large intestine
- Accessory digestive organs – teeth, tongue, gallbladder, salivary glands, liver, and pancreas



---

# Gastrointestinal Tract Activities

- **Motility**
- **Secretion**
- **Chemical digestion** – catabolic breakdown of food
- **Absorption** – movement of nutrients from the GI tract to the blood or lymph
- **Defecation** – elimination of indigestible solid wastes



Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.



---

# GI Tract

- External environment for the digestive process
- **Regulation of digestion involves:**
  - **Mechanical and chemical stimuli** – stretch receptors, osmolarity, and presence of substrate in the lumen
  - **Extrinsic control** by CNS centers
  - **Intrinsic control** by local centers

---

# Receptors of the GI Tract

- **Mechano- and chemoreceptors** respond to:
  - Stretch, osmolarity, and pH
  - Presence of substrate, and end products of digestion
- They initiate **reflexes** that:
  - Activate or inhibit digestive glands
  - Mix lumen contents and move them along


---

# Peritoneum and Peritoneal Cavity

- **Peritoneum** – serous membrane of the abdominal cavity
  - **Visceral** – covers external surface of most digestive organs
  - **Parietal** – lines the body wall
- **Peritoneal cavity**
  - Lubricates digestive organs
  - Allows them to slide across one another



# Peritoneum and Peritoneal Cavity



**(a) Transverse section of abdominal cavity**

---

# Peritoneum and Peritoneal Cavity


- **Mesentery** – double layer of peritoneum that provides:
  - Vascular and nerve supplies to the viscera
  - Hold digestive organs in place and store fat
- **Retroperitoneal organs** – organs outside the peritoneum
- **Peritoneal organs (intraperitoneal)** – organs surrounded by peritoneum

---

# Histology of the Alimentary Canal

- From esophagus to the anal canal the walls of the **GI tract have the same four tunics**
  - From the lumen outward they are the **mucosa, submucosa, muscularis externa, and serosa**
- Each tunic has a predominant tissue type and a specific digestive function

# Histology of the Alimentary Canal



---

# Mucosa

- Moist epithelial layer that lines the lumen of the alimentary canal
- **Three major functions:**
  - **Secretion of mucus**
  - **Absorption of end products of digestion**
  - **Protection against infectious disease**
- Consists of three layers: a lining epithelium, lamina propria, and muscularis mucosae

---

# Mucosa: Epithelial Lining

- Simple columnar epithelium and mucus-secreting goblet cells
- Mucus secretions:
  - **Protect digestive organs from digesting themselves**
  - **Ease food along the tract**
- Stomach and small intestine mucosa contain:
  - **Enzyme-secreting cells**
  - **Hormone-secreting cells (making them endocrine and digestive organs)**

---

# Mucosa: Other Sublayers

- Submucosa – dense connective tissue containing elastic fibers, blood and lymphatic vessels, lymph nodes, and nerves
- Muscularis externa – responsible for segmentation and peristalsis
- Serosa – the protective visceral peritoneum
  - Replaced by the fibrous adventitia in the esophagus
  - Retroperitoneal organs have both an adventitia and serosa




---

# Enteric Nervous System

- Composed of two major intrinsic nerve plexuses:
  - **Submucosal nerve plexus** – regulates glands and smooth muscle in the mucosa
  - **Myenteric nerve plexus** – Major nerve supply that controls GI tract mobility
- Segmentation and peristalsis are largely automatic involving local reflex arcs
- Linked to the CNS via long autonomic reflex arc

# Anatomy of the Oral Cavity: Mouth



(a)

---

# Palate


- **Hard palate** – underlain by palatine bones and palatine processes of the maxillae
  - Assists the tongue in chewing
  - Slightly corrugated on either side of the raphe (midline ridge)
- **Soft palate** – mobile fold formed mostly of skeletal muscle
  - Closes off the nasopharynx during swallowing
  - Uvula projects downward from its free edge
- Palatoglossal and palatopharyngeal arches form the borders of the fauces

---

# Tongue

- Occupies the floor of the mouth and fills the oral cavity when mouth is closed
- Functions include:
  - **Gripping and repositioning food during chewing**
  - **Mixing food with saliva and forming the bolus**
  - **Initiation of swallowing, and speech**
- Intrinsic muscles change the shape of the tongue
- Extrinsic muscles alter the tongue's position
- Lingual frenulum secures the tongue to the floor of the mouth

# Tongue



---

# Tongue

- Superior surface bears three types of papillae
  - Filiform – give the tongue roughness and provide friction
  - Fungiform – scattered widely over the tongue and give it a reddish hue
  - Circumvallate – V-shaped row in back of tongue


---

# Salivary Glands

- Three pairs of extrinsic glands – **parotid, submandibular, and sublingual**
- Intrinsic salivary glands (**buccal glands**) – scattered throughout the oral mucosa
- Function of Saliva:
  - **Cleanses the mouth**
  - **Moistens and dissolves food chemicals**
  - **Aids in bolus formation**
  - **Contains enzymes that break down starch**



# Salivary Glands



(a)

# Saliva: Source and Composition

- Secreted from serous and mucous cells of salivary glands
- **97-99.5% water, hypo-osmotic, slightly acidic solution containing**
  - **Electrolytes** –  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Cl}^-$ ,  $\text{PO}_4^{2-}$ ,  $\text{HCO}_3^-$
  - **Digestive enzyme** – salivary amylase
  - **Proteins** – mucin, lysozyme, defensins, and IgA
  - **Metabolic wastes** – urea and uric acid

---

# Control of Salivation

- Intrinsic glands keep the mouth moist
- Extrinsic salivary glands secrete serous, enzyme-rich saliva in response to:
  - Ingested food which stimulates chemoreceptors and pressoreceptors
  - The thought of food
- Strong sympathetic stimulation inhibits salivation and results in dry mouth

---

# Pharynx

- From the mouth, the oro- and laryngopharynx allow passage of:
  - **Food and fluids to the esophagus**
  - **Air to the trachea**
- Lined with stratified squamous epithelium and mucus glands
- Has two skeletal muscle layers
  - Inner longitudinal
  - Outer pharyngeal constrictors


---

# Esophagus

- Muscular tube going from the **laryngopharynx to the stomach**
- Travels through the mediastinum and pierces the diaphragm
- Joins the stomach at the **cardiac orifice**



# Esophagus




---

# Digestive Processes in the Mouth

- Food is ingested
- Mechanical digestion begins (chewing)
- Propulsion is initiated by swallowing
- Salivary amylase begins chemical breakdown of starch
- The pharynx and esophagus serve as conduits to pass food from the mouth to the stomach




# Deglutition (Swallowing)




**(a) Upper esophagea  
sphincter contracted**

# Deglutition (Swallowing)




**(b) Upper esophageal  
sphincter relaxed**

# Deglutition (Swallowing)




**(c) Upper esophageal  
sphincter contracted**

# Deglutition (Swallowing)




# Deglutition (Swallowing)



---


# Stomach

- Chemical breakdown of proteins begins and food is converted to chyme
- **Cardiac region** – surrounds the cardiac orifice
- **Fundus** – dome-shaped region beneath the diaphragm
- **Body** – midportion of the stomach
- **Pyloric region** – made up of the antrum and canal which terminates at the pylorus
- The **pylorus** is continuous with the duodenum through the **pyloric sphincter**






**(c) In the stomach, surface area is increased by invaginations called gastric glands.**




Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.



Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.





**1** Food or cephalic reflexes initiate gastric secretion.

**2** Gastrin stimulates acid secretion by direct action on parietal cells or indirectly through histamine.

**3** Acid stimulates short reflex secretion of pepsinogen.

**4** Somatostatin release by  $H^+$  is the negative feedback signal that modulates acid and pepsin release.

#### KEY

- Stimulus
- Receptor
- Integrating center
- Efferent pathway

---

# Regulation of Gastric Secretion

- Neural and hormonal mechanisms regulate the release of gastric juice
- Stimulatory and inhibitory events occur in three phases
  - **Cephalic (reflex) phase:** prior to food entry
  - **Gastric phase:** once food enters the stomach
  - **Intestinal phase:** as partially digested food enters the duodenum

---

# Cephalic Phase

- **Excitatory** events include:
  - Sight or thought of food
  - Stimulation of taste or smell receptors
- **Inhibitory** events include:
  - Loss of appetite or depression
  - Decrease in stimulation of the parasympathetic division

---

# Gastric Phase


- **Excitatory** events include:
  - Stomach distension
  - Activation of stretch receptors (neural activation)
  - Activation of chemoreceptors by peptides, caffeine, and rising pH
  - Release of gastrin to the blood
- **Inhibitory** events include:
  - A pH lower than 2
  - Emotional upset that overrides the parasympathetic division



---

# Intestinal Phase

- **Excitatory phase** – low pH; partially digested food enters the duodenum and encourages gastric gland activity
- **Inhibitory phase** – distension of duodenum, presence of fatty, acidic, or hypertonic chyme, and/or irritants in the duodenum
  - Initiates inhibition of **local reflexes and vagal nuclei**
  - Closes the pyloric sphincter







**1** Food or cephalic reflexes initiate gastric secretion.

**2** Gastrin stimulates acid secretion by direct action on parietal cells or indirectly through histamine.


**3** Acid stimulates short reflex secretion of pepsinogen.

**4** Somatostatin release by  $H^+$  is the negative feedback signal that modulates acid and pepsin release.

#### KEY

-  Stimulus
-  Receptor
-  Integrating center
-  Efferent pathway

# Gastric Contractile Activity




---

# Small Intestine: Gross Anatomy

- Runs from pyloric sphincter to the ileocecal valve
- Has three subdivisions: **duodenum, jejunum, and ileum**
- The bile duct and main pancreatic duct:
  - Join the duodenum at the hepatopancreatic ampulla
  - Are controlled by the sphincter of Oddi
- The jejunum extends from the duodenum to the ileum
- The ileum joins the large intestine at the ileocecal valve

**(a)**



Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.


---

# Small Intestine: Microscopic Anatomy

- Structural modifications of the small intestine wall increase surface area
  - **Plicae circulares**: deep circular folds of the mucosa and submucosa
  - **Villi** – fingerlike extensions of the mucosa
  - **Microvilli** – tiny projections of absorptive mucosal cells' plasma membranes



# Small Intestine: Microscopic Anatomy




---

# Liver

- The largest gland in the body
- Superficially has four lobes – right, left, caudate, and quadrate
- The falciform ligament:
  - Separates the right and left lobes anteriorly
  - Suspends the liver from the diaphragm and anterior abdominal wall


**(a)** The liver is the largest of the internal organs, weighing about 1.5 kg (3.3 lb) in an adult. It lies just under the diaphragm, toward the right side of the body.

**Liver**  
**Gallbladder**



Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.





**(b)** Blood flow to the liver comes from two sources. Oxygenated blood containing metabolites from peripheral tissues reaches the liver via the hepatic artery. Blood to the liver via the hepatic portal vein is rich in absorbed nutrients from the gastrointestinal tract (Fig. 21-30) and contains hemoglobin breakdown products from the spleen. Blood leaves the liver in the hepatic vein (not shown). Bile synthesized in the liver is secreted into the **common hepatic duct** for storage in the gallbladder. From there, it is secreted into the lumen of the intestine through the **common bile duct**.


Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

---

# Liver: Associated Structures

- Bile leaves the liver via:
  - Bile ducts, which fuse into the common hepatic duct
  - The common hepatic duct, which fuses with the cystic duct
    - These two ducts form the bile duct


# Gallbladder and Associated Ducts






# Liver: Microscopic Anatomy

- **Hexagonal-shaped liver lobules** are the structural and functional units of the liver
  - Composed of **hepatocyte** (liver cell) plates radiating outward from a central vein
  - **Portal triads** are found at each of the six corners of each liver lobule



# Microscopic Anatomy of the Liver



---

# Liver: Microscopic Anatomy

- Hepatocytes' functions include:
  - **Production of bile**
  - **Processing bloodborne nutrients**
  - **Storage of fat-soluble vitamins**
  - **Detoxification**
- Secreted bile flows between hepatocytes toward the bile ducts in the portal triads

---

# Composition of Bile

- A yellow-green, alkaline solution containing bile salts, bile pigments, cholesterol, neutral fats, phospholipids, and electrolytes
- Bile salts are cholesterol derivatives that:
  - **Emulsify fat**
  - **Facilitate fat and cholesterol absorption**
  - **Help solubilize cholesterol**
- Enterohepatic circulation recycles bile salts
- The chief bile pigment is **bilirubin**, a waste product of heme

---

# The Gallbladder

- Thin-walled, green muscular sac on the ventral surface of the liver
- **Stores and concentrates bile** by absorbing its water and ions
- **Releases bile** via the cystic duct, which flows into the bile duct

---

# Regulation of Bile Release

- Acidic, fatty chyme causes the duodenum to release:
  - **Cholecystokinin (CCK) and secretin** into the bloodstream
- CCK and secretin transported in blood stimulate the liver to produce bile
- **Vagal stimulation** causes weak contractions of the gallbladder




---

# Regulation of Bile Release

- Cholecystokinin causes:
  - **The gallbladder to contract**
  - **The hepatopancreatic sphincter to relax**
- As a result, bile enters the duodenum

# Regulation of Bile Release




---

# Pancreas

- Location
  - Lies deep to the greater curvature of the stomach
  - The head is encircled by the duodenum and the tail abuts the spleen

# Duodenum and Related Organs




---

# Pancreas

- Location
  - Lies deep to the greater curvature of the stomach
  - The head is encircled by the duodenum and the tail abuts the spleen
- **Exocrine function**
  - Secretes pancreatic juice which breaks down all categories of foodstuff
  - Acini (clusters of secretory cells) contain zymogen granules with digestive enzymes
- The pancreas also has an **endocrine function** – release of insulin and glucagon


# Acinus of the Pancreas



**(a)**



# Pancreatic Activation



---

# Composition and Function of Pancreatic Juice


- Water solution of enzymes and electrolytes (primarily  $\text{HCO}_3^-$ )
  - Neutralizes acid chyme
  - Provides optimal environment for pancreatic enzymes
- Enzymes are released in inactive form and activated in the duodenum

---

# Regulation of Pancreatic Secretion

- **Secretin and CCK** are released when fatty or acidic chyme enters the duodenum
- CCK and secretin enter the bloodstream
- Upon reaching the pancreas:
  - **CCK induces the secretion of enzyme-rich pancreatic juice**
  - **Secretin causes secretion of bicarbonate-rich pancreatic juice**
- Vagal stimulation also causes release of pancreatic juice

# Regulation of Pancreatic Secretion



**TABLE 21-1     The Digestive Hormones**


	STIMULUS FOR RELEASE	PRIMARY TARGET(S)	PRIMARY EFFECT(S)	OTHER INFORMATION
<b>STOMACH</b>				
<b>Gastrin</b>	Peptides and amino acids; neural reflexes	ECL cells and parietal cells	Stimulates gastric acid secretion and mucosal growth.	Somatostatin inhibits release.
<b>INTESTINE</b>				
<b>Cholecystokinin (CCK)</b>	Fatty acids and some amino acids	Gallbladder, pancreas, stomach	Stimulates gallbladder contraction and pancreatic enzyme secretion.  Inhibits gastric emptying and acid secretion.	Promotes satiety.  Some effects may be due to CCK as a neurotransmitter.
<b>Secretin</b>	Acid in small intestine	Pancreas, stomach	Stimulates bicarbonate secretion.  Inhibits gastric emptying and acid secretion.	
<b>Motilin</b>	Fasting: periodic release every 1.5–2 hours	Gastric and intestinal smooth muscle	Stimulates migrating motor complex.	Inhibited by eating a meal.
<b>Gastric inhibitory peptide (GIP)</b>	Glucose, fatty acids, and amino acids in small intestine	Beta cells of pancreas	Stimulates insulin release (feedforward mechanism).  Inhibits gastric emptying and acid secretion.	
<b>Glucagon-like peptide 1 (GLP-1)</b>	Mixed meal that includes carbohydrates or fats in the lumen	Endocrine pancreas	Stimulates insulin release.  Inhibits glucagon release and gastric function.	Promotes satiety.

---

# Motility in the Small Intestine


- The most common motion of the small intestine is segmentation
  - It is initiated by intrinsic pacemaker cells (Cajal cells)
  - Moves contents steadily toward the ileocecal valve
- After nutrients have been absorbed:
  - Peristalsis begins with each wave starting distal to the previous
  - Meal remnants, bacteria, mucosal cells, and debris are moved into the large intestine






Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

**(a) Peristaltic contractions create forward movement.**




**(b) Segmental contractions are responsible for mixing.**




Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

**(a) Peristaltic contractions create forward movement.**



Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

**(b) Segmental contractions are responsible for mixing.**



**No net forward movement**

Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.



---

# Control of Motility

- **Local enteric neurons** of the GI tract coordinate intestinal motility
- **Cholinergic neurons** cause:
  - Contraction and shortening of the circular muscle layer
  - Shortening of longitudinal muscle
  - Distension of the intestine

---

# Control of Motility

- Other impulses relax the circular muscle
- **The gastroileal reflex and gastrin:**
  - Relax the ileocecal sphincter
  - Allow chyme to pass into the large intestine




---

# Large Intestine

- Has three unique features:
  - Teniae coli – three bands of longitudinal smooth muscle in its muscularis
  - Haustra – pocketlike sacs caused by the tone of the teniae coli
  - Epiploic appendages – fat-filled pouches of visceral peritoneum

# Large Intestine




---

# Valves and Sphincters of the Rectum and Anus

- Three valves of the rectum stop feces from being passed with gas
- The anus has two sphincters:
  - Internal anal sphincter composed of smooth muscle
  - External anal sphincter composed of skeletal muscle
- These sphincters are closed except during defecation



# Structure of the Anal Canal



(b)

---

# Bacterial Flora

- The bacterial flora of the large intestine consist of:
  - Bacteria surviving the small intestine that enter the cecum and
  - Those entering via the anus
- These bacteria:
  - Colonize the colon
  - Ferment indigestible carbohydrates
  - Release irritating acids and gases (flatus)
  - Synthesize B complex vitamins and vitamin K

---

# Functions of the Large Intestine

- Other than digestion of enteric bacteria, no further digestion takes place
- Vitamins, water, and electrolytes are reclaimed
- Its major function is propulsion of fecal material toward the anus
- Though essential for comfort, the colon is not essential for life



---

# Motility of the Large Intestine


- Haustral contractions
  - Slow segmenting movements that move the contents of the colon
  - Haustra sequentially contract as they are stimulated by distension
- Presence of food in the stomach:
  - Activates the gastrocolic reflex
  - Initiates peristalsis that forces contents toward the rectum

---

# Defecation

- Distension of rectal walls caused by feces:
  - Stimulates contraction of the rectal walls
  - Relaxes the internal anal sphincter
- Voluntary signals stimulate relaxation of the external anal sphincter and defecation occurs

# Defecation



---

# Chemical Digestion: Carbohydrates


- Absorption: via cotransport with  $\text{Na}^+$ , and facilitated diffusion
  - Enter the capillary bed in the villi
  - Transported to the liver via the hepatic portal vein
- Enzymes used: salivary amylase, pancreatic amylase, and brush border enzymes

# Chemical Digestion: Proteins

- Absorption: similar to carbohydrates
- Enzymes used: pepsin in the stomach
- Enzymes acting in the small intestine
  - Pancreatic enzymes – trypsin, chymotrypsin, and carboxypeptidase
  - Brush border enzymes – aminopeptidases, carboxypeptidases, and dipeptidases



***InterActive Physiology®:***  
**Digestion and Absorption, pages 5 and 8**






---

# Chemical Digestion: Fats

- Absorption: Diffusion into intestinal cells where they:
  - Combine with proteins and extrude chylomicrons
  - Enter lacteals and are transported to systemic circulation via lymph
- Glycerol and short chain fatty acids are:
  - Absorbed into the capillary blood in villi
  - Transported via the hepatic portal vein
- Enzymes/chemicals used: bile salts and pancreatic lipase

# Chemical Digestion: Fats




---

# Fatty Acid Absorption

- Fatty acids and monoglycerides enter intestinal cells via diffusion
- They are combined with proteins within the cells
- Resulting chylomicrons are extruded
- They enter lacteals and are transported to the circulation via lymph

# Fatty Acid Absorption



---

# Chemical Digestion: Nucleic Acids

- Absorption: active transport via membrane carriers
- Absorbed in villi and transported to liver via hepatic portal vein
- Enzymes used: pancreatic ribonucleases and deoxyribonuclease in the small intestines

---

# Electrolyte Absorption

- Most ions are actively absorbed along the length of small intestine
  - $\text{Na}^+$  is coupled with absorption of glucose and amino acids
  - Ionic iron is transported into mucosal cells where it binds to ferritin
- Anions passively follow the electrical potential established by  $\text{Na}^+$



---

# Electrolyte Absorption

- $K^+$  diffuses across the intestinal mucosa in response to osmotic gradients
- $Ca^{2+}$  absorption:
  - Is related to blood levels of ionic calcium
  - Is regulated by vitamin D and parathyroid hormone (PTH)

---

# Water Absorption

- 95% of water is absorbed in the small intestines by osmosis
- Water moves in both directions across intestinal mucosa
- Net osmosis occurs whenever a concentration gradient is established by active transport of solutes into the mucosal cells
- Water uptake is coupled with solute uptake, and as water moves into mucosal cells, substances follow along their concentration gradients

---

# Malabsorption of Nutrients

- Results from anything that interferes with delivery of bile or pancreatic juice
- Factors that damage the intestinal mucosa (e.g., bacterial infection)
- Gluten enteropathy (adult celiac disease) – gluten damages the intestinal villi and reduces the length of microvilli
  - Treated by eliminating gluten from the diet (all grains but rice and corn)

---

# Cancer

- Stomach and colon cancers rarely have early signs or symptoms
- Metastasized colon cancers frequently cause secondary liver cancer
- Prevention is by regular dental and medical examinations

---

# Cancer

- Colon cancer is the 2nd largest cause of cancer deaths in males (lung cancer is 1st)
- Forms from benign mucosal tumors called polyps whose formation increases with age
- Regular colon examination should be done for all those over 50