Mathematics Department Math 1411 - Worksheet #7 Rasha shadid Q. Are the factions  $f(x) = (X+1)^3+2$  and  $g(x) = X+\frac{1}{X}$ One-to-one? Q2 let f(x)= 8x3+3. Show that (i) f' exist (ii) Show that  $f'(x) = \frac{1}{2} \sqrt[3]{x-3}$ (iii) find df at x=2 On if guis the inverse function of fix P(4) = 5 / f(4) = = = - find g'(5) Qu Find y' for each of the following: - $2 y = ln \left( \frac{1}{3+x^2} \right)$ O y = x Vln x 3 y = J lu + dt Q5 Fint the following integrals:

STUDENTS-HUB; com3

$$\frac{Q_1}{g(x)} \quad \text{is } 1-1$$

$$g(x) \quad \text{is } not \quad L1 \quad \left(\text{Since} \quad g(2) - g(\frac{1}{2})\right)$$

(ii) 
$$f^{-1}(f(x)) = f^{-1}(8x^3 + 3) = X$$
  
And  $f(f^{-1}(x)) = f(\frac{1}{2}\sqrt[3]{x-3}) = X$ 

And 
$$F(F(X)) = F(\frac{1}{2})^n$$
  
Thun  $\frac{1}{2}\sqrt[3]{X-3}$  is the inverse function of  $8x^3+3$ 

(iii) 
$$\left(\frac{df^{-1}}{dx}\right)$$
 =  $\frac{1}{6}$ 

$$Q_3$$
  $Q'(5) = \frac{3}{2}$ 

Q4 (1) 
$$y' = \frac{1 + \ln x}{2 \sqrt{\ln x}}$$
 (2)  $y' = \frac{1}{2x+2} - \frac{1}{3+x^2}(2x)$ 

(3) 
$$y' = 3 x^2 ln x^3 - 2x ln x^2$$

$$Q_5$$
.  $Q_5$   $\frac{1}{8} \ln |2x^4+3| + C$ 

Mathematics Department

Math 141 - Worksheet #8

Ch. 7.3 - 7.5

Rasha Shadid

Name:\_\_\_\_

Q Simplify the following expression:-

(3) e + lm √e6

6) log 27 + log 18 - ln e

Q2: Find the derivative of the following

 $0 \quad y = 3 \quad \log_8 \left( \log_2 x \right)$ 

2) y= Jen + d+

P3 Evalute the following integral:

 $\int \frac{e^{-1/x^2}}{x^3} dx$ 

STUDENTS HUB Com

Uploaded By: anonymous

By: Find the limit of the following: Rim Sin x - X 3  $\lim_{X \to (\frac{T}{2})^{-}} \left(\frac{T}{2} - X\right) \tan X$ 

 $\lim_{x \to 0} \left( e^x + x \right)^{1/x}$   $\lim_{x \to 0} \left( e^x + x \right)^{1/x}$   $\lim_{x \to 0} \left( e^x + x \right)^{1/x}$ 

(5) lim (x3+e) = 20 x

6 lim Sinx lux x-sot Short Answers:-

$$Q_2 \qquad 0 \qquad y' = \frac{1}{\ln 2} \left( \frac{1}{x \ln x} \right)$$

$$\int_{0}^{\infty} \frac{-1/x^{2}}{1 - 1/2} e^{-1/x^{2}} + C$$

① 
$$I = e^{x} + \ln |e^{x} - 1| + C$$
  
②  $I = e^{x} + \ln |e^{x} - 1| + C_{1}$ , since  $C_{1} = c^{-1}$   
 $= e^{x} + \ln |e^{x} - 1| + C_{1}$ , since  $C_{1} = c^{-1}$ 

$$(3) \quad | \quad \qquad (4) \quad e$$

$$(5) e^{3}$$
  $(6) C$ 

STUDENTS-HUB.com

Uploaded By: anonymous

lim Sinx lnx O(-00) lim CSC X lm x lim - SMX tanx OR l'Hopital
Rule -lim sinx lim tanx x sot

Mathematics Department Math 1411 - Worksheet #9 "7.6, 7.7, 7.8" . Rasha. Shadid

· Name: - - - -

[Q]: Find the exact value of the following expression if its exist

a)  $Sin\left(\cos^{-1}\left(\frac{\sqrt{2}}{2}\right)\right)$ 

B Sin' (Sin 71)

Cot  $\left(Sin'\left(\frac{1}{2}\right) - Sec'(2)\right)$ 

Q2 Find the solution of the following equation  $ln(x-e) = x^2 - tan^2(sec^Tx)$ 

Q3 Evalute the following integral

(a)  $\int \frac{\sec^2 x}{\sqrt{4 - \tan^2 x}}$ 

STUDENTS-HUB.com

Uploaded By: anonymous

Qu Simplify the following expression
Sinh (2 lnx)

(D5) Evalute the following integral

a) Je-x cash x dx

(b) Sech x Seeh x tanh x dx

Discuss the growth of following Palr

X<sup>2</sup> and VYXY+3X<sup>2</sup>+1

$$Q_1$$
  $Q_2$   $\frac{1}{\sqrt{2}}$ 

$$\bigcirc$$
  $\frac{}{3}$ 

$$Q_2$$
  $X = 2e$ 

$$\int_{3}^{2} = \sin^{-1}\left(\frac{\tan x}{2}\right) + C$$

(b) 
$$T = \frac{1}{4} \ln |x^{4}+9| + \frac{1}{2} \tan^{-1}(\frac{x^{2}}{3}) + C$$

$$Q_{4}$$
  $\frac{\chi^{4}-1}{2\chi}$ 

$$\Box = \frac{1}{2} \times - \frac{1}{4} e^{-2x} + C$$

STUDENTS-HUBK.com 
$$\sqrt{\frac{x^2}{4x^4+3x^2+1}} = \frac{1}{2}$$