


Flow Diagram for the separation of cations into qualitative Groups.



[HCl]

$\text{AgCl}, \text{PbCl}_2, \text{Hg}_2\text{Cl}_2$
 Group I

Group II, III, IV

[HCl, H_2S]

Group II

$\text{Bi}_2\text{S}_3, \text{CuS}, \text{As}_2\text{S}_3, \text{Sb}_2\text{S}_3$

Group III, IV

[$\text{NH}_3, \text{NH}_4^+, \text{H}_2\text{S}$]

Group III

$\text{NiS}, \text{FeS}, \text{Al}(\text{OH})_3, \text{ZnS}$

Group IV

$\text{Na}^+, \text{NH}_4^+, \text{Ba}^{2+}, \text{Ca}^{2+}$

در جدول
 علی جدول

① precipitation → separation ②


Prelaboratory Assignment

Group I. The Insoluble Chlorides (Ag^+ , Pb^{2+} , Hg_2^{2+})

Group I. The Insoluble Chlorides


Date _____ lab sec. _____ Name _____

ليسوا راسب أبيض.



من فلان الـ HCl بفضل الحروب الأول من الثاني عن الثالث. $\text{[HCl]} \rightarrow$ group reagent


Group I



$[\text{H}_2\text{O}, \text{heat}]$


$\text{ppt} \leftarrow$

هو اوي الذي
يزدب NH_3




نظام
راسب.

$[\text{NH}_3]$



Black

$[\text{HNO}_3]$ مع القاذبة



white


X

known :- (كلمة هوسونين) فيه

1- 2 drops HCl على البسوخنة
بمسوخة راسب أبيض.

2- 10 drops H_2O
بمسوخة بـ
bath


$\rightarrow$ supernatant.



هول تاندا
heat. بال

يزدب بال
suber
notant.







$[\text{K}_2\text{CrO}_4]$




yellow

اذا هارندي راسب
أمنر بيني تغلي برني طم.

G.II (flow diagram)








total charge = 0






Subgroup II B

Procedure
E, F, G




white








Black

X








deep blue

suber natant.



dark brown

yellow



Orange

X



X

Sub group II A

procedure

C, D


لا ترسبني وسط محلول لونها البنية فيه



ترسبني وسط

عني

excess to force the forward reaction $\rightarrow$



For calibration curve

$[\text{SCN}^{-}] = [\text{FeSCN}^{+2}] \rightarrow$ because $[\text{Fe}^{+3}]$ is in large excess compared to SCN^{-}

$A = \epsilon C$

For equilibrium concentrations

$C = \frac{A}{\epsilon}$ or $\frac{A}{\text{slope}}$

C in Molarity = $\frac{\# \text{ of moles}}{\text{Volume (L)}}$

$\# \text{ of moles} = M \times V$

M of FeSCN^{+2} for calibration curve = M of SCN^{-}

Example ① volume
mls of $\text{NaSCN} = 2 \text{ ml}$
M of $\text{SCN}^{-} = 0.002 \text{ M}$

$\# \text{ of moles} = M \times V = 0.002 \times \frac{2}{1000} = \frac{4}{1000 \times 1000} = 4 \times 10^{-6} \text{ moles}$

M of $\text{SCN}^{-} = \frac{\# \text{ of moles}}{\text{total V (L)}} = \frac{4 \times 10^{-6}}{\frac{100}{1000}} = \frac{4 \times 10^{-6} \times 1000}{100} = 4 \times 10^{-5} \text{ M}$

Example ②

$\# \text{ of moles} = M \times V = 0.002 \times \frac{4}{1000} = 8 \times 10^{-6} \text{ moles}$

$M = \frac{8 \times 10^{-6}}{\frac{100}{1000}} = 8 \times 10^{-5} \text{ M}$

Example ③

$\frac{12 \times 10^{-6}}{0.1} = 12 \times 10^{-5} \text{ M}$

Example ④

$\frac{16 \times 10^{-6}}{0.1} = 16 \times 10^{-5} \text{ M}$


①

shahd Abu Rajab



Cosmic rays

الأشعة طول موجي
في الأتوم
والفلك مجي

X-rays violet
UV-light indigo
Visible blue
yellow
red orange




$\text{slope} = \frac{A_2 - A_1}{\frac{[\text{FeSCN}^{+2}]_2 - [\text{FeSCN}^{+2}]_1}{2}}$



(2)

For equilibrium concentrations to determine $K_{eq} = K_c$
 First we should determine the initial concentrations of $Fe^{+3} + SCN^-$



② $M_{initial} = \frac{10 \times 10^{-6}}{\frac{10}{1000}} = \frac{10 \times 10^{-6} \times 1000}{10} = 1 \times 10^{-3}$

$[Fe^{+3}]_{initial}$

$M = \frac{\# \text{ of moles}}{\text{Volume in liter}}$

$[SCN^-]_{initial} = \frac{4 \times 10^{-6}}{\frac{10}{1000}} = \frac{4 \times 10^{-6} \times 1000}{10} = 4 \times 10^{-4}$

Example ② Fe^{+3}

of moles = $0.002 M \times \frac{5}{1000} = 10 \times 10^{-6} = 1 \times 10^{-5}$

$[Fe^{+3}]_{initial} = \frac{10 \times 10^{-6}}{\frac{10}{1000}} = 1 \times 10^{-3}$

$[SCN^-]_{initial} = \frac{8 \times 10^{-6}}{\frac{10}{1000}} = 8 \times 10^{-4} M$

③ $\frac{12 \times 10^{-6}}{\frac{10}{1000}} = 12 \times 10^{-4}$

④ $\frac{16 \times 10^{-6}}{\frac{10}{1000}} = 16 \times 10^{-4}$

3

Spectrophotometric determination of an Equilibrium Constant

Date _____ lab sec. _____ Name _____

DATA

A. Set of Standard Solutions

1. Molar concentration $\text{Fe}(\text{NO}_3)_3$ _____2. Molar concentration NaSCN _____

Standard Solutions	1	2	3	4	5
Volume NaSCN (mL)	0	2	4	6	8
Moles SCN^- ($M \times V$) $\frac{\text{moles}}{V(L)}$	0	4×10^{-6}	8×10^{-6}	12×10^{-6}	16×10^{-6}
$[\text{SCN}^-]$ IN 100 mL solution	0	4×10^{-5}	8×10^{-5}	12×10^{-5}	16×10^{-5}
$[\text{FeSCN}^{2+}]$ $\frac{\text{total volume}}$	0	4×10^{-5}	8×10^{-5}	12×10^{-5}	16×10^{-5}
Absorbance, A					

Construct a standardization curve for A Vs $[\text{FeSCN}^{2+}]$

B. Set of Equilibrium Solutions

1. Molar concentration $\text{Fe}(\text{NO}_3)_3$ 0.002 M2. Molar concentration NaSCN 0.002 M

3. Solutions "

Solutions number	1	2	3	4
a. Volume $\text{Fe}(\text{NO}_3)_3$ (mL)	5	5	5	5
b. Moles Fe^{3+} , initial	$10 \times 10^{-6} = 1 \times 10^{-5}$	1×10^{-5}	1×10^{-5}	1×10^{-5}
c. Volume NaSCN (mL)	0	2	3	4
d. Moles SCN^- , initial	0	4×10^{-6}	6×10^{-6}	8×10^{-6}
e. Absorbance				

total volume 10 mL

example to determine # of initial moles of SCN^-
 For SCN^- initial = # of moles = $M \times V = 0.002 \times \frac{2}{1000} = 4 \times 10^{-6}$
 example take 2 mL

of moles = $M \times V$

Example ~~took~~ from calibration curve

1. $[FeSCN^{+2}]_{eq} = 8 \times 10^{-5} M$ = suppose its like that

2. $[Fe^{+3}]_{initial} = 1 \times 10^{-5}$

of moles of $FeSCN^{+2}$ in solution at eq = $M \times V = 8 \times 10^{-5} \times \frac{10}{1000} = 8 \times 10^{-7}$

2a. moles of Fe^{+3} complexed = moles of $FeSCN^{+2}$ at eq = 8×10^{-7}

2b. moles of Fe^{+3} uncomplexed = initial moles - complexed

2c. moles of Fe^{+3} at equilibrium

$$= 1 \times 10^{-5} - 8 \times 10^{-7} = 1 \times 10^{-5} - 0.08 \times 10^{-5}$$

$$= 0.92 \times 10^{-5}$$

2d. $[Fe^{+3}]_{eq} = \frac{\# \text{ of moles}}{V(L)} = \frac{0.92 \times 10^{-5}}{0.01} = 0.92 \times 10^{-3}$

$0.01 = 10/1000$ total volume = 10 ml

3. $[SCN^{-}]$

3a. moles of SCN^{-} complexed = moles of $FeSCN^{+2}$ at equilibrium = $8 \times 10^{-7} \text{ mol}$

3b. moles of SCN^{-} uncomplexed = initial moles - complexed

$$= 4 \times 10^{-6} - 8 \times 10^{-7}$$

$$= 4 \times 10^{-6} - 0.8 \times 10^{-6}$$

$$= 3.2 \times 10^{-6}$$

3c. $[SCN^{-}]_{eq} = \frac{\# \text{ of moles}}{V(L)} = \frac{3.2 \times 10^{-6}}{0.01} = 3.2 \times 10^{-4}$

$\uparrow$ = at complex.

4. $K_c = \frac{[FeSCN^{+2}]_{eq}}{[Fe^{+3}]_{eq} [SCN^{-}]_{eq}} = \frac{8 \times 10^{-5}}{(0.92 \times 10^{-3})(3.2 \times 10^{-4})}$

→ From calibration curve

$$= \frac{8 \times 10^{-5}}{2.94 \times 10^{-7}} = \frac{8 \times 10^{-6}}{2.94 \times 10^{-7}} = \frac{27.16}{1 \times 10^{-1}} = 27.16 \times 10$$

$$= 2.7 \times 10^2$$

literature value = 138 at 25°C

≈ 270

5

CALCULATIONS:


Determination of K_c

	1	2	3	4
1. $[\text{FeSCN}^{2+}]$, from calibration curve	from calibration curve			
2. $[\text{Fe}^{3+}]_{\text{initial}}$	1×10^{-3}	1×10^{-3}	1×10^{-3}	1×10^{-3}
a. Moles FeSCN^{2+} in solution at equilibrium				
b. Moles Fe^{3+} complexed				
c. Moles Fe^{3+} uncomplexed	initial - complexed			
d. $[\text{Fe}^{3+}]$ equilibrium, uncomplexed (mole/liter)				
3. $[\text{SCN}^-]_{\text{initial}}$				
a. Moles SCN^- complexed = moles of FeSCN^{2+} in equlb				
b. Moles SCN^- uncomplexed				
c. $[\text{SCN}^-]$ at equilibrium, uncomplexed (mole/liter)				
4. $K_c = \frac{[\text{FeSCN}^{2+}]}{[\text{Fe}^{3+}][\text{SCN}^-]}$				

5. Average $K_c = \frac{K_{c1} + K_{c2} + K_{c3} + K_{c4}}{4}$


6. Average deviation. Sum all of the absolute differences between each K_c and the average K_c and divide by 4 (the number of measurements).

Determination of energy of activation



The rxn is 2nd order (1st order in each reactant)



The rxn occurs in several steps with the first step is:



when the I^- is present in large excess the rxn is pseudo first order
0.5M its conc remains const

$$\ln \left(\frac{[A]_t}{[A]_0} \right) = -kt$$

$$\text{or } t = -\frac{\ln \left(\frac{[A]_t}{[A]_0} \right)}{k}$$



when $[A]_t = \frac{1}{2} [A]_0 \rightarrow t = t_{1/2}$



$$k = \ln \frac{n}{t_{1/2}} = A e^{\left(\frac{-E_a}{RT} \right)} \rightarrow \text{Arrhenius equation}$$

$$R = 8.314 \text{ J mol}^{-1} \text{ K}^{-1}$$

T in Kelvin


$$t_{1/2} = \frac{\ln(n)}{A} e^{\left(\frac{E_a}{RT} \right)}$$

$$\ln t_{1/2} = \ln \left[\frac{\ln(n)}{A} \right] + \frac{E_a}{RT}$$




$S_2O_3^{2-}$
0.01 M

10 ml




$S_2O_8^{2-}$
0.01 M

20 ml



KI
0.5 M

20 ml



$$K_{eq} = \frac{[C][D]}{[A][B]}$$

تساوی در حالت تعادل


Exp # 1

Le Chatelier's Principle

- 1- Temperature effect → change in equilibrium constant
- 2- Concentration effect


A. Saturated NaCl Solution

concentration effect




add HCl(aq) shift the equilibrium to the left because it increase the Cl^- ion

کاهش غلظت Cl^- باعث افزایش $NaCl$ می شود
تفاعل هو ایستاد و متوازن است



B. $SbCl_3 - H_2O$ Equilibrium

concentration effect




add HCl the $SbOCl(s)$ will dissolve the equilibrium shift to the left

add H_2O again the equilibrium shifts to the right and $SbOCl(s)$ will ppt

solid
H₂O
پرسو
پرسو
پرسو
Cl

C. $CrO_4^{2-} - Cr_2O_7^{2-}$ Equilibrium

concentration effect









Yellow

Orange

add H^+ from HNO_3
it will react with OH^-
and shift the eqn to the left.


D. Equilibrium involving silver ion Ag^+

① املاح منوع




Free Ag^+

لوصية Cu بدل Ag بطلع عنا لون أزرق سمادي.






بطلع عنا لون

↓ add 6M HNO_3 this will react with NH_3



$AgI \rightarrow$ yellow

1.9 test tube.





أول انما نفضل
درجف اللون
تتوصل لاسك

اللون الزماني
green
blue


(2) أبيض بالعادة

1g KBr



(3) صمغ ركب
بنتقوده
بج و


save this solution



continue adding H_2O until 15 ml is added




save this solution for Part G

كيف بررم سدادله لواء



شيف اللوناب

common ion effect



temp

the rxn is endothermic: $\Delta H +$
So when heating the solution the rxn will shift to the right and the color change from blue to green.

*Solution 1: Blank 1M $CoCl_2$ only
*Solution 2: place solution 1 in a boiling water bath

*Solution 3 & 4: Saturate both with $NaCl$ + compare the color with solution 1
*Heat solution 3 to examine temp eff compare with solution 1, explain the color

*Decant ~ 2 ml of solution 4 into small test tube, add 5-10 drops of 1M $AgNO_3$ take the supernatant liquid and explain your observation

برجى يلكى راب انين
عن سدادله لواء