













	Mean diameter	Mean wall thickness	Endothelium Elastic tissue Smooth muscle Fibrous tissue	
Artery	4.0 mm	1.0 mm		
Arteriole	30.0 μm	6.0 μm		
Capillary	8.0 μm	0.5 μm		
Venule	20.0 μm	1.0 μm		
Vein	5.0 mm	0.5 mm		

Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.




Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

(a) Ventricular contraction




- 1 Ventricle contracts.
- 2 Semilunar valve opens.
- 3 Aorta and arteries expand and store pressure in elastic walls.

(b) Ventricular relaxation




- 1 Isovolumic ventricular relaxation occurs.
- 2 Semilunar valve shuts, preventing flow back into ventricle.
- 3 Elastic recoil of arteries sends blood forward into rest of circulatory system.




Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

Valves in the veins prevent backflow of blood.





Valve open

Valve closed




When the skeletal muscles compress the veins, they force blood toward the heart (the skeletal muscle pump).




MEAN ARTERIAL BLOOD PRESSURE

is determined by




Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

(a) Active hyperemia




Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.


(b) Reactive hyperemia



Arteriole diameter is controlled by tonic release of norepinephrine.




Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.




Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

(a) Blood flow through four identical vessels (A–D) is equal. Total flow into vessels equals total flow out.




(b) When vessel B constricts, resistance of B increases and flow through B decreases. Flow diverted from B is divided among the lower-resistance vessels A, C, and D.




Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings

(a) When precapillary sphincters are relaxed, blood flows through all capillaries in the bed.





(b) If precapillary sphincters constrict, blood flow bypasses capillaries completely and flows through metarterioles.






Faster


(a) Continuous capillaries have leaky junctions.



(b) Fenestrated capillaries have large pores.




Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.



Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

(a) Filtration in systemic capillaries



Hydrostatic pressure P_{cap} forces fluid out of the capillary.

↳ let fluids out


Colloid osmotic pressure of proteins within the capillary pulls fluid into the capillary.

KEY

P_{cap} = Capillary hydrostatic pressure

π = Colloid osmotic pressure


(b) Relationship between capillaries and lymph vessels



The excess water and solutes that filter out of the capillary are picked up by the lymph vessels and returned to the circulation.

(a) Filtration in systemic capillaries

Net pressure = hydrostatic pressure – colloid osmotic pressure




KEY

P_{cap} = Capillary hydrostatic pressure

π = Colloid osmotic pressure


Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

(b) Relationship between capillaries and lymph vessels







The excess water and solutes that filter out of the capillary are picked up by the lymph vessels and returned to the circulation.

Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.



Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.






When Blood P increase → Baroreceptors


Higher firing

Cardiovascular control center


Symp in higher Blood P
 Sy ↓ Para ↑

Para






Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.




Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.



Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

(c) Stable fibrous plaque



A lipid core accumulates beneath the endothelium.


Fibrous scar tissue cap forms to wall off the lipid core.

Smooth muscle cells divide and contribute to thickening of the intima.

Calcifications are deposited within the plaque.

Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

(d) Vulnerable plaque



Platelets that are exposed to collagen activate and initiate a blood clot.

Macrophages may release enzymes that dissolve collagen and convert stable plaques to unstable plaques.

Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.