


Taylor Theorem :-

$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n + \dots$$



$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n$$

$$f(x) \approx f(a) + f'(a)(x-a) \quad \text{linear estimation}$$

$$\text{Error} = \frac{f''(a)}{2!}(x-a)^2 + \dots$$

$$f(x) \approx f(a) + f'(a)(x-a) + \frac{f''(a)}{2}(x-a)^2$$

$$\text{Error} = \frac{f'''(a)}{3!}(x-a)^3 + \dots$$

in general

$$f(x) \approx f(a) + f'(a)(x-a) + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n$$

$$\text{Error} = \frac{f^{(n+1)}(a)}{(n+1)!}(x-a)^{n+1} + \dots \quad (\text{infinite Terms}).$$

Taylor :-

$$\text{Error} = \frac{f(c)(x-a)^{n+1}}{(n+1)!} \quad c \text{ between } a, x$$

$$|\text{Error}| \leq \max_{a \leq x \leq c} \left| \frac{f^{(n+1)}(c)}{(n+1)!} (x-a)^{n+1} \right|$$

↑ Error up
c is less also ↑ error

$P_n(x)$

$$\Rightarrow f(x) \approx f(a) + f'(a)(x-a) + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n + \frac{f^{(n+1)}(c)}{(n+1)!}(x-a)^{n+1}$$

↓ Error
 $E_n(x)$

$$e^x, \quad a=0$$


$$e^x = f(0) + f'(0)(x-0) + \frac{f''(c)}{2!}(x-0)^2$$

$$e^x = 1 + x + \frac{e^c x^2}{2!}$$

$$e^x \approx 1+x \text{ with error } \frac{e^c x^2}{2!}$$

$$e^x = f(0) + f'(0)(x-0) + \frac{f''(0)(x-0)^2}{2!} + \frac{f'''(c)(x-0)^3}{3!}$$

$$e^x \approx 1 + x + \frac{x^2}{2} \quad \text{error} = \frac{e^c x^3}{6}$$




$$\begin{aligned} e^x &\approx 1 + 0.1 + \frac{0.01}{2} & \text{error } \frac{e^c (0.001)}{6} < 1 \times 10^{-3} \\ &\approx 1.105 & c \in [0, 0.1] \end{aligned}$$

Upper bound for error $\frac{e^c (0.001)}{6} \leq \frac{e^{0.001}}{6} \leq \frac{1.0005}{6} \leq 0.0005$


Intermediate Value Theorem (IVT)

- $f(x)$ is continuous
- L between $f(a)$ and $f(b)$
- Then $\exists c \in (a, b)$ such that $f(c) = L$




bolzano

- $f(x)$ is continuous
- $f(a) = f(b) < 0$
- Then $\exists c \in (a, b)$ such that $f(c) = 0$



mean value theorem (MVT)

- $f(x)$ is continuous on $[a, b]$
- $f(x)$ is differentiable on (a, b)
- then $\exists c \in (a, b)$ such that $f'(c) = \frac{f(b) - f(a)}{b - a}$



sis

suppose that p^n is an approximation to P

the error is $E_p = P - p^n$

the relative error $R_p = \frac{E_p}{P} = \frac{P - p^n}{P}$

X:- 1. let $x = 3.141592$

$$x^n = 3.14$$

متوازن متقاربة میں

$$Ex = 3.141592 - 3.14 = 0.001592$$

$$Rx = \frac{0.001592}{3.141592} = 0.000507$$

2. let $y = 1,000,000$

$$y^n = 999,996 \quad \begin{array}{l} \text{متوازن متقاربة} \\ \text{میں اکٹھے میں کی} \end{array}$$

$$Ey = 4$$

$$Ry = \frac{4}{1,000,000} = 4 * 10^{-6}$$

3. let $z = 0.000,012$

$$z^n = 0.000,009$$

متوازن متقاربة
و لا فنا ای فنزلہ

$$Ez = 0.000,003$$

$$Rz = 0.25$$

Normalized decimal Form:-

$$\pm 0.d_1d_2d_3\ldots \times 10^n$$

$$d_1 \neq 0$$

$$x^2 = 2$$

$$x^2 - 2 = 0 \quad \begin{array}{c} - + + + \\ \hline 1.5 \end{array} \quad \begin{array}{l} \text{جواب} \\ \text{نہیں} \end{array}$$

$$0 = \frac{1+2}{2} = 1.5$$

$$1 = \frac{1+1.5}{2} = 1.25$$

$$2 = \frac{1+1.25}{2} = 1.125 = 0.1125 * 10^1$$

2 significant digits \Rightarrow Error $\leq 10^{-2}$

بعد اول مترال بعده

Def: the number \hat{P} is said to approximate P to d significant digits if d is the largest positive integer for which

$$\frac{|P - \hat{P}|}{|P|} < \frac{10^{-d}}{2}$$

$$\text{i.e } 2|R_{\text{rel}}| \leq 10^{-d}$$

ex:-

$$1. X = 3.141592$$

$$x^{\text{chop}} = 3.14$$

$$R_x = 3.141592 - 3.14 = 0.001592$$

$$R_x = \frac{0.001592}{3.141592} = 0.000507$$

$$2|R_x| = 0.001014 \approx 10^{-3} \\ < 10^{-4}$$

example 2. $2|R_y| = 8 \times 10^{-6} < 10^{-3}$

$$\begin{array}{c} 10^{-2} \\ 10^{-3} \\ 10^{-4} \\ 10^{-5} \\ \boxed{10^{-5}} \\ 10^{-6} \end{array}$$

3. $2|R_z| = 0.5 \not< 10^{-4}$
no significant bits.

- if $P = \pm 0.d_1 d_2 \dots d_n d_{n+1} \dots \times 10^n$ is the normalized decimal form of the number P , $d_1 \neq 0$, then the k^{th} digit chopped floating point representation of P is

$$f_{\text{chop}}(P) = \pm 0.d_1 d_2 \dots d_k * 10^n$$

the k^{th} digit round off floating point representation of P is

$$f_{\text{round}}(P) = \pm 0.d_1 d_2 \dots d_{k-1} r_k \times 10^n$$

where r_k is obtained by rounding d_{k+1}, d_{k+2}, \dots

$P = 0.1234 \mid 444445$

4 digits chopped

$$f(1) = 0.1235$$

round

Final

use 4 digits arithmetic (round). حفازك بعد اول فنزلة غير مفترضة

$$\frac{\frac{3}{7} + \frac{5}{8} + \frac{11}{15}}{21} = ?? \quad \text{or} \quad \frac{\frac{3}{7} + 0.5967 + \frac{11}{15}}{21} = ??$$

$$\frac{(0.4286 + 0.5967) + 0.7333}{21}$$


$$0.4286 + 0.5967 = 1.0253 \approx 1.025$$

$$1.025 + 0.7333 = 1.7583 \approx 1.758$$

$$\frac{1.758}{21} = 0.08371$$

Order of estimation

$$e^x \approx 1+x$$



$$e^x \approx 1+x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

$$e^x \approx 1+x$$

$$e^h \approx 1+h \quad h \approx 0 \quad \text{order of approximation.}$$

$$\text{Error} = \frac{h^2}{2!} \approx O(h^2)$$

$$e^{0.1} \approx 1+0.1 \approx 1.1 \quad \text{error} = C h^2$$

$$e^{0.1} = 1.105170918 \quad \begin{aligned} & \downarrow \\ & = C(0.1)^2 \\ & = C(0.01) \\ & \leq 10^{-2} \end{aligned}$$

$$e^h \approx 1+h + \frac{h^2}{2!}$$

$$\text{Error} = C h^3 = O(h^3)$$

$$e^{0.1} = 1+0.1 + \frac{0.01}{2} = 1.105$$

$$\text{Error} \approx C(0.1)^3$$

$$\approx C(0.001) \leq 10^{-3}$$

$$\sin(0.1) \approx 0.1$$

$$\sin h \approx h \text{ with error } O(h^3).$$

$$\sinh h \approx h - h^3 \text{ with error } O(h^5)$$

$$\text{suppose } e^h \approx 1+h \quad \text{Error} = O(h^2) \quad (0.01)$$

$$\sin h = h - \frac{h^3}{3!} \quad \text{Error} = O(h^5) \quad (0.00001)$$

$$e^h + \sin h \approx 1+2h - \frac{h^3}{3!} \quad \text{with Error } O(h^2) + O(h^5)$$

نحوی
نحوی جوں میں ہے

$$\approx 1+2h+O(h^2)$$

defi- Order of approximation

assume that $f(h)$ is approximated by $p(h)$ and there exists a real constant $M > 0$ and a positive integer n so that

$$\frac{|f(h) - p(h)|}{|h^n|} \leq M \quad \text{for small } h$$

we say $p(h)$ approximate $f(h)$ with order of approximation $O(h^n)$ and we write $f(h) = p(h) + O(h^n)$

$$|f(h) - p(h)| \leq M|h^n|$$

$$f(h) - p(h) \approx Ch^n$$

Ex: Show that $p(h) = 1+h$ estimate of $f(h) = e^h$ with order $O(h^2)$

or

Show that $e^h = 1+h+O(h^2)$

$$e^h = 1+h + \frac{h^2}{2!} + \frac{h^3}{3!} + \dots$$

$$\frac{|e^h - (1+h)|}{|h^2|} = \frac{\frac{h^2}{2} + \frac{h^3}{3!} + \frac{h^4}{4!} + \dots}{h^2} = \frac{\frac{1}{2} + \frac{h}{3!} + \frac{h^2}{4!} + \frac{h^3}{5!} + \dots}{h^2}$$

↓ harmonic series
 $(\sum \frac{1}{n})$ diverges

$$< \frac{1}{2} + \frac{1}{3!} + \frac{1}{4!} + \frac{1}{5!} + \dots$$

$$< \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \dots \rightarrow$$

$$\text{geometric series} = \frac{1/2}{1-1/2} = 1$$

$$e^h = 1+h+O(h^2)$$

Exercise

Show that

$$-\sin h = h - \frac{h^3}{3!} + O(h^5)$$

$$2. f(h) = \sum_{k=0}^n f(h) \times h^k + O(h^{n+1})$$

Theory:-

$$\text{assume that } f(h) = P(h) + O(h^n)$$

$$g(h) = Q(h) + O(h^m)$$

$$\text{and } r = \min [m, n] \\ \text{then}$$

$$f(h) \pm g(h) = P(h) + Q(h) + O(h^r)$$

$$f(h) \cdot g(h) = P(h) Q(h) + O(h^r)$$

$$\frac{f(h)}{g(h)} = \frac{P(h)}{Q(h)} + O(h^r) \quad Q(h), Q(h) \neq 0.$$

Ex:-

$$f(h) = P(h) + O(h^3)$$

$$g(h) = Q(h) + O(h^2)$$

$$\frac{f(h)}{g(h)} = \frac{P(h)}{Q(h)} + O(h^2)$$

Ex:- (loss of significant)

$$f(x) = x(\sqrt{x+1} + \sqrt{x})$$

$$g(x) = \frac{x}{\sqrt{x+1} + \sqrt{x}}$$

use 6 digits arithmetic and round to find $f(500)$, $g(500)$

$$\begin{aligned} f(500) &= 500(\sqrt{501} - \sqrt{500}) \\ &= 500(22.3830 - 22.3607) \\ &= 500(0.0223200) = 11.1500. \end{aligned}$$

$$g(500) = \frac{500}{\sqrt{501} + \sqrt{500}} = \frac{500}{22.3830 + 22.3607} = 11.1748$$

exact answer = 11.174755...

المethode الثانية أحسن لأن في الملة المدخلة

signigant digits

$$\frac{3}{17} = 0.176470588 + \epsilon$$

Note:-

$$P = \tilde{P} + \epsilon_P$$

$$q = \tilde{q} + \epsilon_q$$

$$P + q = \tilde{P} + \tilde{q} + \epsilon_P + \epsilon_q \\ = \tilde{P} + \tilde{q} + \epsilon_{P+q}$$

$$P \cdot q = (\tilde{P} + \epsilon_P)(\tilde{q} + \epsilon_q) \\ = \tilde{P}\tilde{q} + \tilde{P}\epsilon_q + \tilde{q}\epsilon_P + \epsilon_P\epsilon_q \\ = \tilde{P}\tilde{q} + \epsilon_{Pq}$$

$$P = 9.8 \times 10^6 + 35 \times 10^{-9}$$

$$\tilde{q} = 3.6 \times 10^7 + 2.4 \times 10^{-9}$$

$$[a_0, b_0] = [a, b]$$

$$c_0 = b_0 - \frac{f(b_0)(b_0 - a_0)}{f(b_0) - f(a_0)}$$

$$f(c_0)$$

If $f(c_0) = 0$ done.

else if $f(c_0) \cdot f(a_0) < 0 \Rightarrow [a_1, b_1] = [a_0, c_0]$

else $[a_1, b_1] = [c_0, b_0]$

$$c_n = b_n - \frac{f(b_n)(b_n - a_n)}{f(b_n) - f(a_n)}$$

example

Solve $x \sin x = 1$.

$$f(x) = x \sin x - 1$$

$$f(0) = -1$$

$$f(2) = 0.81859485$$

$$c_0 = b_0 - \frac{f(b_0)(b_0 - a_0)}{f(b_0) - f(a_0)}$$

$$= 2 - \frac{0.81859485(2 - 0)}{0.81859485 - (-1)} = 1.09975017$$

$$F(c_0) = 1.09975017 \sin(1.09975017) - 1$$

$$= -0.02001912$$

$$[a_1, b_1] = [1.09975017, 2]$$

$$c_1 = b_1 - \frac{f(b_1)(b_1 - a_1)}{f(b_1) - f(a_1)} = 2 - \frac{0.81859485(2 - 1.09975017)}{0.81859485 - (-0.02001912)}$$

$$= 1.12124074$$

$$f(c_1) = 0.00983461$$

$$[a_2, b_2] = [1.09975017, 1.12124074]$$

$$c_2 = 1.11416120$$

$$c_3 = 1.11415714$$

Section 2.1

Fixed point iteration

To solve $f(x)=0$ we solve $x=g(x)$ [where $f(x)=x-g(x)$]
 i.e to find the roots of $F \rightarrow$ we find the fixed point of $g(x)$.

Def:- P is a fixed point of g iff $g(P)=P$.

$$1. g(x) = \frac{1}{x}$$


fixed points $1, -1$.

$$g(P) = P$$

$$\frac{1}{P} = P \Rightarrow P^2 = 1 \Rightarrow P = \pm 1$$

$$2. g(x) = x+1 . \text{ No fixed points}$$

$$3. g(x) = x . \text{ all points are fixed points.}$$



Def:- Fixed point iteration:-

start with P_0 , $P_{n+1} = g(P_n)$, $n=0, 1, 2, 3, \dots$

$$P_1 = g(P_0)$$

$$P_2 = g(P_1)$$

⋮

Theorem:-

If the fixed point iteration converges to P , then P is the fixed point of g .

$$\lim_{n \rightarrow \infty} P_n = P \Rightarrow \lim_{n \rightarrow \infty} g(P_{n+1}) = \lim_{n \rightarrow \infty} g(P_n) = g(\lim_{n \rightarrow \infty} P_n) = g(P)$$

Since $P_{n+1} = g(P_n)$ $\downarrow = P$.

example 2:

Solve $x^2 - 2x - 3 = 0 \Rightarrow f(x) = 0$.
 $(x-3)(x+1) = 0$.

$$\begin{aligned} x &= 3 \\ x &= -1 \end{aligned}$$

$$x^2 = 2x + 3$$

$$x = \sqrt{2x+3} = g(x).$$

if $P_0 = 4$.

$$P_1 = g(4) = g(P_0) = \sqrt{11} = 3.31662.$$

$$P_2 = g(P_1) = g(3.31662) = \sqrt{9.63325} = 3.10375$$

$$P_3 = 3.03439$$

$$P_4 = 3.01184$$

Note that 3 is a fixed point of

$$P_n \rightarrow 3$$

$$g(x) = \sqrt{2x+3} \text{ because } g(3) = 3$$

way 2:- X $\overset{?}{\underset{\text{divergence.}}{\longrightarrow}}$

$$2x = x^2 - 3$$

$$x = \frac{x^2 - 3}{2} = g(x)$$

$$P_0 = 4$$

$$P_1 = g(4) = 6.5$$

$$P_2 = g(6.5) = 19.625$$

$$P_3 = 191.07$$

way 3:-

$$x(x-2) = 3 \Rightarrow x = \frac{3}{x-2} = g(x).$$

$$P_0 = 4$$

$$P_1 = g(4) = \frac{3}{2} = 1.5$$

$$P_2 = -6$$

$$P_3 = -0.375$$

$$P_4 = -1.26315$$

$$P_5 = -0.919355$$

$$P_6 = -1.02762$$

$$P_7 = -0.990876$$

Theorem:- (fixed point Theorem I).

assume $g \in C[a,b]$ if $g(x) \in [a,b]$ for all $x \in [a,b]$ then g has a fixed point in $[a,b]$ furthermore if $|g'(x)| \leq k < 1$ for all $x \in [a,b]$ then g has a unique fixed point.

Proof:-

if $g(a)=a$ or $g(b)=b$ done.

if not $g(a) > a$ and $g(b) < b$.

let $h(x) = g(x) - x$, h continuous.

$$h(a) = g(a) - a > 0$$

$$h(b) = g(b) - b < 0$$

by bolzano $\exists c \in \mathbb{C}$ such that $h(c)=0$.

$$g(c) - c = 0$$

$$\boxed{g(c) = c}$$

Uniqueness

Suppose $\exists P_1, P_2$ such that $g(P_1) = P_1, g(P_2) = P_2$.

Using mean value theorem on (P_1, P_2) .

$\exists c \in (P_1, P_2)$ such that $\left| \frac{g(P_2) - g(P_1)}{P_2 - P_1} \right| = |g'(c)| < 1$

$$\frac{P_2 - P_1}{P_2 - P_1} = 1 \Rightarrow 1 < 1 \rightarrow \text{Contradiction}$$

Theorem:- (fixed point iteration theorem) $P_1 = P_2$. \times

assume that $g(x)$ and $g'(x)$ are continuous on a balanced interval

$(a,b) = (P-S, P+S)$ that contains a unique fixed point P and that the started value P_0 is chosen in this interval.

1. if $g'(x) \leq k < 1$ for all $x \in (a,b)$ then the FPI converge.

$P_{n+1} = g(P_n)$ will converge (attractive Fixed point).

2. if $g'(x) > 1$ for all $x \in (a,b)$ then the fixed point iteration diverges (we call it repulsive Fixed point).

Note:-

If p is given we can replace the above two conditions by

1. if $|g'(p)| < 1 \rightarrow$ the FPI converges.
2. if $|g'(p)| \geq 1 \rightarrow$ the FPI diverges.

Convergence.


$$|g'(x)| < 1$$

$$-1 < g'(x) < 0$$

$$0 < g'(x) < 1$$

$$0 < g'(x) < 1$$


monotone convergence.



$$P_1 = g(P_0)$$

$$-1 < g'(x) < 0$$


alternating convergence



divergence $|g'(x)| > 1$

$$g'(x) > 1 \quad g'(x) < -1$$

$$g'(x) > 1$$



example

investigate the nature of the FPT and show your answer by examples for

$$g(x) = 1 + x - \frac{x^2}{4}$$

solution

$$x = g(x).$$

$$x = 1 + x - \frac{x^2}{4}.$$

$$x^2 = 4$$

$$x = \pm 2 \text{ (Fixed points)}$$

when $x = 2$,

$$g'(x) = 1 - \frac{x}{2}$$

$|g'(2)| = 0 \Leftrightarrow$ Convergence Fixed point. (attractive Fixed point)

to show that:-

$$\text{let. } P_0 = 1.6.$$

$$P_1 = g(1.6) = 1.96.$$

$$P_2 = g(1.96) = 1.996.$$

$$\text{if } P_n \rightarrow 2.$$

$$P_0 = 2.5.$$

$$P_1 = g(2.5)$$

at $x = -2$

~~$|g(x)|$~~ $|g'(-2)| = 2 > 1$ diverge \rightarrow FPI diverge. (Repulsive fixed point).

$$P_0 = -2.05.$$

$$P_1 = g(-2.05) \approx -2.1 \dots$$

$$P_2 = g(-2.1) = -2.2.$$


$P_n \rightarrow$ divergence.

Proof:-

by mean value.

$$|P_1 - P| = |g(P_0) - g(P)| = |g'(c)| (P_0 - P) < (P_0 - P)$$

$\rightarrow P_1$ is closer to P from P_0 .



$P_0 \leftarrow P \leftarrow P_1 \leftarrow P_2$

$$\underbrace{P_0 \leftarrow P \leftarrow P_1 \leftarrow P_2}_{\text{الخط}} \leq k_2 (P_{n-1} - P) \leq k_2^2 (P_{n-2} - P)$$

$$\leq k \cdot k \cdot k_2 (P_{n-3} - P)$$

$$\leq k^n (P_0 - P)$$

$$\lim_{n \rightarrow \infty} |P_n - P| = 0$$

$$\lim_{n \rightarrow \infty} P_n = P.$$

② $|P - P| = |g'(c)| |P_0 - P|$ $\frac{\nexists}{> 1}$ $|P_0 - P| > |P_0 - P|$

Theorem:-

a. $|P_n - P| \leq k^n |P_0 - P|$.
 $\frac{\nexists}{\text{error}}$

exists such
upper bound for error

k is the upper bound

$$k = g'(P) \rightarrow \text{given } k, P.$$

b. $|P_n - P| \leq \frac{k^n}{1-k} |P_0 - P|$ (exercise).

\rightarrow we can find n

Example:-

$$x^3 - x + 5 = 0$$

Use Fixed point iteration to find all the roots, find k for each case

QWALIAK.

XBAKASAK.

XBAKASAK.


$$F(x) = x^3 - x + 5$$

$$F(0) = 5$$

$$F(-1) = 5$$

$$F(-2) = -11$$

$$F(2) = 1$$



$$x^3 = x + 5$$

$$x = \sqrt[3]{x+5} = (x+5)^{1/3}$$

$$g(x) = x$$

$$g'(x) = \frac{1}{3}(x+5)^{-2/3}$$

$$= \frac{1}{3\sqrt[3]{(x+5)^2}} < 1$$

for all x
 $0 < x$

$$P_0 = 1.5$$

start
iteration

For $x > 0$.

$$x+5 > 5$$

$$(x+5)^2 > 25$$

$$\sqrt[3]{x+5} > (25)^{\frac{1}{3}} > 2$$

$$\frac{1}{3\sqrt[3]{(x+5)}} < \frac{1}{2}$$


$$\frac{1}{3\sqrt[3]{(x+5)}} < \frac{1}{6}$$

$$k = \frac{1}{6}$$

DISCUSSION

$$f(x) = \frac{\cos(x-1)}{1+e^{x-2}}$$

max point. قيم صيغ نطلع



5.

$$x^4 - 3x^2 - 3 = 0$$

$$P_0 = 1$$

$$10^{-2}$$

$$[1, 2]$$

$$x^4 = 3x^2 + 3.$$

$$x = \sqrt[4]{3x^2 + 3}.$$

$$P_1 = g(1) = \sqrt[4]{6} = 1.5650$$

$$P_2 = 1.79358$$

$$P_3 = 1.88595$$

$$P_4 = 1.92285$$

لزغنا حتى 5 iteration.

$$P_5 = 1.93751$$

ثبتنا منزلي

$$P_6 = 1.94332$$

$$y_c^{(2.2)} \\ P_n = P_{n-1} - \frac{P_{n-1}^5 - 7}{5P_{n-1}^4}$$

$$g(x) = x - \frac{x^5 - 7}{5x^4}$$

$$g'(x) = x - \frac{f(x)}{f'(x)}$$

$$g(x) = x - \frac{x}{5} + \frac{7}{5x^4}$$

$$g(x) = \frac{4x}{5} + \frac{7}{5x^4}$$

$$P = 7^{1/5} \\ P_n = g(P_{n-1})$$

$$x = 7^{1/5}$$

$$x^5 = 7$$

$$x^5 - 7 = 0$$

$$f(x) = x^5 - 7$$

$$g'(7^{\frac{1}{15}}) = \frac{4}{5} - \frac{28}{5(7^{\frac{1}{15}})^5}$$

$$= \frac{4}{5} - \frac{28}{5 \cdot 7} = \frac{4}{5} - \frac{4}{5} = 0. \text{ Method } \mathcal{E}^{-1}$$

newton method.

2.2, 2.4, 2.5, 6, 16, 17

2.1,

P&P
14
2.2

Solve

$$x = \tan x \quad \text{in } [4, 5]$$

$$g(x) = \sec^2 x > 1.$$

$$x = \tan^{-1} x$$

$$g(x) = \tan^{-1} x$$

$$g'(x) = \frac{1}{1+x^2} < 1$$

$$P_0 = 4.5.$$

$$P_1 = \tan^{-1}(4.5)$$

$$= 1.352127$$

$$P_2 = \tan^{-1}(P_1)$$

$$= 0.93$$

$$x = \tan x = \tan(x - \pi) = \tan(x + \pi)$$

$$x = \tan(x - \pi)$$


$$\tan^{-1} x = x - \pi$$

$$x = \tan^{-1} x + \pi$$

$$P_0 = 4.5.$$

$$P_1 = 4.49372$$

$$P_2 = 4.49342$$



(4)
2.1

$$\text{Let } f(x) = (x-1)^{10}$$

$$P = 1$$

$$P_n = 1 + \frac{1}{n}$$

Show that if $|f(P_n)| < 10^{-3}$

but $|P - P_n| < 10^{-3}$ requires $n > 1000$

$$1. |P - P_n| < \epsilon$$

$$2. |C_{n+1} - C_n| < \epsilon$$

$$3. |f(P_n)| < \epsilon$$

$$F(p_n) = \left(\frac{1}{n}\right)^{10} < 10^{-3} \text{ for } n > 1.$$

$$|P - p_n| < 10^{-3} = |1 - 1 - \frac{1}{n}| < 10^{-3}$$

$$\left|\frac{1}{n}\right| < 10^{-3}$$

$$\frac{1}{n} < 10^{-3} \Rightarrow n > 1000.$$

15
2.1

$$p_n = \sum_{k=1}^{\infty} \frac{1}{k}$$

Show that p_n diverge even though $\lim_{n \rightarrow \infty} (p_n - p_{n-1}) = 0$.

$$P = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{6} + \dots$$

$$\lim_{n \rightarrow \infty} p_n = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \dots + \frac{1}{n} = \sum_{n=0}^{\infty} \frac{1}{n} \text{ harmonic series}$$

$$p_n - p_{n-1} = \left(1 + \frac{1}{2} + \dots + \frac{1}{n-1} + \frac{1}{n}\right) - \left(1 + \frac{1}{2} + \dots + \frac{1}{n-1}\right) = \frac{1}{n}$$

$$\lim_{n \rightarrow \infty} (p_n - p_{n-1}) = \lim_{n \rightarrow \infty} \frac{1}{n} = 0.$$

$$c_n = \frac{a_n + b_n}{2} \text{ stop.}$$

$$F(c_n) \leq \epsilon \text{ or } |c_n - c_{n-1}| \leq \epsilon$$

$$\text{stop if } F(c_n) \leq \epsilon \text{ and } \frac{|c_n - c_{n-1}|}{c(c_{n-1})} \leq 1 \times 10^{-6}.$$

Solve this equ

$$3x^2 - e^x = 0$$


$$f(0) = -1$$

$$f(1) = 0 \cdot 28 > 0$$

$$f(2) = 12 - e^2 > 0$$

$$f(3) = 27 - e^3 > 0$$

$$f(4) = 48 - e^4 < 0$$



Newton method

$$f'(P_0) = \frac{f(P_0) - 0}{P_0 - A}$$

$$P_0 - P_1 = \frac{f(P_0)}{f'(P_0)}$$


$$P_1 = P_0 - \frac{f(P_0)}{f'(P_0)}$$

$$P_2 = P_1 - \frac{f(P_1)}{f'(P_1)}$$

$$P_{n+1} = P_n - \frac{f(P_n)}{f'(P_n)}$$

$$x = x - \frac{f(x)}{f'(x)}$$

$g(x)$ ← Newton fixed point function



Th:- Newton Raphson theorem

assume $f \in C^2[a,b]$ and $\exists P \in [a,b]$ such that $f(P)=0$, if $f'(P) \neq 0$ then there exist a $\delta > 0$ such that the sequence

$$\{P_{k\epsilon}\}_{k=0}^{\infty} \text{ which is defined by } P_{k\epsilon} = g(P_{(k-1)\epsilon}) = P_{(k-1)\epsilon} - \frac{f(P_{(k-1)\epsilon})}{f'(P_{(k-1)\epsilon})}$$

will converge to P for any initial approximation $P_0 \in [P-\delta, P+\delta]$

example:-

estimate $5^{\frac{3}{7}}$

$$x = 5$$

$$x^7 = 5^3$$

$$f(x) = x^7 - 125$$

$$P_1 = P_0 - \frac{f(P_0)}{f'(P_0)}$$

$$f'(x) = 7x^6$$

$$\begin{aligned} P_{n+1} &= P_n - \frac{f(P_n)}{f'(P_n)} \\ &= P_n - \frac{P_n^7 - 125}{7P_n^6} \\ &= \frac{6}{7}P_n + \frac{125}{7P_n^6} \end{aligned}$$

$$P_0 = 2$$

$$P_1 = \frac{6}{7}(2) + \frac{125}{7(2)^6} = 1.71429$$

$$P_2 = \frac{6}{7}(1.71429) + \frac{125}{7(1.71429)^6} = 2.07$$

Proof the theorem

$$g(x) = x - \frac{f(x)}{f'(x)}$$

$$g'(x) = 1 - \frac{f'(x)f'(x) - f(x)f''(x)}{(f'(x))^2}$$

$$g'(x) = \frac{f(x)f''(x)}{(f'(x))^2}$$

$$g'(P) = \frac{f(P)f''(P)}{(f'(P))^2} = 0$$

theory

→ by Fixed point iteration \rightarrow the Fixed point iteration will converge.

- if $e_{n+1} \approx A e_n$ where $e_n = P - P_n$
- or $e_{n+1} \approx \frac{1}{100} e_n$ (error smaller than the first) (the best one)
- $e_{n+1} \approx \frac{1}{2} e_n$ results \downarrow

جواب ایسا

Definition

P is a root of multiplicity M of $f(x)$ if $f(x) = (x-P)^M h(x)$, $h(P) \neq 0$.

- $f(x) = x^3 - 3x + 2$

1 is a root of $f(x)$

what is the multiplicity of 1 ?

$$\begin{array}{r} x^2 + x - 2 \\ \hline x-1 \quad | \quad x^3 - 3x + 2 \\ \quad x^3 + x^2 \\ \hline \quad x^2 - 3x + 2 \\ \quad x^2 + x \\ \hline \quad -2x + 2 \\ \quad +2x + 2 \\ \hline \quad 0 \end{array}$$

$$\begin{array}{r} x+2 \\ \hline x-1 \quad | \quad x^2 + x - 2 \\ \quad x^2 - x \\ \hline \quad 2x - 2 \\ \quad 2x - 2 \\ \hline \quad 0 \end{array}$$

$$f(x) = (x-1)(x^2 + x - 2)$$

1 has multiplicity 2 (quadratic root) $M=2$

-2 is a simple root ($M=1$)

Theory:-

P is a root of multiplicity M of $f(x)$ iff.

$$f(P)=0, f'(P)=0, \dots, f^{(m-1)}(P)=0 \text{ but } f^{(m)}(P) \neq 0$$

$$f^{(m)}(P) \neq 0$$

Example:-

$$f(x) = x^3 - 3x + 2$$

$$f(1) = 0$$

$$f'(x) = 3x^2 - 3$$

$$f'(1) = 0$$

$$f''(x) = 6x$$

$$f''(1) = 6$$

$$M = 2$$

$$e_{n+1} \approx Ae_n$$

$$\frac{e_{n+1}}{e_n} \approx A$$

$$\lim_{n \rightarrow \infty} \frac{e_{n+1}}{e_n} = A \rightarrow \text{linear Convergence.}$$

$$\text{if } e_{n+1} \approx Ae_n^2$$

$$\lim_{n \rightarrow \infty} \frac{e_{n+1}}{e_n^2} \approx A \rightarrow \text{quadratic Convergence.}$$

Definition:- Order of convergence

assume $p_n \rightarrow p$ and $e_n = p - p_n$, if there exists two positive numbers A, R such that

$$\lim_{n \rightarrow \infty} \frac{|e_{n+1}|}{|e_n|^R} = A$$

Then the sequence is said to converge to p with order of convergence R , A is called the Asymptotic error constant.

if $R=1$, we call it linear convergence.

if $R=2$, we call it quadratic convergence.

example:-

Show that $p_n = \frac{1}{n^3}$ converges to $\overset{\downarrow}{p}$ linearly??

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{|e_{n+1}|}{|e_n|} &= \lim_{n \rightarrow \infty} \frac{\left|0 - \frac{1}{(n+1)^3}\right|}{\left|0 - \frac{1}{n^3}\right|} = \lim_{n \rightarrow \infty} \frac{\frac{1}{(n+1)^3}}{\frac{1}{n^3}} = \lim_{n \rightarrow \infty} \frac{n^3}{(n+1)^3} \\ &= \left(\lim_{n \rightarrow \infty} \frac{n}{n+1}\right)^3 = 1 \end{aligned}$$

$\frac{1}{n^3} \rightarrow 0$ linearly
converge to,

Example:-

$$f(x) = x^{101} - x^{100} - x + 1$$

$$f(1) = 0$$

$$f'(x) = 101x^{100} - 100x^{99} - 1$$

$$f'(1) = 101 - 100 - 1 = 0$$

$$f''(x) = (101)(100)x^{99} - (100)(99)x^{98}$$

$$f''(1) \neq 0$$

$$M = 2.$$

Theorem:- Convergence of Newton method

if we use Newton iteration,

1. if P is a simple root, then

$$\lim_{n \rightarrow \infty} \frac{|e_{n+1}|}{|e_n|^2} = \left| \frac{f''(P)}{2f'(P)} \right|$$

$\left[\begin{array}{l} P \text{ is a simple root} \\ \text{Convergence is quadratic} \\ A = \left| \frac{f''(P)}{2f'(P)} \right|, R=2 \end{array} \right]$

2. if P has multiplicity $M > 1$, then

$$\lim_{n \rightarrow \infty} \frac{|e_{n+1}|}{|e_n|} = \frac{M-1}{M}$$

$\left[\begin{array}{l} \text{Convergence is linear} \\ A = \frac{M-1}{M}, R=1 \end{array} \right]$

example

$$f(x) = x^3 - 3x + 2$$

$$f'(x) = 3x^2 - 3$$

$$f(x) = (x-1)^2(x+2)$$

$$f''(x) = 6x$$

-2 is a simple roots

convergence is fast $R=2$ ($\frac{|e_{n+1}|}{|e_n|^2} = \left| \frac{f''(P)}{2f'(P)} \right|$)

$$A = \left| \frac{f''(-2)}{2f'(-2)} \right| = \left| \frac{-12}{2(9)} \right| = \frac{2}{3}$$

$$P=1, M=2$$

linear convergence ($R=1$)

$$P_{n+1} = P_n - \frac{f(P_n)}{f'(P_n)}$$

$$P_0 = -2.4$$

n	P_n	e_n	$\frac{ e_{n+1} }{ e_n }$
0	-2.4	0.4	
1	-2.0761904	0.0761904	0.4761
2	-2.0035960	0.003596	0.6194
3	-2.00000858	0.000008589	0.6642

\downarrow $\frac{2}{3} \approx A$

Fast Convergence.

$$P_0 = 1.2$$

n	P_n	e_n	$\frac{ e_{n+1} }{ e_n }$
0	1.2	-0.2	
1	1.103030	-0.10303	0.515
2	1.052356	-0.052356	0.5081
3	1.0264008	-0.02640081	0.4962

\downarrow $A \approx \frac{1}{2}$

Slow Convergence.

Theory:- Accelerated newton method

if P is a root of multiplicity M then the iteration

$$P_{n+1} = P_n - \frac{Mf(P_n)}{f'(P_n)}$$
 will converge quadratically to P .

Ex:-

For the previous example. $f(x) = (x-1)^2(x+2)$

has multiplicity 2, if we use the accelerated Newton iteration

$$P_{n+1} = P_n - \frac{2f(P_n)}{f'(P_n)}$$
 will get quadratic convergence!

$$P_0 = 1.2$$

n	P_n	e_n	$\frac{ e_{n+1} }{ e_n ^2}$
0	1.2	-0.2	
1	1.0060606	-0.00606	0.15
2	1.000006087	-0.000006087	0.15

Proof $L(p) \approx g(p) \approx 0, g(p) = p$

$$g(x) = g(p) + g'(p)(x-p) + \frac{g''(c)}{2}(x-p)^2$$

$$P_{n+1} = g(P_n) \approx p + 0 + \frac{g''(c)}{2}(P_n - p)^2$$

Secant method:-


$$\frac{f(p_1) - 0}{p_1 - p_2} = \frac{f(p_1) - f(p_0)}{p_1 - p_0}$$

$$p_1 - p_2 = \frac{f(p_1)(p_1 - p_0)}{f(p_1) - f(p_0)}$$

$$p_2 = p_1 - \frac{f(p_1)(p_1 - p_0)}{f(p_1) - f(p_0)}$$

$$p_3 = p_2 - \frac{f(p_2)(p_2 - p_1)}{f(p_2) - f(p_1)}$$

$$p_n = p_{n-1} - \frac{f(p_{n-1})(p_{n-1} - p_{n-2})}{f(p_{n-1}) - f(p_{n-2})}$$



Theorem:-

If we use Secant method to get $p_n \rightarrow p$. Then.

$$\lim_{n \rightarrow \infty} \frac{|e_{n+1}|}{|e_n|^{1.618}} = \left| \frac{f''(p)}{2f'(p)} \right|^{0.618}$$

$$\rightarrow R = 1.618 = \frac{1 + \sqrt{5}}{2}$$

Ex:-

$$f(x) = (x+2)(x-1)^2$$

$$p_0 = -2.15, \quad p_1 = -2.4$$

and we use Secant method.

n	p_n	e_n	$\frac{ e_{n+1} }{ e_n ^{1.618}}$
0	-2.6	0.6	
1	-2.4	0.4	
2	-2.106598	0.106598	
3	-2.02264	0.02264	
4	-2.00161	0.00161	

False position method

Speed

1

Cost

1

Convergence

Very accurate

Secant method

1.6

1

depends on
 P_0, P_1

Newton method

2

2

depends on
 P_0

2.6 Fixed point iteration for system of equation

$$x^2 \cos y + y \sin x = 10$$

$$y \ln x + x^2 \cos y = 5$$


$$x^2 - y^2 = 1$$

$$x^2 + y^2 = 2$$

$$2x^2 = 3$$

$$x^2 = \frac{3}{2}$$

$$x = \pm \sqrt{\frac{3}{2}}$$



$$x^2 - y^2 = x + 3$$

$$x^2 + y^2 = e^x - 1$$

$$2x^2 = x + 3 + e^x - 1$$

$$2x^2 - x - e^x - 2 = 0$$

$$x = g_1(x, y)$$

$$y = g_2(x, y)$$

$$(P_0, g_0)$$

$$P_1 = g_1(P_0, g_0)$$

$$P_2 = g_2(P_1, g_1)$$

$$g_1 = g_2(P_0, g_0)$$

$$g_2 = g_2(P_1, g_1)$$

$$P_{n+1} = g_1(P_n, g_n)$$

$$g_{n+1} = g_2(P_n, g_n)$$

Definition:-

(P, g) is a Fixed Point of the system

$x = g_1(x, y), y = g_2(x, y)$ if $P_x = g_1(P, g)$ and $g_y = g_2(P, g)$

Defi:-

Fixed point iteration for the system

$x = g_1(x, y), y = g_2(x, y)$ is given (P_0, g_0) then

$$P_{n+1} = g_1(P_n, g_n)$$

$$g_{n+1} = g_2(P_n, g_n) \quad n=1, 2, 3, \dots$$


Ex:-

$$f_1(x, y) = x^2 - 2x - y + 0.5 = 0$$

$$f_2(x, y) = x^2 + 4y^2 - 4 = 0$$

estimate the solutions ?

$$\begin{aligned} & x^2 + 4y^2 = 4 \\ & \frac{x^2}{4} + y^2 = 1 \end{aligned}$$



$$x = \frac{x^2 - y + 0.5}{2} = g_1(x, y).$$

$$y = \frac{-x^2 - 4y^2 + 8y + 4}{8} = g_2(x, y).$$

$$(P_0, g_0) = (0, 1)$$

$$P_1 = g_1(0, 1) = \frac{0 - 1 + 0.5}{2} = -0.25$$

$$g_1 = g_2(0, 1) = \frac{0 - 4 - 8 + 4}{8} = 1$$

$$P_4 = -0.2221680$$

$$g_4 = 0.9938121$$

$$P_5 = -0.222194$$

$$g_5 = 0.9938095$$

$$(P_0, g_0) = (2, 0) \quad (\text{diverges})$$

$$P_1 = g_1(2, 0) = 2.25$$

$$g_1 = g_2(2, 0) = 0$$

$$\text{Let } g_1(x, y) = \frac{-x^2 + 4x + y - 0.5}{2}$$

$$g_2(x, y) = \frac{-x^2 - 4y^2 + 11x + 4}{11}$$

$$(P_0, g_0) = (2, 1)$$

$$(2, 1) \rightarrow \text{Converge}$$

This: Fixed point iteration - For system of equation:-

assume $g_1(x,y)$, $g_2(x,y)$ and their partial derivative are continuous on a region that contains^{*} the fixed point (P,Q) , if the starting point (P_0, Q_0) is chosen sufficiently close to (P,Q) and.

$$\left| \frac{dg_1}{dx} \right| + \left| \frac{dg_1}{dy} \right| < 1 \text{ and } \left| \frac{dg_2}{dx} \right| + \left| \frac{dg_2}{dy} \right| < 1 \text{ in that region}$$

then the FPI will converge.

Note:-

if (P,Q) is given we apply the condition at (P,Q) only.

to proof
المطلب المطلوب

Fixed point \rightarrow we talk about g 's
Newton \rightarrow we talk about F .

if $|x| < 0.5$ and $0.5 < y < 1.5$ \rightarrow دوّن تمارين الفرق.

$$\left| \frac{dg_1}{dx} \right| + \left| \frac{dg_2}{dy} \right| = |x| + 0.5 < 1$$

+
أكبر قيمة
0.5

$$\left| \frac{dg_2}{dx} \right| + \left| \frac{dg_2}{dy} \right| = \frac{|x|}{4} + |1-y| < \frac{1}{4} + 0.5 < 1$$

+
أكبر قيمة
0.5
+
أكبر قيمة
1.5

حتى نثبت أن النقطة المكانة تحت تمارين فرق \leftarrow divergence \rightarrow تمارين فرق \rightarrow لا تتحقق الافتراض
السابقين او لا تتحقق سطر واحد من الافتراض.

Example (linear system)

$$3x + 2y + 7z = 10 \rightarrow x = \frac{10 - 2y - 7z}{3} = g_1(x, y, z)$$

$$2x + 4y - z = 4 \rightarrow y = \frac{4 + z - 2x}{4} = g_2(x, y, z)$$

$$x + 5y + 10z = 15 \rightarrow z = \frac{15 - x - 5y}{10} = g_3(x, y, z).$$

$$P_1 = g_1(P_0, g_0, r_0)$$

$$g_1 = g_2(P_0, g_0, r_0)$$

$$r_1 = g_3(P_0, g_0, r_0)$$

$$\rightarrow \begin{aligned} P_1 &= g_1(P_0, g_0, r_0) \\ g_1 &= g_2(P_1, g_0, r_0) \\ r_1 &= g_3(P_1, g_0, r_0) \end{aligned} \quad \left. \begin{array}{l} \\ \\ \end{array} \right\} \text{Gauss-Sidel method}$$

$$P_{n+1} = P_n - \frac{f(P_n)}{f'(P_n)}$$

$$f_1(x, y) = 0$$

$$f_2(x, y) = 0$$

$$\begin{pmatrix} P_{n+1} \\ g_{n+1} \end{pmatrix} = \begin{pmatrix} P_n \\ g_n \end{pmatrix} - J^{-1} \begin{pmatrix} f_1(P_n, g_n) \\ f_2(P_n, g_n) \end{pmatrix}$$

↓
Jacobian: (P_n, g_n)

$$h: (x, y) \rightarrow (f_1(x, y), f_2(x, y))$$

$$h' = J = \begin{pmatrix} \frac{\partial f_1}{\partial x} & \frac{\partial f_1}{\partial y} \\ \frac{\partial f_2}{\partial x} & \frac{\partial f_2}{\partial y} \end{pmatrix}$$

$$\boxed{\overrightarrow{P}_{n+1} = \overrightarrow{P}_n - J^{-1} \#}$$

2.7 Newton Method

given $F_1(x, y) = 0, F_2(x, y) = 0$

and $F_1(p, q) = 0, F_2(p, q) = 0$

starting with (P_0, q_0) close to (p, q) then using taylor expansion in two dimension at (P_0, q_0)

$$F_1(x, y) \approx F_1(P_0, q_0) + \left. \frac{dF_1}{dx} \right|_{(P_0, q_0)} (x - P_0) + \left. \frac{dF_1}{dy} \right|_{(P_0, q_0)} (y - q_0)$$

$$F_2(x, y) \approx F_2(P_0, q_0) + \left. \frac{dF_2}{dx} \right|_{(P_0, q_0)} (x - P_0) + \left. \frac{dF_2}{dy} \right|_{(P_0, q_0)} (y - q_0)$$

substitute (P, q) above

$$0 = F_1(P_0, q_0) + \left. \frac{dF_1}{dx} \right|_{(P_0, q_0)} (P - P_0) + \left. \frac{dF_1}{dy} \right|_{(P_0, q_0)} (q - q_0)$$

$$0 = F_2(P_0, q_0) + \left. \frac{dF_2}{dx} \right|_{(P_0, q_0)} (P - P_0) + \left. \frac{dF_2}{dy} \right|_{(P_0, q_0)} (q - q_0)$$

$$\begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} f_1(P_0, q_0) \\ f_2(P_0, q_0) \end{bmatrix} + \begin{bmatrix} \frac{dF_1}{dx} & \frac{dF_1}{dy} \\ \frac{dF_2}{dx} & \frac{dF_2}{dy} \end{bmatrix}_{(P_0, q_0)} \begin{bmatrix} P - P_0 \\ q - q_0 \end{bmatrix}$$

$$\boxed{\begin{bmatrix} -F_1 \\ -F_2 \end{bmatrix}_{(P_0, q_0)} = J_{(P_0, q_0)}^{-1} \begin{bmatrix} P - P_0 \\ q - q_0 \end{bmatrix}} \rightarrow \text{Direct method.}$$

$$-J^{-1} \begin{bmatrix} F_1 \\ F_2 \end{bmatrix} = \begin{bmatrix} P - P_0 \\ q - q_0 \end{bmatrix}$$

$$\begin{bmatrix} P_0 \\ q_0 \end{bmatrix} - J^{-1} \begin{bmatrix} F_1 \\ F_2 \end{bmatrix}_{(P_0, q_0)} = \begin{bmatrix} P_1 \\ q_1 \end{bmatrix} \quad \text{inverse way.}$$

- Inverse method.

$$\begin{bmatrix} p_{n+1} \\ q_{n+1} \end{bmatrix} = \begin{bmatrix} p_n \\ q_n \end{bmatrix} - J^{-1}_{(p_n, q_n)} \begin{bmatrix} f_1(p_n, q_n) \\ f_2(p_n, q_n) \end{bmatrix}$$

- Direct method

$$- \begin{bmatrix} f_1(p_n, q_n) \\ f_2(p_n, q_n) \end{bmatrix} = J_{(p_n, q_n)} \begin{bmatrix} \Delta x \\ \Delta y \end{bmatrix}$$

$$\Delta x = p_{n+1} - p_n \rightarrow p_{n+1} = \Delta x + p_n$$

$$\Delta y = q_{n+1} - q_n \rightarrow q_{n+1} = \Delta y + q_n$$

- example

Solve Used Newton method.
- inverse method.

$$x^2 - 2x - y = 0.5 \rightarrow F_1(x, y) = x^2 - 2x - y - 0.5 = 0 = f_1(x, y),$$

$$x^2 + 4y^2 = 4 \rightarrow x^2 + 4y^2 - 4 = 0 = f_2(x, y).$$

$$(p_0, q_0) = (2, 0.25)$$

$$J = \begin{pmatrix} 2x-2 & -1 \\ 2x & 8y \end{pmatrix}_{(2, 0.25)} = \begin{pmatrix} 2 & -1 \\ 4 & 2 \end{pmatrix}.$$

$$\begin{pmatrix} p_1 \\ q_1 \end{pmatrix} = \begin{pmatrix} 2 \\ 0.25 \end{pmatrix} - \begin{pmatrix} 2 & -1 \\ 4 & 2 \end{pmatrix}^{-1} \begin{pmatrix} 0.25 \\ 0.25 \end{pmatrix} \quad \begin{aligned} f_1(2, 0.25) &= 0.25 \\ f_2(2, 0.25) &= 0.25 \end{aligned}$$

$$= \begin{pmatrix} 2 \\ 0.25 \end{pmatrix} - \frac{1}{8} \begin{bmatrix} 2 & 1 \\ -4 & 2 \end{bmatrix} \begin{pmatrix} 0.25 \\ 0.25 \end{pmatrix} = \begin{pmatrix} 1.90625 \\ 0.3125 \end{pmatrix}.$$

$$\begin{pmatrix} p_2 \\ q_2 \end{pmatrix} = \begin{pmatrix} 1.90625 \\ 0.3125 \end{pmatrix} - \begin{pmatrix} 1.8125 & -1 \\ 3.8125 & 2.5 \end{pmatrix}^{-1} \begin{pmatrix} 0.008789 \\ 0.024414 \end{pmatrix},$$

$$= \begin{pmatrix} 1.900691 \\ 0.31213 \end{pmatrix}$$

→ Direct method

$$-\begin{pmatrix} f_1(2, 0.25) \\ f_2(2, 0.25) \end{pmatrix} = \begin{pmatrix} 2 & -1 \\ 4 & 2 \end{pmatrix} \begin{pmatrix} \Delta x \\ \Delta y \end{pmatrix}$$

$$-\begin{pmatrix} 0.25 \\ 0.25 \end{pmatrix} = \begin{pmatrix} 2 & -1 \\ 4 & 2 \end{pmatrix} \begin{pmatrix} \Delta x \\ \Delta y \end{pmatrix}.$$

$$\Delta x = \frac{\begin{vmatrix} -0.25 & -1 \\ -0.25 & 2 \end{vmatrix}}{\begin{vmatrix} 2 & -1 \\ 4 & 2 \end{vmatrix}} = -\frac{0.75}{8} = -0.09375$$

$$\begin{aligned} P_1 &= \Delta x + P_0 \\ &= -0.09375 + 2 \\ &= 1.90625. \end{aligned}$$

$$\Delta y = \frac{\begin{vmatrix} -0.25 & 2 & -0.25 \\ -0.25 & 4 & -0.25 \end{vmatrix}}{8} = -\frac{0.5+1}{8} = \frac{0.5}{8} = 0.0625$$

$$\begin{aligned} \Delta y &= g_1 + g_0 \\ g_1 &= \Delta y + g_0 \\ &= 0.0625 + 0.25 \\ &= 0.3125. \end{aligned}$$

discussion

2.4

□ $f(x) = (x-p)^m h(x)$.

$$\Leftrightarrow f(p)=0, f'(p)=0 \dots f^{(m-1)}(p)=0 \quad \text{but} \quad f^{(m)}(p) \neq 0.$$

$f(p)=0$.

$$f'(x) = m(x-p)^{m-1} h(x) + (x-p)^m h'(x).$$

$$f'(p)=0.$$

$$f''(p)=0.$$

$(x-p)$ is a factor of $f(x)$.

$(x-p)^2$ is a factor of $f'(p)$.

8

$$g(x) = x - \frac{mf(x)}{f'(x)} \quad \text{it will converge quadratically to } p.$$

p is a root of multiplicity m for $f(x)$.

$$g'(p)=0 \quad \text{اى تسلیم}$$

$$f(x) = (x-p)^m h(x), h(p) \neq 0.$$

$$f'(x) = m(x-p)^{m-1} h(x) + (x-p)^m h'(x).$$

$$g(x) = x - \frac{m(x-p)^m h(x)}{m(x-p)^{m-1} h(x) + (x-p)^m h'(x)}$$

$$= x - \frac{m(x-p) h(x)}{m h(x) + (x-p) h'(x)}$$

$$g'(x) = 1 - \frac{(m h(x) + (x-p) h'(x))(m h(x) + (x-p) h'(x)) - m(x-p) h(x) \cancel{x^2}}{[m h(x) + (x-p) h'(x)]^2}$$

$$g'(p) = 1 - \frac{(m h(p))^2}{m h(p)^2}$$

$$= 0.$$