

Ex Solve the equation $x = e^{-x} + 1$ with $P_0 = 1.5$ using Newton's method. Find five iterations.

Solution $f(x) = x - e^{-x} - 1$, $P_0 = 1.5$

$$f'(x) = 1 + e^{-x}$$

$$P_{n+1} = P_n - \frac{P_n - e^{-P_n} - 1}{1 + e^{-P_n}}$$

1.5 = Ans.

$$P_0 = 1.5, P_1 = 1.273638286 \Rightarrow 0.23$$

$$P_2 = 1.278462001 \Rightarrow 0.003$$


$$P_3 = 1.278464543 \Rightarrow 0.000002$$

$$P_4 = 1.278464543 \Rightarrow 0$$

$$P_5 = 1.278464543 \Rightarrow 0$$

[5] The Secant Method

$$f(x) = 0, P_0, P_1$$



$$P_2 = P_1 - \frac{f(P_1)(P_1 - P_0)}{f(P_1) - f(P_0)}$$

2 condition ←

$$P_3 = P_2 - \frac{f(P_2)(P_2 - P_1)}{f(P_2) - f(P_1)}$$

$$P_{n+1} = P_n - \frac{f(P_n)(P_n - P_{n-1})}{f(P_n) - f(P_{n-1})}$$

Ex Estimate $\sqrt[3]{20}$ using $P_0 = 2$, $P_1 = 3$
 using a secant method. Find two Iteration
 $P_2 = ??$ اكتب قيم P_0, P_1

Solution $f(x) = x^3 - 20$, $P_2 = 3 - \frac{f(3)(3-2)}{f(3)-f(2)} = 3 - \frac{7}{7-12} = 2.63157$


$P_3 = 2.63157 - \frac{f(2.63157)(2.63157-3)}{f(2.63157)-f(3)}$

في هذه الطريقة ممكن ان يكون الجواب سالب

Q Find the Point on $y = x^2$ that is the closet to Point $(3, 1)$

في هذه المسئلة
استخدم شريطة

x x
3 1



Distance = $\sqrt{(x-\hat{x})^2 + (y-\hat{y})^2}$
 $\downarrow \downarrow \downarrow$
 $(x-3)^2 + (x^2-1)^2$

Distance = $\sqrt{(x-3)^2 + (x^2-1)^2}$

$d^2 = (x-3)^2 + (x^2-1)^2$

$2dd' = 2(x-3) + 2(x-1)(2x) = 0$
 $f(x)$

$f(0) =$

$f(1) = > 0$

$f(2) = < 0$

$\frac{1+2}{2} = 1.5$

* Multiplicity of roots

$$f(x) = (x-2)^3 (x+1)$$

Root : $p=2$
 $p=-1$

Def Let p be a root for $f(x)$

$$\text{if } f(p) = f'(p) = f''(p) = \dots = f^{(M-1)}(p) = 0$$

$$\text{but } f^{(M)}(p) \neq 0$$

then we say p has Multiplicity M .

Another Def Let p be a root for $f(x)$, we say that p

has Multiplicity M if we can write $f(x)$ as :

$$f(x) = (x-p)^M \cdot h(x) ; h(p) \neq 0 \quad \text{Mult.} \quad \text{step}$$

Ex : $f(x) = (x-2)^3 (x+1)$ $p=2, p=-1$

Find Multiplicity for each root

Sol $p=2 : f(x) = (x-2)^3 (x+1), h(x) = x+1$
 $h(2) = 3 \neq 0$
 $M=3$

or $f(2) = 0 : f'(x) = (x-2)^3 + 3(x-2)^2 (x+1)$
 $f'(2) = 0$
 $f''(x) = 3(x-2)^2 + 3(x-2)^2 + (x+1) \cdot 6(x-2)$
 $f''(2) \neq 0 ?$
 $f'''(x) =$

$$f'''(2) \neq 0 \Rightarrow M=3$$

$$P = -1 \quad f(x) = (x+1)(x-2)^3$$

$$(x-2)^3 \text{ has } h(x)$$

$$h(-1) \neq 0 \Rightarrow M=1$$

or $f(-1) = 0$

$$f'(-1) = 27 \neq 0$$

لا يكون صفر

Note $f(x) = \ln x$

$P=1$

$h(1) = 0 \Rightarrow$ so differ. it

$$df = \ln x + \frac{x-1}{x} + 1 = \frac{1}{x}$$

$$ddf = \frac{1}{x} + \frac{1}{x^2} \neq 0 \Rightarrow M=2$$

* Notations * P has Multiplicity = 1 $\Rightarrow$ P is called simple root

* P has $M > 1 \Rightarrow$ P is called Multiple root

$M=2 \Rightarrow$ double root

$M=3 \Rightarrow$ cubic root

Ex $f(x) = (x-1) \ln x$, find M for root

Sol $P=1$ root

$$f'(x) = \frac{x-1}{x} + \ln x \Rightarrow f'(1) = 0$$

$$f''(x) = \frac{x-x+1}{x^2} + \frac{1}{x} \Rightarrow f''(1) = 2 \neq 0$$

$\Rightarrow M=2$, $P=1$ is double root

Ex $f(x) = (x-1)^3 \ln x$, $P=1$ is cubic root $M=3$

* $f(x) = x^3 - 70$
 $f'(x) = 3x^2 \neq 0 \Rightarrow$ simple root

* Order of Convergence: is the number that Measures the speed of convergence of any numerical method (any sequence P_n)
 $R > 0$ يعبر عن

* Notation sequence $P_n \Big|_{n=0}^{\infty} \rightarrow P_{\text{exact root}}$

$$|E_n| = |P - P_n| \quad \text{الخطأ الحقيقية - التقريب}$$

$$|E_{n+1}| = |P - P_{n+1}|$$

Def Let $\{P_n\}$ be a sequence converge to P then

if $\lim_{n \rightarrow \infty} \frac{|E_{n+1}|}{|E_n|^R} = A$, then we say the order of convergence

is $R > 0$, where A is called the asymptotic error constant ($A > 0$)

Note Find Order of Convergence $\Rightarrow P$ & A

Def

Sequence $[P_n]_{n=0}^{\infty} \rightarrow P$

1-3-2018

$$|E_n| = |P - P_n|$$

$$\lim_{n \rightarrow \infty} \frac{|E_{n+1}|}{|E_n|^R} = A, R, A > 0$$

تقريب السرعة

we say P_n converge to P with order of convergence = R

* A: asymptotic error constant

For n large $\frac{|E_{n+1}|}{|E_n|^R} \approx A$

$$\Rightarrow |E_{n+1}| \approx A |E_n|^R$$

constant

I claim: $R \uparrow$, convergence is faster.

Method 1

Method 2

فهي $R=1$

$R=2$

R أكبر < أسرع

* Suppose $|E_n| \approx 0.1$

$$|E_{n+1}| \approx A |E_n|^1 \approx A(0.1)$$

$$|E_{n+1}| \approx A |E_n|^2 = A(0.1)^2 = 0.01$$

السرعة تقل ، الكونفرينس قريب ، السرعة تزيد R & A

Ex Find the order of convergence of the sequence $= \frac{1}{6^n}$

Solution of $\frac{1}{6} \leftarrow \frac{1}{36} \leftarrow \frac{1}{216} \leftarrow \dots$, $P_n = \frac{1}{6^n}$

$P = 0$ اللت تبعد على

$$\lim_{n \rightarrow \infty} \frac{|E_{n+1}|}{|E_n|^R} = \lim_{n \rightarrow \infty} \frac{|P_{n+1} - P|}{|P_n - P|^R}$$

$$= \lim_{n \rightarrow \infty} \frac{\left| \frac{1}{6^{n+1}} - 0 \right|}{\left| \frac{1}{6^n} - 0 \right|^R}$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{\frac{1}{6^{n+1}}}{\frac{1}{6^{nR}}} = \lim_{n \rightarrow \infty} \frac{6^{nR}}{6^{n+1}} = \lim_{n \rightarrow \infty} 6^{nR-n-1}$$

$\frac{6^R}{6} = 6^{R-1}$

$$\Rightarrow \lim_{n \rightarrow \infty} 6^{-1} \cdot 6^{n(R-1)} = \left[\frac{1}{6} \right] \lim_{n \rightarrow \infty} 6^{n(R-1)}$$

جواب

$$= \frac{1}{6} \begin{cases} \infty, & R > 1 & \times \\ 0, & R < 1 & \times \\ \frac{1}{6}, & R = 1 & \checkmark \end{cases}$$

مقدار

$$\Rightarrow R = 1, A = \frac{1}{6}$$

* Notes $R = 1 \Rightarrow$ Convergence is linear

$R = 2 \Rightarrow$ = = Quadratic عمر

$R = 3 \Rightarrow$ = = Cubic

نظریه Newton Method, $P_{n+1} = P_n - \frac{f(P_n)}{f'(P_n)}$ for $f(x)$

assume $\{P_n\} \rightarrow P$

1 if P is simple root ($M = 1$), then $R = 2$
 $A = \left| \frac{f''(P)}{2f'(P)} \right|$

(i.e. : $\lim_{n \rightarrow \infty} \frac{|E_{n+1}|}{|E_n|^2} = \left| \frac{f''(p)}{2f'(p)} \right|$)

② if p is a multiple root $(M > 1)$
 then $R = 1$, $A = \frac{M-1}{M}$

simple $\Rightarrow$ بسيطة
 multiple $\Rightarrow$ متعددة

(i.e. $\lim_{n \rightarrow \infty} \frac{|E_{n+1}|}{|E_n|} = \frac{M-1}{M}$)

* Secant Method : $P_{n+1} = P_n - \frac{f(P_n)(P_n - P_{n-1})}{f(P_n) - f(P_{n-1})}$

① if p simple root

$R = 1.618$, $A = \left| \frac{f''(p)}{2f'(p)} \right|^{0.618}$

② if p multiple root

$R = 1$, A depend on M

* Bisection

$R = 1$, $A = 0.5$ always

طبيقة

* False-Position

$R = 1$, A depend on $f(x)$

لا يتغير على الدوام

* FPT :

depends on $g(x)$

Ex $P(\text{known})$ في هذا المثال

$$f(x) = (x+2)(x-1)^2$$

$$P = -2 \Rightarrow M = 1$$

$$P = 1 \Rightarrow M = 2$$

1 Newton Find R, A theoretically for each root

$$[P = -2] : R = 2, A = \left| \frac{f'(-2)}{2f'(-2)} \right|$$

$$f(x) = (x+2)(x^2-2x+1) = x^3 - 3x + 2$$

$$f'(x) = 3x^2 - 3$$

$$f''(x) = 6x$$

$$\lim_{n \rightarrow \infty} \frac{|E_{n+1}|}{|E_n|^2} = \frac{2}{3} = A$$

$$\Rightarrow A = \left| \frac{f'(-2)}{2f'(-2)} \right| = \frac{12}{18} = \frac{2}{3} = A$$

$$[P = 1] : R = 1, A = \frac{M-1}{M} = \frac{2-1}{2} = \frac{1}{2}$$

$$\lim_{n \rightarrow \infty} \frac{|E_{n+1}|}{|E_n|} = \frac{1}{2} = 0.5$$

2 R, A, Secant

$$[P = -2] : R = 1.618, A = \left| \frac{f''(-2)}{2f'(-2)} \right| = 0.618$$

$$\lim_{n \rightarrow \infty} \frac{|E_{n+1}|}{|E_n|^{1.618}} = 0.77835$$

نظرياً لا يتوقف $P=1$, $R=1$, A

3 Bisection , $P, A \Rightarrow P=-2$, $P=1$
 $R=1$, $A=0.5$

$$\lim_{n \rightarrow \infty} \frac{|E_{n+1}|}{|E_n|} = 0.5$$

4 False-Position : $R=1$, $A=?$

Ex $f(x) = x^3 - 27$

6-3-2018

Solution $P = 3$ (Known)

أولاً P $\rightarrow$ 3

✂

① Find R, A if we used Newton method (theoretically)

② Start with 3.5 using three iteration of Newton

Find R, A numerically

① $f'(x) = 3x^2$

$f'(3) = 27 \neq 0$

$\Rightarrow M=1$ simple root

$R=2, A = \left| \frac{f''(3)}{2f'(3)} \right| = \left| \frac{18}{2(27)} \right| = \frac{18}{54} = \frac{1}{3} = 0.333$

* this means $\lim_{n \rightarrow \infty} \frac{|E_{n+1}|}{|E_n|^2} = \frac{1}{3}$

② $P_{n+1} = P_n - \frac{f(P_n)}{f'(P_n)}$ P is known

$\frac{|E_{k+1}|}{|E_k|} = A$

$\frac{P - P_{n+1}}{(P - P_n)^R}$

P_k	$ E_k = 3 - P_k $	$\frac{ E_{k+1} }{ E_k }$
3.5	0.5	0.272108
3.068027211	0.068027211	0.323533
3.001497216	0.001497216	0.332355
3.000000747	0.000000747	

$0.068027211 / 0.5^2$

$\downarrow \frac{1}{3}$

Ex $f(x) = (x-2)^2(x+1)$

Solution Root : $P=2, M=2, P=-1, M=1$

$R=1$

$$f(x) = x^3 - 3x^2 + 4$$

$$f'(x) = 3x^2 - 6x$$

$$f''(x) = 6x - 6$$

<u>Newton</u>	$P=2$	$E_k = P - P_k $	<u>Newton</u>	$P=-1$
$P_0 = 2.5$		0.3	$P_0 = -1.5$	
$P_1 = 2.26667$		0.26662	$P_1 = -1.111$	
$P_2 = 2.13856$		0.13056	$P_2 = -1.00740$	
$P_3 = 2.070777$			$P_3 = -1.000036311$	
$P_4 = 2.03579$			$P_4 = -1.0000000001$	
$P_5 = 2.0180008$			$P_5 = -1$	
$\vdots$				
$P_{25} = 2.000000547$				

* Accelerated Newton Iteration

Newton for Multiple roots is slow $R=1$

P root $M > 1$

$$P_{n+1} = P_n - \frac{M f(P_n)}{f'(P_n)} \quad \text{Accelerated Newton}$$

$R=2$

Ex For previous examples $P=2$, $R=1$, $M=2$

use accelerated Newton method

$$P_{n+1} = P_n - \frac{2(P_n^3 - 3P_n^2 + 4)}{3P_n^2 - 6P_n}$$

$$P_0 = 2.5$$

$$R=2 \text{ في هذه الطريقة}$$

$$P_1 = 2.0333$$

$$P_2 = 2.000182 \quad [R=2]$$

$$P_3 = 2.0000005$$

Find A theo. ①

$$\frac{1}{1} \frac{1}{1^2}$$

②

* FPI, R, A

Theorem Let P be a fixed pt for $g(x)$

$$\text{If } g'(P) = g''(P) = \dots = g^{(n-1)}(P) = 0$$

$$\text{but } g^{(n)}(P) \neq 0$$

Multiplicity $\Rightarrow$ لا تكون له $\Rightarrow$ نفس root

then FPI will converge to P with $\underline{R=n}$, $A = \left| \frac{g^{(n)}(P)}{n!} \right|$

$\Rightarrow$ * Theoretically $\Leftarrow$

Ex $g(x) = \frac{x}{2} + \frac{2}{x}$ $P=2$ Fixed Point

① Find R, A, Theoretically if we used FPI

② Prove your claim in (a) using $P_0 = 2.5$ & three Iteration of FPI

$$x = \frac{x}{2} + \frac{2}{x} \Rightarrow \text{f.p.} =$$

Solution $g'(x) = \frac{1}{2} - \frac{2}{x^2}$

$$g'(2) = \frac{1}{2} - \frac{1}{2} = 0$$

$$g''(x) = \frac{4}{x^3}$$

$$g''(2) \neq 0 = \frac{4}{2^3} = \frac{1}{2}$$

$$P_{k+1} = \frac{Ans}{2} + \frac{2}{Ans}$$

$R=n$
FPI

$$\boxed{R=2}, A = \left| \frac{g''(2)}{2!} \right| = 0.25$$

P_k	$P_k - P_{k-1}$	$ E_k $ ← <u>مقدار الخطأ</u>	$\frac{ E_{k+1} }{ E_k ^2}$
2.5		0.5	0.20
2.05		0.05	0.2439
2.000609756		0.000609756	0.25013
2.000000093		0.000000093	↓ 0.25 = $\frac{1}{4}$

Proof need to show $\lim_{k \rightarrow \infty} \frac{|E_{k+1}|}{|E_k|^n} = \left| \frac{g^{(n)}(P)}{n!} \right|$
 $P_{k+1} = g(P_k)$

Apply Taylor's Exp. for $g(x)$ about P

$$g(x) = g(P) + g'(P)(x-P) + \frac{g''(P)}{2}(x-P)^2 + \dots + \frac{g^{(n-1)}(P)}{(n-1)!}(x-P)^{n-1} + \frac{g^{(n)}(c)}{n!}(x-P)^n$$

$$g(x) = P + \frac{g^{(n)}(c)}{n!}(x-P)^n$$

$$\boxed{x = P_k} : g(P_k) = P + \frac{g^{(n)}(c)}{n!}(P_k - P)^n$$

$$P_{k+1} - P = \frac{g^{(n)}(c)}{n!}(P_k - P)^n \quad c \text{ between } P \text{ \& } P_k$$

$$\left| \frac{P_{k+1} - P}{(P_k - P)^n} \right| = \left| \frac{g^{(n)}(c)}{n!} \right|$$

$$\Rightarrow \lim_{k \rightarrow \infty} \frac{|E_{k+1}|}{|E_k|^n} = \left| \frac{g^{(n)}(P)}{n!} \right|$$

$$P_k \approx P$$

$$P_{n+1} = P_n - \frac{f(P_n)}{f'(P_n)}$$

Q Prove that if P is a simple root for $f(x)$, then Newton iteration will converge to P with $R=2$, $A = \left| \frac{f''(P)}{2f'(P)} \right|$
 $M=1 \Rightarrow f'(P) \neq 0$
 $\text{or } g'(P) = 0$

Proofs Apply Taylor Exp. for $f(x)$ about $P_n \Rightarrow \text{center}$

$$f(x) = f(P_n) + f'(P_n)(x - P_n) + \frac{f''(c)(x - P_n)^2}{2}$$

$$\text{Let } x = P \quad f(P) = f(P_n) + f'(P_n)(P - P_n) + \frac{f''(c)(P - P_n)^2}{2}$$

Divide by $f'(P_n)$ (قسمة الكل بـ $f'(P_n)$)

root P is $f(P) = 0$
 $f(P_n) \rightarrow 0$

$$0 = \frac{f(P_n)}{f'(P_n)} + \frac{P - P_n + \frac{f''(c)}{2f'(P_n)}(P - P_n)^2}{1}$$


$$P - (P_n - \frac{f(P_n)}{f'(P_n)}) = -\frac{f''(c)}{2f'(P_n)}(P - P_n)^2$$

$$|P - P_{n+1}| = \left| \frac{-f''(c)}{2f'(P_n)} \right| |P - P_n|^2$$

$$\frac{|E_{n+1}|}{|E_n|^2} = \left| \frac{f''(c)}{2f'(P_n)} \right|$$

$$\lim_{n \rightarrow \infty} \frac{|E_{n+1}|}{|E_n|^2} = \left| \frac{f''(P)}{2f'(P)} \right|$$

Since when $n \rightarrow \infty$ $c \approx P_n \approx P$



Q Prove that if P is a multiple root for $f(x)$ ($m > 1$) then Newton has $R=1$, $A = \frac{m-1}{m}$

Hint Apply Taylor about $x = P$ for $f(x)$
 center $\Rightarrow$ not x

$$f(x) = f(p) + f'(p)(x-p) + \dots + \frac{f^{(M-1)}(p)}{(M-1)!} (x-p)^{M-1} + \frac{f^{(M)}(p)}{M!} (x-p)^M$$

$$f(x) = \frac{f^{(M)}(p)}{M!} (x-p)^M \Rightarrow f'(x) = \frac{M f^{(M)}(p) (x-p)^{M-1}}{M(M-1)!}$$

مثال

$$f'(x) = \frac{f^{(M)}(p)}{(M-1)!} \quad (M! = M(M-1)!)$$

نفس

$$\frac{f(x)}{f'(x)} = \frac{x-p}{M} \Rightarrow \frac{f(p)}{f'(p)} = \frac{p-p}{M} \Rightarrow$$

↓

$$\frac{\frac{f^{(M)}(p)}{M!} (x-p)^M}{\frac{f^{(M)}(p)}{(M-1)!} (x-p)^{M-1}} = \frac{(x-p)^{M-(M-1)}}{M} = \frac{f(x)}{f'(x)}$$

let $x = p_n \Rightarrow \frac{f(p_n)}{f'(p_n)} = \frac{p_n - p}{M}$

للقرينة

$$\frac{p_n}{f'(p_n)} - \frac{f(p_n)}{f'(p_n)} = p_n - \frac{p_n - p}{M}$$

$$-p \quad p_{n+1} = p_n - \frac{p_n - p}{M}$$

$$\frac{p - p_{n+1}}{p - p_n} = 1 - \frac{1}{M} = \frac{M-1}{M}$$

$$\frac{|E_{n+1}|}{|E_n|} = \frac{M-1}{M} \quad \text{Constant}$$

$$\lim \frac{|E_{n+1}|}{|E_n|} = \frac{M-1}{M} \quad \text{Constant}$$