



Student's Name : Ayham Hashesh

Instructor : Mahamoud Ghannam

Numerical Methods

طرق التحليل العددي



BITRZET UNIVERSITY


1161301

أيهام حشيش

* Chapter 1 :-

1.1 Review of calculus

* Continuity : f is continuous on $[a, b]$




* $\forall$ for all

$$f(x) \in C[a, b]$$

$$f(x) \in C'[a, b]$$


* f is increasing & continuous on $[a, b]$

$$f'(x) > 0$$



* f is decreasing & continuous on $[a, b]$

$$f'(x) < 0$$



* Theorem if f is continuous on $[a, b]$ then

f has an absolute Max (upper bound) = M

& f has an absolute min (lower bound) = m

$$\Rightarrow m \leq f(x) \leq M \quad \forall x \in [a, b]$$


من الأمثلة أو النطاق الحرجة

المشتقة الأولى

max على

min على

Ex $f(x) = x^2 - 8x + 3$, $[2, 6]$ find M, m



$$f(2) = -9 \quad M$$


$$f(6) = -9 \quad M$$

$$f'(x) = 2x - 8 = 0$$

$$\boxed{x=4} \Rightarrow f(4) = -13 \quad m$$

$x \in [2, 6]$

Ex $f(x) = \sin x$, $[0, 1]$



$$M = f(1) = \sin 1$$

$$m = f(0) = \sin 0 = 0$$

$$0 \leq f(x) \leq \sin 1$$


$x \in [0, 1]$

Ex $f(x) = \frac{2}{(x+1)}$, $1 \leq x \leq 4$


$\Rightarrow$ decreasing

$$\Rightarrow M = f(1) = \frac{2}{64} = \frac{1}{32}$$

$$m = f(4) = \frac{2}{7^3}$$



* Intermediate Value Theorem [IVT]




$f \in C[a, b]$ & L
between $f(a)$, $f(b)$

$$\Rightarrow \exists c \text{ in } [a, b] \text{ s.t. } f(c) = L$$

* Bolzano Theorem ^{صيغة أويلر} ^{موجب} ^{والأخر سالب}

$$f \in C[a, b] \text{ & } f(a)f(b) < 0$$

$$\Rightarrow \exists c \in (a, b) \text{ s.t. } f(c) = 0$$



$x = c$ is a root (zero) for $f(x)$

$x = c$ = = solution for the equation $\boxed{f(x) = 0}$


* Mean Value Theorem [MVT]

$$\Rightarrow f \in C[a, b] \text{ & } f' \in C[a, b]$$

$$\Rightarrow \exists c \in (a, b) \text{ s.t. :}$$

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$|f(b) - f(a)| = |f'(c)| |b - a|$$



Recall : $|x \mp y| \leq |x| + |y|$

* Rolle's Theorem

f cont on $[a, b]$ & differentiable on (a, b) & $f(a) = f(b)$

$\Rightarrow c \in (a, b)$ s.t. $f'(c) = 0$

* Taylor Series [Taylor expansion]

$$[1] \quad e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots$$

$e^x = 1 + x + \frac{x^2}{2}$, for x close to zero center

$$[2] \quad \sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$$

$$[3] \quad \cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$$

given $f(x)$ & a center $x=a$, The Taylor expansion (series) for $f(x)$ about $x=a$ is given by

$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)(x-a)^2}{2!} + \frac{f'''(a)(x-a)^3}{3!} + \dots$$

$\Rightarrow$ used to approximate $f(x)$ for x close to a

* Taylor Theorem

Taylor expansion for $f(x)$ about $x=a$ &

$$f(x) \approx f(a) + f'(a)(x-a) + \frac{f''(a)(x-a)^2}{2!} + \dots + \frac{f^{(n)}(a)(x-a)^n}{n!}$$

the error of this equation is

$$E(x) = \frac{f^{(n+1)}(c)(x-a)^{n+1}}{(n+1)!}, \text{ where } c \text{ is a constant between } x \text{ and } a$$

$f(x)$ $P_3(x)$ Ex

$$e^x = \left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} \right)$$

[1] Find Error

$$E = \frac{f^{(4)}(c)}{24} x^4$$

$$f(x) = e^x = f'(x) = f''(x) = f'''(x)$$

$$E = \frac{e^c}{24} x^4, \quad c \text{ between } x \text{ \& } 0$$

[2] Find an upper bound for |Error| when approximately $e^{0.5}$

$$\text{Now } e^{0.5} \approx 1 + 0.5 + \frac{0.5^2}{2} + \frac{0.5^3}{6}$$

$$E(x) = \frac{e^c}{24} x^4$$

c between 0.5 & 0

$$0 < c < 0.5$$

$$1 = e^0 < e^c < e^{0.5}$$

$$|E(0.5)| = \frac{e^c (0.5)^4}{24} \leq \frac{e^{0.5} (0.5)^4}{24}$$

في الامتحان المثل جواب رقم

1.3 Error

13-2-2018

If P is approximated by $\hat{P}$ then

① absolute error $|E| = |P - \hat{P}|$

القيمة الحقيقية
القيمة التقريبية

② Relative error $= |R| = \frac{|P - \hat{P}|}{|P|}$

Ex $P = 1.2351$, $\hat{P} = 1.2$

Find $|E|$, $|R|$ Solution $|E| = |1.2351 - 1.2|$
 $= 0.0351$

$$|R| = \frac{0.0351}{1.2351} = 0.0284$$

Ex $999\ 995 = P$, $1\ 000\ 000 = \hat{P}$

Solution $|E| = 5$ $|R| = \frac{5}{999\ 995} = 0.000\ 005\ 000\ 025$

abs $|E|$ precision just Relative

Recall $e^x \approx \underbrace{1 + x + \frac{x^2}{2} + \frac{x^3}{6}}_{P_3(x)}$

$$|E| = \left| e^x - \left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} \right) \right|$$

* Source of Error

① Round-off error : a number is approximated by a number

② Truncation error : a formula is estimated by a formula

Round-off Error

* Finite digit Arithmetic

⇒ Computer has a Finite numbers of digit

* Floating Point Representation

$$\begin{aligned} 0.02341 &= 0.2341 * 10^{-1} \\ &= 2.341 * 10^{-2} \\ &= 2341.0 * 10^{-5} \end{aligned}$$

* Normalized Floating Point

any number is represented by $\underbrace{0.d_1d_2d_3\dots d_k}_{\text{significant digit}} * 10^n$

where $d_1 \neq 0$

$$0.000000012 \Rightarrow \underbrace{0.12}_{\text{significant digit}} * 10^{-7} \quad [\text{Normalized}]$$

or [ناتج ابدأ من اليسار حتى نجد رقم غير صفر]

$$\text{Ex } 0.0100234 * 10^{-4} \Rightarrow 6 \text{ significant digit}$$

$$1.200 \Rightarrow 4 \text{ significant digit}$$

* Rule : Start Counting from left from the first non-zero digit حالة من اليسار

$$0.000000012$$

الأرقام الحاسبة تبدأ من اليمين
10 حانات من اليمين

* Two Types of Round-off : [1] Chopping : $0.d_1d_2\dots d_kd_{k+1}d_{k+2}\dots$
 $\approx 0.d_1d_2\dots d_k$

[2] Rounding : $0.d_1d_2\dots d_kd_{k+1}d_{k+2}\dots$
 $\approx 0.d_1d_2\dots r_k$

$\begin{cases} d_k & \text{if } d_{k+1} < 5 \\ d_{k+1} & \text{if } d_{k+1} > 5 \end{cases}$

* We Assume that Both [chopping & Rounding] are Finite digit Arithmetic $\Leftarrow$ [significant digit]

Ex Using 5 digit arithmetic estimate

[1] $2.0147985 \Rightarrow$ [a] chopping $\Rightarrow 2.0147$
 [b] Rounding $\Rightarrow 2.0148$

[2] $0.000314295 \Rightarrow$ [a] chopping $\Rightarrow 0.00031429$
 [b] Rounding $\Rightarrow 0.00031430$

Ex Estimate $\frac{1}{7} + \frac{1}{8} + \frac{1}{9}$ using 2-digit rounding

Solution $0.14 + 0.13 + 0.11$
 $\underline{0.27} + 0.11 = 0.38$

تقريب $\frac{1}{7}, \frac{1}{8}, \frac{1}{9}$
 مجموع $[\frac{1}{8}, \frac{1}{7}]$
 التقريب الكلي التالي $\frac{1}{9}$
 التقريب الكلي النهائي

* Order of Operations :

[1] Brackets

[3] $\div, *$ [From Left to Right]

[2] Powers

[4] $+, -$ [= = = =]

Ex Estimate $\frac{\frac{1}{6} \div \frac{1}{3} + (0.23)^3}{\cos e^2 * \sqrt{0.01348}}$ Using 3-digit rounding

Note default in calculator [D] $\Rightarrow$ Degree mode, we use Rad mode always

Solution $\frac{0.167 \div 0.333 + (0.23)^3}{(\cos(2.72)^2) * \sqrt{0.0135}}$

Notes ① $e = 2.72$

② $0.23 \Rightarrow$ لو أكبر من ٣ خانة يجب التقريب

③ $(0.23)^3 = (0.23)(0.23)(0.23)$

كتابة مباشرة

④ لو كانت الجمع قبل القسمة = تقسم الأرقام الممنوعة

$\Rightarrow \frac{0.502 + 0.0122}{\cos(7.46) * 0.116}$

$\Rightarrow \frac{0.514}{0.0509} = 10.1$

6 sig

* Loss of significant :- $P = 0.123479$ $P - q = 0.000001$ [1 sig]
 $q = 0.123478$ Loss = عدد من رقمين في نفس المكان

Ex $\ln(x) = -\ln(x+1) \Rightarrow \text{sol } \ln\left(\frac{x}{x+1}\right)$ [L.O.S] في النهاية

* Propagation of error :- error may increase due to operation
 يمكن أن تزداد

Ex $P = \hat{P} + E_p$ $q = \hat{q} + E_q$ $\nearrow$ التقريب

$P + q = \hat{P} + \hat{q} + E_p + E_q$

* Order of Approximation :- Notation or Method used to express the truncation error for some estimation

Note When I use ch instead of x , this means h is a variable close to 0 $\Rightarrow |h| < 1$

ex $e^h = 1 + h + \frac{h^2}{2} + \dots$ $\Rightarrow E = \frac{f^{(3)}(c)}{3!} h^3 = \frac{e^c h^3}{6}$

C is constant between 0 & h [truncation Error] $E \approx ch^3$

We can express this error by $O(h^3)$ order of approximation

Def : Order of approximation if $f(h) = p(h) + O(h^n)$ then $f(h) \approx p(h)$
 with $E_{\text{trunc}} \approx C * h^n$

Ex $\sin h \approx h - \frac{h^3}{3!} + \frac{h^5}{5!} \Rightarrow$ Order of approximation : $O(h^7)$

* Order of approximation :

15-2-2018

$$f(h) = p(h) + o(h^n)$$

$$f(h) \approx p(h)$$

$$E \approx C \cdot h^n$$

$$f(h) = p(h) + o(h^5)$$

$\bar{p}(h)$ is a better approximation for $f(h)$

$$f(h) = \bar{p}(h) + o(h^6)$$

Ex $\sinh h \approx h - \frac{h^3}{3!} + \frac{h^5}{5!}$, what is the order of approx ?

$$E = \frac{f^{(7)}(c)}{7!} h^7 = \frac{\cosh c}{7!} h^7 = C \cdot h^7$$

order of approx : $o(h^7)$

Theorem : $f(h) = p(h) + o(h^n)$

$g(h) = q(h) + o(h^m)$ then

$$f(h) \pm g(h) = p(h) \pm q(h) + o(h^r) \quad \text{where}$$

$$① r = \min \{m, n\}$$

$$1 \approx 1 + 0.1 \quad \text{لأن}$$

$$② h^K + o(h^r) = o(h^r) \quad \text{if } K \geq r$$

فإن h^K أكبر أو تساوي h^r لأن

Ex $f(h) = 1 - 2h + h^2 + o(h^3)$
 $g(h) = h^2 + 2h^4 + o(h^5)$

Find ① $f(h) + g(h)$

$$① \Rightarrow 1 - 2h + h^2 + h^2 + 2h^4 + o(h^3)$$

$$1 - 2h + 2h^2 + o(h^3)$$

$$\textcircled{2} f(h) \cdot g(h) = (1 - 2h + h^2)(h^2 + 2h^4) + o(h^3)$$

$$= h^2 + o(h^3)$$

$\textcircled{3}$ estimate $f(0.1)g(0.1)$ and find error

$$f(0.1)g(0.1) \approx (0.1)^2 = 0.01$$

$$E \approx C \propto h^3$$

$$E = C (0.1)^3 = C \propto 0.001$$