

## Actrapid®

100 IU/ml Solution for injection in vial Insulin human (rDNA)

Read all of this leaflet carefully before you start using your insulin.

- Keep this leaflet. You may need to read it again.

- If you have any further questions, ask your doctor, nurse or your pharmacist.

This medicine has been prescribed for you.
 Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.

 If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor, nurse or your pharmacist.

# 1. What Actrapid® is and what it is used for

Actrapid® is human insulin used to treat diabetes. Diabetes mellitus is a disease where your body does not produce enough insulin to control the level of your blood sugar. Actrapid® is a fast-acting insulin. This means that it will start to lower your blood sugar about ½ hour after you take it, and the effect will last for approximately 8 hours. Actrapid® is often given in combination with longer-acting insulin products.

### 2. Befo

## Do not I

- human ir ingredier informat.
- blood su in an em
- ► If the pr Each vial plastic ca when yo your pha
- ▶ If it has been fro
- ► If the in colourle

#### Before u

- ► Check tl right typ
- ► Remove

#### Take spe

- If you have liver, or we thyroid o
- If you d
  a hypo a
  empty st
- ➤ If you e want to affect yo
- ► If you a and con



# Actrapid®

100 IU/ml Solution for injection in vial Insulin human (rDNA)

Read all of this leaflet carefully before you start using your insulin.

- Keep this leaflet. You may need to read it again.

- If you have any further questions, ask your doctor, nurse or your pharmacist.

This medicine has been prescribed for you.
 Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.

 If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor, nurse or your pharmacist.

# 1. What Actrapid® is and what it is used for

Actrapid® is human insulin used to treat diabetes. Diabetes mellitus is a disease where your body does not produce enough insulin to control the level of your blood sugar. Actrapid® is a fast-acting insulin. This means that it will start to lower your blood sugar about ½ hour after you take it, and the effect will last for approximately 8 hours. Actrapid® is often given in combination with longeracting insulin products.

### 2. Befo

# Do not I

- human in ingredier informat
- blood su in an em
- ► If the pr Each vial plastic ca when yo your pha
- ▶ If it has been fro.
- ► If the in colourle

#### Before u

- ► Check tl right typ
- ► Remove

#### Take spe

- ► If you h liver, or v thyroid g
- ► If you d a hypo a empty st
- ➤ If you e want to affect yo
- ► If you a and con