




ELEVENTH EDITION
CAMPBELL
BIOLOGY
URRY • CAIN • WASSERMAN
MINORSKY • REECE

© 2017 Pearson Education, Inc.

Chapter 10

Photosynthesis

Lecture Presentations by
Nicole Tunbridge and
Kathleen Fitzpatrick

The Process That Feeds the Biosphere

- Plants and other photosynthetic organisms contain organelles called **chloroplasts**
- **Photosynthesis** is the process that converts solar energy into chemical energy within chloroplasts
- Directly or indirectly, photosynthesis nourishes almost the entire living world
- Photosynthesis occurs in *plants*, *algae*, certain other *unicellular eukaryotes*, and some *prokaryotes*

© 2017 Pearson Education, Inc.

- **Autotrophs** are “self-feeders” that sustain themselves without eating anything derived from other organisms
- Autotrophs are the **producers** of the biosphere, producing *organic molecules from CO_2 and other inorganic molecules*
- Almost all plants are **photoautotrophs**, using the energy of sunlight to make organic molecules

© 2017 Pearson Education, Inc.

Figure 10.1



© 2017 Pearson Education, Inc.

Figure 10.1a



Other organisms also benefit from photosynthesis.

© 2017 Pearson Education, Inc.

Figure 10.2

Examples of photoautotrophs



(a) Plants



(b) Multicellular alga




10 μm

(c) Unicellular eukaryotes



(d) Cyanobacteria 40 μm



1 μm

(e) Purple sulfur bacteria

- **Heterotrophs** obtain organic material from *other organisms*
- Heterotrophs are the **consumers** of the biosphere
- Some eat other living organisms; others, called **decomposers**, consume *dead organic material or feces*
- Almost all heterotrophs, including humans, depend on photoautotrophs for food and O₂

© 2017 Pearson Education, Inc.

Concept 10.1: Photosynthesis converts light energy to the chemical energy of food

- Chloroplasts are *structurally similar* to, and likely evolved from, *photosynthetic bacteria*
- The structural organization of these organelles allows for the chemical reactions of photosynthesis

© 2017 Pearson Education, Inc.

Chloroplasts: The Sites of Photosynthesis in Plants


- *Leaves* are the major locations of photosynthesis in plants
- Chloroplasts are found mainly in cells of the **mesophyll**, *the interior tissue of the leaf*
- Each mesophyll cell contains *30–40 chloroplasts*
- CO₂ enters and O₂ exits the leaf through *microscopic pores* called **stomata**

© 2017 Pearson Education, Inc.

- A chloroplast has an envelope of two membranes surrounding a dense fluid called the **stroma**
- **Thylakoids** are connected sacs in the chloroplast that compose a third membrane system
- Thylakoids may be stacked in columns called **grana**
- **Chlorophyll**, the pigment that gives leaves their green color, resides in the thylakoid membranes

© 2017 Pearson Education, Inc.

Figure 10.4



© 2017 Pearson Education, Inc.

Photosynthesis: *Splitting of Water*


- Photosynthesis is a complex series of reactions that can be summarized as the following equation:

$$6 \text{ CO}_2 + 12 \text{ H}_2\text{O} + \text{Light energy} \rightarrow \text{C}_6\text{H}_{12}\text{O}_6 + 6 \text{ O}_2 + 6 \text{ H}_2\text{O}$$
- The overall chemical change during photosynthesis is the reverse of the one that occurs during *cellular respiration*
- Chloroplasts split H₂O into **hydrogen** and **oxygen**, incorporating *the electrons of hydrogen into sugar molecules* and *releasing oxygen gas as a by-product*

© 2017 Pearson Education, Inc.

Photosynthesis as a Redox Process



- Photosynthesis *reverses the direction of electron flow compared to respiration*
- Photosynthesis is a redox process in which **H₂O is oxidized** and **CO₂ is reduced**
- Remember:** In respiration, O₂ is reduced to H₂O and sugar molecules are oxidized to CO₂
- Photosynthesis is an **endergonic process**; the energy is provided by light



The Two Stages of Photosynthesis: *A Preview*

- Photosynthesis consists of the **light reactions** (*the photo part*) and **Calvin cycle** (*the synthesis part*)
- The light reactions (in the thylakoids)**
 - Split H₂O
 - Release O₂
 - Reduce the electron acceptor **NADP⁺** to **NADPH**
 - Generate ATP from ADP by **photophosphorylation**


© 2017 Pearson Education, Inc.



- **The Calvin cycle (in the stroma)**
 - Forms sugar from CO_2 , using ATP and NADPH
 - Begins with **carbon fixation**, incorporating CO_2 into organic molecules


© 2017 Pearson Education, Inc.

Figure 10.6_3



© 2017 Pearson Education, Inc.

Figure 10.6_4




Concept 10.2: The light reactions convert solar energy to the chemical energy of ATP and NADPH

- Chloroplasts are solar-powered chemical factories
- Their thylakoids *transform light energy into the chemical energy of ATP and NADPH*

© 2017 Pearson Education, Inc.

The Nature of Sunlight

- Light is *electromagnetic energy*, also called electromagnetic radiation
- Electromagnetic energy travels in rhythmic *waves*
- Light also behaves as if it consists of discrete particles, called *photons*
- **Wavelength** is the distance between crests of electromagnetic waves




© 2017 Pearson Education, Inc.

- The **electromagnetic spectrum** is the entire range of electromagnetic energy, or radiation
- **Visible light** consists of wavelengths (380 nm to 750 nm) that produce colors we can see
- *Visible light also includes the wavelengths that drive photosynthesis*

© 2017 Pearson Education, Inc.

Figure 10.7




© 2017 Pearson Education, Inc.

Photosynthetic Pigments: The Light Receptors

- **Pigments** are substances that absorb visible light
- *Different pigments absorb different wavelengths*
- Wavelengths that are not absorbed are **reflected** or **transmitted**
- *Leaves appear green because chlorophyll reflects and transmits green light*


© 2017 Pearson Education, Inc.

Figure 10.8



© 2017 Pearson Education, Inc.

Animation: Light Energy and Pigments



© 2017 Pearson Education, Inc.


- There are three types of pigments in chloroplasts:
 - **Chlorophyll *a***, the key light-capturing pigment
 - **Chlorophyll *b***, an accessory pigment
 - **Carotenoids**, a separate group of accessory pigments

© 2017 Pearson Education, Inc.

- A **spectrophotometer** measures a pigment's ability to absorb various wavelengths
- This machine sends light through pigments and measures the fraction of light transmitted at each wavelength


© 2017 Pearson Education, Inc.

Figure 10.9



© 2017 Pearson Education, Inc.

- An **absorption spectrum** is a graph plotting a pigment's light absorption versus wavelength




- The absorption spectrum of chlorophyll *a* suggests that violet-blue and red light work best for photosynthesis


© 2017 Pearson Education, Inc.

Figure 10.10


- An **action spectrum** profiles the relative effectiveness of different wavelengths of radiation in driving a process (e.g. photosynthesis)



(a) Absorption spectra



(b) Action spectrum



(c) Engelmann's experiment

Data from T. W. Engelmann, *Bacterium photometricum*. Ein Beitrag zur vergleichenden Physiologie des Licht- und Farbensinnes, *Archiv. für Physiologie* 30:95–124 (1883).

© 2017 Pearson Education, Inc.

- The action spectrum for photosynthesis is **broad**er than the absorption spectrum of chlorophyll
- *Accessory pigments*, such as chlorophyll b, broaden the spectrum used for photosynthesis
- Other accessory pigments (**carotenoids**) may also broaden the spectrum of colors that drive photosynthesis
- Some carotenoids function in *photoprotection*; they absorb excessive light that would damage chlorophyll or react with oxygen


© 2017 Pearson Education, Inc.

Excitation of Chlorophyll by Light


- When a pigment absorbs light, it goes from a ground state to an excited state, which is unstable
- When excited electrons fall back to the ground state, excess energy is released as **heat**
- In isolation, some pigments also emit light, an afterglow called **fluorescence**

© 2017 Pearson Education, Inc.

Figure 10.12



(a) Excitation of isolated chlorophyll molecule



(b) Fluorescence

© 2017 Pearson Education, Inc.

A Photosystem: A Reaction-Center Complex Associated with Light-Harvesting Complexes

- A **photosystem** consists of a **reaction-center** complex surrounded by **light-harvesting complexes**
- The **reaction-center complex** is an association of proteins holding a special pair of chlorophyll *a* molecules and a primary electron acceptor
- The **light-harvesting complex** consists of pigment molecules bound to proteins

© 2017 Pearson Education, Inc.


- Light-harvesting complexes transfer the energy of photons to the chlorophyll *a* molecules in the reaction-center complex
- *These chlorophyll *a* molecules are special because they can transfer an excited electron to a different molecule*

© 2017 Pearson Education, Inc.

- A **primary electron acceptor** in the reaction center accepts excited electrons and is reduced as a result
- Solar-powered transfer of an electron from a chlorophyll *a* molecule to the primary electron acceptor is the first step of the light reactions

© 2017 Pearson Education, Inc.

Figure 10.13



(a) How a photosystem harvests light

© 2017 Pearson Education, Inc.

- There are two types of photosystems in the thylakoid membrane
- **Photosystem II (PS II) functions first** (the numbers reflect order of discovery)
- The reaction-center chlorophyll *a* of PS II is called **P680** because it is best at absorbing a wavelength of 680 nm

© 2017 Pearson Education, Inc.

- **Photosystem I (PS I)** is best at absorbing a wavelength of 700 nm
- The reaction-center chlorophyll *a* of PS I is called **P700**

© 2017 Pearson Education, Inc.

Linear Electron Flow


- During the light reactions, there are two possible routes for electron flow: cyclic and linear
- **Linear electron flow**, the primary pathway, involves both photosystems and produces ATP and NADPH using light energy

© 2017 Pearson Education, Inc.

- There are eight steps in linear electron flow:
 1. A photon hits a pigment in a light-harvesting complex of PS II, and its energy is passed among pigment molecules until it excites P680
 2. An excited electron from P680 is transferred to the primary electron acceptor (we now call it P680⁺)

© 2017 Pearson Education, Inc.

Figure 10.14_1




© 2017 Pearson Education, Inc.

3. H_2O is split by enzymes, and the electrons are transferred from the hydrogen atoms to P680^+ , thus reducing it to P680
 - P680^+ is the strongest known biological oxidizing agent
 - The H^+ are released into the thylakoid space
 - O_2 is released as a by-product of this reaction

© 2017 Pearson Education, Inc.

Figure 10.14_2




© 2017 Pearson Education, Inc.

4. Each electron “falls” down an electron transport chain from the primary electron acceptor of PS II to PS I. Energy released by the fall drives the creation of a proton gradient across the thylakoid membrane
5. Potential energy stored in the proton gradient drives production of ATP by chemiosmosis

© 2017 Pearson Education, Inc.

Figure 10.14_3




© 2017 Pearson Education, Inc.

6. In PS I (like PS II), transferred light energy excites P700, which loses an electron to the primary electron acceptor
 - P700⁺ (P700 that is missing an electron) accepts an electron passed down from PS II via the electron transport chain

© 2017 Pearson Education, Inc.

Figure 10.14_4




© 2017 Pearson Education, Inc.

7. Each electron “falls” down an electron transport chain from the primary electron acceptor of PS I to the protein ferredoxin (Fd)
8. NADP^+ reductase catalyzes the transfer of electrons to NADP^+ , reducing it to NADPH
 - The electrons of NADPH are available for the reactions of the Calvin cycle
 - This process also removes an H^+ from the stroma


© 2017 Pearson Education, Inc.

Figure 10.14_5



© 2017 Pearson Education, Inc.

Figure 10.15




© 2017 Pearson Education, Inc.

Cyclic Electron Flow

- In **cyclic electron flow**, electrons cycle back from Fd to the PS I reaction center via a plastocyanin molecule (Pc)
- Cyclic electron flow uses only photosystem I and produces ATP, but not NADPH
- No oxygen is released

© 2017 Pearson Education, Inc.

Figure 10.16




© 2017 Pearson Education, Inc.

- ATP and NADPH are produced on the side facing the stroma, where the Calvin cycle takes place
- In summary, light reactions generate ATP and increase the potential energy of electrons by moving them from H_2O to NADPH


© 2017 Pearson Education, Inc.

Figure 10.UN03



© 2017 Pearson Education, Inc.

Figure 10.18



© 2017 Pearson Education, Inc.

Figure 10.22c

LIGHT REACTIONS	CALVIN CYCLE REACTIONS
• Are carried out by molecules in the thylakoid membranes • Convert light energy to the chemical energy of ATP and NADPH • Split H_2O and release O_2 to the atmosphere	• Take place in the stroma • Use ATP and NADPH to convert CO_2 to the sugar G3P • Return ADP, inorganic phosphate, and $NADP^+$ to the light reactions

© 2017 Pearson Education, Inc.