

Chapter 4 Introduction to Probability

Learning Objectives

- 1. Obtain an appreciation of the role probability information plays in the decision making process.
- 2. Understand probability as a numerical measure of the likelihood of occurrence.
- 3. Know the three methods commonly used for assigning probabilities and understand when they should be used.
- 4. Know how to use the laws that are available for computing the probabilities of events.
- 5. Understand how new information can be used to revise initial (prior) probability estimates using Bayes' theorem.

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.



Uploaded By: 1210711@student.birzeit.edu



Chapter 4 **Solutions:**

1. Number of experimental Outcomes = (3)(2)(4) = 24

2.
$$\binom{6}{3} = \frac{6!}{3!3!} = \frac{6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{(3 \cdot 2 \cdot 1)(3 \cdot 2 \cdot 1)} = 20$$

$$ABC \quad ACE \quad BCD \quad BEF$$

$$ABD \quad ACF \quad BCE \quad CDE$$

$$ABE \quad ADE \quad BCF \quad CDF$$

$$ABF \quad ADF \quad BDE \quad CEF$$

$$ACD \quad AEF \quad BDF \quad DEF$$

3.
$$P_3^6 = \frac{6!}{(6-3)!} = (6)(5)(4) = 120$$

BDF BFD DBF DFB FBD FDB

4. a.



b. Let: H be head and T be tail

(H,H,H) (T,H,H) (H,H,T) (T,H,T) (H,T,H) (T,T,H) (H,T,T) (T,T,T)

c. The outcomes are equally likely, so the probability of each outcomes is 1/8.

5.
$$P(E_i) = 1/5$$
 for $i = 1, 2, 3, 4, 5$

 $P(E_i) \ge 0$ for i = 1, 2, 3, 4, 5

 $P(E_1) + P(E_2) + P(E_3) + P(E_4) + P(E_5) = 1/5 + 1/5 + 1/5 + 1/5 = 1$

The classical method was used.

4 - 2

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.

STUDENTS-HUB.com



6. $P(E_1) = .40, P(E_2) = .26, P(E_3) = .34$

The relative frequency method was used.

- 7. No. Requirement (4.4) is not satisfied; the probabilities do not sum to 1. $P(E_1) + P(E_2) + P(E_3) + P(E_4) = .10 + .15 + .40 + .20 = .85$
- 8. a. There are four outcomes possible for this 2-step experiment; planning commission positive council approves; planning commission positive council disapproves; planning commission negative council disapproves.
 - b. Let p = positive, n = negative, a = approves, and d = disapproves



9.
$$\binom{50}{4} = \frac{50!}{4!46!} = \frac{50 \cdot 49 \cdot 48 \cdot 47}{4 \cdot 3 \cdot 2 \cdot 1} = 230,300$$

10. a. Use the relative frequency approach:

P(California) = 1,434/2,374 = .60

b. Number not from 4 states = 2,374 - 1,434 - 390 - 217 - 112 = 221

P(Not from 4 States) = 221/2,374 = .09

- c. P(Not in Early Stages) = 1 .22 = .78
- d. Estimate of number of Massachusetts companies in early stage of development = $(.22)390 \approx 86$

4 - 3

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.

STUDENTS-HUB.com



e. If we assume the size of the awards did not differ by states, we can multiply the probability an award went to Colorado by the total venture funds disbursed to get an estimate. Estimate of Colorado funds = (112/2374)(\$32.4) = \$1.53 billion

Authors' Note: The actual amount going to Colorado was \$1.74 billion.

11. a. Total drivers = 858 + 228 = 1086

$$P(\text{Seatbelt}) = \frac{858}{1086} = .79 \text{ or } 79\%$$

- b. Yes, the overall probability is up from .75 to .79, or 4%, in one year. Thus .79 does exceed his .78 expectation.
- c. Northeast $\frac{148}{200} = .74$

Midwest
$$\frac{162}{216} = .75$$

South
$$\frac{296}{370} = .80$$

West
$$\frac{252}{300} = .84$$

The West with .84 shows the highest probability of use.

d. Probability of selection by region:

Northeast
$$\frac{200}{1086} = .184$$

Midwest
$$\frac{216}{1086} = .200$$

South
$$\frac{370}{1086} = .340$$

West
$$\frac{300}{1086} = .286$$

South has the highest probability (.34) and West was second (.286).

e. Yes, .34 for South + .286 for West = .626 shows that 62.6% of the survey came from the two highest usage regions. The .79 probability may be high.

If equal numbers for each region, the overall probability would have been roughly

$$\frac{.74 + .75 + .80 + .84}{4} = .7825$$

4 - 4

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.

STUDENTS-HUB.com



Introduction to Probability

Although perhaps slightly lower, the .78 to .79 usage probability is a nice increase over the prior year.

12. a. Use the counting rule for combinations:

$$\binom{55}{5} = \frac{55!}{5!50!} = \frac{(55)(54)(53)(52)(51)}{(5)(4)(3)(2)(1)} = 3,478,761$$

One chance in 3,489,761

- b. Very small: 1/3,478,761 = .000000287
- c. Multiply the answer in part (a) by 42 to get the number of choices for the six numbers.

Number of Choices = (3,478,761)(42) = 146,107,962

Probability of Winning = 1/146,107,962 = .0000000684

- 13. Initially a probability of .20 would be assigned if selection is equally likely. Data does not appear to confirm the belief of equal consumer preference. For example using the relative frequency method we would assign a probability of 5/100 = .05 to the design 1 outcome, .15 to design 2, .30 to design 3, .40 to design 4, and .10 to design 5.
- 14. a. $P(E_2) = 1/4$
 - b. P(any 2 outcomes) = 1/4 + 1/4 = 1/2
 - c. P(any 3 outcomes) = 1/4 + 1/4 + 1/4 = 3/4
- 15. a. $S = \{ace of clubs, ace of diamonds, ace of hearts, ace of spades\}$
 - b. $S = \{2 \text{ of clubs}, 3 \text{ of clubs}, \dots, 10 \text{ of clubs}, J \text{ of clubs}, Q \text{ of clubs}, K \text{ of clubs}, A \text{ of clubs}\}$
 - c. There are 12; jack, queen, or king in each of the four suits.
 - d. For a: 4/52 = 1/13 = .08

For b: 13/52 = 1/4 = .25

For c: 12/52 = .23

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.



Uploaded By: 1210711@student.birzeit.edu



Chapter 4 16. a. (6)(6) = 36 sample points

b.

		Die 2						
		1	2	3	4	5	6	_
	1	2	3	4	5	6	7	
Die 1	2	3	4	5	6	7	8	Total for Both
	3	4	5	6	7	8	9	
	4	5	6	7	8	9	10	
	5	6	7	8	9	10	11	
	6	7	8	9	10	11	12	

- c. 6/36 = 1/6
- d. 10/36 = 5/18
- e. No. P(odd) = 18/36 = P(even) = 18/36 or 1/2 for both.
- f. Classical. A probability of 1/36 is assigned to each experimental outcome.
- 17. a. (4,6), (4,7), (4,8)
 - b. .05 + .10 + .15 = .30
 - c. (2,8), (3,8), (4,8)
 - d. .05 + .05 + .15 = .25
 - e. .15

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.



Uploaded By: 1210711@student.birzeit.edu



Introduction to Probability

18. a.
$$P(\text{no meals}) = \frac{11}{496} = .022$$

b. P(at least four meals) = P(4) + P(5) + P(6) + P(7 or more)

$$= \frac{36}{496} + \frac{119}{496} + \frac{114}{496} + \frac{139}{496}$$
$$= .073 + .240 + .230 + .280$$
$$= .823$$

c. *P*(two or fewer meals)

$$= P(2) + P(1) + P(0)$$

$$= \frac{30}{496} + \frac{11}{496} + \frac{11}{496}$$
$$= .060 + .022 + .022$$
$$= .104$$

19. a/b. Use the relative frequency approach to assign probabilities. For each sport activity, divide the number of male and female participants by the total number of males and females respectively.

Activity	Male	Female
Bicycle Riding	.18	.16
Camping	.21	.19
Exercise Walking	.24	.45
Exercising with Equipment	.17	.19
Swimming	.22	.27

c.
$$P(\text{Exercise Walking}) = \frac{(28.7+57.7)}{248.5} = .35$$

d.
$$P(Woman) = \frac{57.7}{(28.7+57.7)} = .67$$

$$P(\mathrm{Man}) = \frac{28.7}{(28.7 + 57.7)} = .33$$

- 20. a. P(N) = 54/500 = .108
 - b. P(T) = 48/500 = .096
 - c. Total in 5 states = 54 + 52 + 48 + 33 + 30 = 217

P(B) = 217/500 = .434

Almost half the Fortune 500 companies are headquartered in these five states.

4 - 7

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.

STUDENTS-HUB.com



21. a. Using the relative frequency method, divide each number by the total population of 281.4 million.

Age	Number	Probability
Under 19	80.5	.2859
20 to 24	19.0	.0674
25 to 34	39.9	.1417
35 to 44	45.2	.1604
45 to 54	37.7	.1339
55 to 64	24.3	.0863
65 and over	35.0	.1243
Total	281.4	1.0000

- b. P(20 to 24) = .0674
- c. P(20 to 34) = .0674 + .1417 = .2091
- d. P(45 or older) = .1339 + .0863 + .1243 = .3445
- 22. a. P(A) = .40, P(B) = .40, P(C) = .60
 - b. $P(A \cup B) = P(E_1, E_2, E_3, E_4) = .80$. Yes $P(A \cup B) = P(A) + P(B)$.
 - c. $A^{c} = \{E_{3}, E_{4}, E_{5}\}$ $C^{c} = \{E_{1}, E_{4}\}$ $P(A^{c}) = .60$ $P(C^{c}) = .40$
 - d. $A \cup B^c = \{E_1, E_2, E_5\}$ $P(A \cup B^c) = .60$
 - e. $P(B \cup C) = P(E_2, E_3, E_4, E_5) = .80$
- 23. a. $P(A) = P(E_1) + P(E_4) + P(E_6) = .05 + .25 + .10 = .40$

 $P(B) = P(E_2) + P(E_4) + P(E_7) = .20 + .25 + .05 = .50$

$$P(C) = P(E_2) + P(E_3) + P(E_5) + P(E_7) = .20 + .20 + .15 + .05 = .60$$

b. $A \cup B = \{E_1, E_2, E_4, E_6, E_7\}$

 $P(A \cup B) = P(E_1) + P(E_2) + P(E_4) + P(E_6) + P(E_7)$ = .05 + .20 + .25 + .10 + .05 = .65

- c. $A \cap B = \{E_4\}$ $P(A \cap B) = P(E_4) = .25$
- d. Yes, they are mutually exclusive.
- e. $B^{c} = \{E_{1}, E_{3}, E_{5}, E_{6}\}; P(B^{c}) = P(E_{1}) + P(E_{3}) + P(E_{5}) + P(E_{6})$ = .05 + .20 + .15 + .10 = .50
- 24. Let E = experience exceeded expectations M = experience met expectations
 - a. Percentage of respondents that said their experience exceeded expectations = 100 - (4 + 26 + 65) = 5%

P(E) = .05

b. $P(M \cup E) = P(M) + P(E) = .65 + .05 = .70$

4 - 8

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.

STUDENTS-HUB.com



- 25. Let M = male young adult living in his parents' home F = female young adult living in her parents' home
 - a. $P(M \cup F) = P(M) + P(F) P(M \cap F)$ = .56 + .42 - .24 = .74
 - b. $1 P(M \cup F) = 1 .74 = .26$
- 26. Let Y = high one-year returnM = high five-year return
 - a. P(Y) = 9/30 = .30

$$P(M) = 7/30 = .23$$

- b. $P(Y \cap M) = 5/30 = .17$
- c. $P(Y \cup M) = .30 + .23 .17 = .36$

$$P(\text{Neither}) = 1 - .36 = .64$$

27.

		Big		
		Yes	No	
D 10	Yes	849	3645	4494
Pac-10	No	2112	6823	8935
		2,961	10,468	13,429

a.
$$P(\text{Neither}) = \frac{6823}{13,429} = .51$$

b.
$$P(\text{Either}) = \frac{2961}{13,429} + \frac{4494}{13,429} - \frac{849}{13,429} = .49$$

c.
$$P(Both) = \frac{849}{13,429} = .06$$

- 28. Let: B = rented a car for business reasonsP = rented a car for personal reasons
 - a. $P(B \cup P) = P(B) + P(P) P(B \cap P)$ = .54 + .458 - .30 = .698

b.
$$P(\text{Neither}) = 1 - .698 = .302$$

29. a.
$$P(E) = \frac{1033}{2851} = .36$$

 $P(R) = \frac{854}{2851} = .30$
 $P(D) = \frac{964}{2851} = .34$

4 - 9

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.

STUDENTS-HUB.com

Uploaded By: 1210711@student.birzeit.edu

Introduction to Probability



b. Yes; $P(E \cap D) = 0$

c. Probability =
$$\frac{1033}{2375} = .43$$

d. 964(.18) = 173.52

Rounding up we get 174 of deferred students admitted from regular admission pool.

Total admitted = 1033 + 174 = 1207

P(Admitted) = 1207/2851 = .42

30. a.
$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{.40}{.60} = .6667$$

b.
$$P(B|A) = \frac{P(A \cap B)}{P(A)} = \frac{.40}{.50} = .80$$

c. No because $P(A | B) \neq P(A)$

31. a.
$$P(A \cap B) = 0$$

- b. $P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{0}{.4} = 0$
- c. No. $P(A | B) \neq P(A)$; \therefore the events, although mutually exclusive, are not independent.
- d. Mutually exclusive events are dependent.
- 32. a. Total sample size = 2000

Dividing each entry by 2000 provides the following joint probability table.

	Health Insurance				
Age	Yes	No	Total		
18 to 34	.375	.085	.46		
35 and over	.475	.065	.54		
	.850	.150	1.00		

- Let A = 18 to 34 age group
 - B = 35 and over age group
 - Y = Insurance coverage
 - N = No insurance coverage
- b. P(A) = .46P(B) = .54

Of population age 18 and over

46% are ages 18 to 34 54% are ages 35 and over

4 - 10

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.

STUDENTS-HUB.com



c. P(N) = .15

d.
$$P(N|A) = \frac{P(N \cap A)}{P(A)} = \frac{.085}{.46} = .1848$$

e.
$$P(N|B) = \frac{P(N \cap B)}{P(B)} = \frac{.065}{.54} = .1204$$

f.
$$P(A|N) = \frac{P(A \cap N)}{P(N)} = \frac{.085}{.150} = .5677$$

g. Probability of no health insurance coverage is .15. A higher probability exists for the younger population. Ages 18 to 34: .1848 or approximately 18.5% of the age group. Ages 35 and over: .1204 or approximately 12% of the age group. Of the no insurance group, more are in the 18 to 34 age group: .5677, or approximately 57% are ages 18 to 34.

33. a.

		Reason for Applying		
	Quality	Cost/Convenience	Other	Total
Full Time	.218	.204	.039	.461
Part Time	.208	.307	.024	.539
	.426	.511	.063	1.00

- b. It is most likely a student will cite cost or convenience as the first reason probability = .511. School quality is the first reason cited by the second largest number of students - probability = .426.
- c. P(Quality | full time) = .218/.461 = .473
- d. P(Quality | part time) = .208/.539 = .386
- e. For independence, we must have $P(A)P(B) = P(A \cap B)$.

From the table, $P(A \cap B) = .218$, P(A) = .461, P(B) = .426

P(A)P(B) = (.461)(.426) = .196

Since $P(A)P(B) \neq P(A \cap B)$, the events are not independent.

- 34. a. Let O =flight arrives on time
 - $O^{c} = flight arrives late$
 - S = Southwest flight
 - U = US Airways flight
 - J = JetBlue flight

Given: P(O | S) = .834 P(O | U) = .751 P(O | J) = .701

P(S) = .40 P(U) = .35 P(J) = .25

$$P(\mathbf{O} \mid \mathbf{S}) = \frac{P(\mathbf{O} \cap \mathbf{S})}{P(\mathbf{S})}$$

:. $P(O \cap S) = P(O \mid S)P(S) = (.834)(.4) = .3336$

4 - 11

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.

STUDENTS-HUB.com



Similarly

$$P(O \cap U) = P(O \mid U)P(U) = (.751)(.35) = .2629$$

$$P(O \cap J) = P(O \mid J)P(J) = (.701)(.25) = .1753$$

Joint probability table

	On time	Late	Total
Southwest	.3336	.0664	.40
US Airways	.2629	.0871	.35
JetBlue	.1753	.0747	.25
Total:	.7718	.2282	1.00

b. Southwest Airlines; P(S) = .40

c.
$$P(O) = P(S \cap O) + P(U \cap O) + P(J \cap O) = .3336 + .2629 + .1753 = .7718$$

d.
$$P(S|O^c) = \frac{P(S \cap O^c)}{P(O^c)} = \frac{.0664}{.2282} = .2910$$

Similarly, $P(U|O^{\circ}) = \frac{.0871}{.2282} = .3817$

$$P(\mathbf{J}|\mathbf{O}^{c}) = \frac{.0747}{.2282} = .3273$$

Most likely airline is US Airways; least likely is Southwest

35. a. The joint probability table is given.

Occupation	Male	Female	Total
Managerial/Professional	0.17	0.17	0.34
Tech./Sales/Admin.	0.10	0.17	0.27
Service	0.04	0.07	0.12
Precision Production	0.11	0.01	0.12
Oper./Fabricator/Labor	0.10	0.03	0.13
Farming/Forestry/Fishing	0.02	0.00	0.02
Total	0.54	0.46	1.00

b. Let MP = Managerial/Professional F = Female

$$P(MP|F) = \frac{P(MP \cap F)}{P(F)} = \frac{.17}{.46} = .37$$

c. Let PP = Precision Production M = Male

$$P(PP|M) = \frac{P(PP \cap M)}{P(M)} = \frac{.11}{.54} = .20$$

d. No. From part (c), $P(PP \mid M) = .20$. But, P(PP) = .12, so $P(PP \mid M) \neq P(PP)$

4 - 12

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.

STUDENTS-HUB.com



36. a. Let A = makes 1st free throw B = makes 2nd free throw Introduction to Probability

Assuming independence, $P(A \cap B) = P(A)P(B) = (.89)(.89) = .7921$

- b. $P(A \cup B) = P(A) + P(B) P(A \cap B) = (.89)(.89) .7921 = .9879$
- c. P(Miss Both) = 1 P(at least one) = 1 .9878 = .0121
- d. For this player use P(A) = .58

 $P(A \cap B) = (.58)(.58) = .3364$

 $P(A \cup B) = .58 + .58 - .3364 = .8236$

P(Miss Both) = 1 - .8236 = .1764

The probability Reggie Miller makes both free throws is .7921, while the center's probability is .3364. The probability Reggie Miller misses both free throws is only .0121, while the center's probability is a much higher, .1764. The opponent's strategy should be to foul the center and not Reggie Miller.

37. Let C = event consumer uses a plastic card
 B = event consumer is 18 to 24 years old
 B^c = event consumer is over 24 years old

Given information:

$$P(C) = .37$$

$$P(B|C) = .19$$

$$P(B^{c}|C) = .81$$

P(B) = .14

a.
$$P(C|B) = \frac{P(C \cap B)}{P(B)}$$

but $P(C \cap B)$ is unknown. So first compute

$$P(C \cap B) = P(C)P(B|C)$$

= .37(.19) = .0703

Then

$$P(C|B) = \frac{P(C \cap B)}{P(B)} = \frac{.0703}{.14} = .5021$$

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied or distributed without the prior consent of the publisher.

STUDENTS-HUB.com

Uploaded By: 1210711@student.birzeit.edu



b.
$$P(C|B^{\circ}) = \frac{P(C \cap B^{\circ})}{P(B^{\circ})}$$

but $P(C \cap B^c)$ and $P(B^c)$ are unknown. However, they can be computed as follows.

 $P(C \cap B^{\circ}) = P(C)P(B^{\circ}|C)$ = .37(.81) = .2997

 $P(B^{c}) = 1 - P(B) = 1 - .14 = .86$

Then

$$P(C|B^{c}) = \frac{P(C \cap B^{c})}{P(B^{c})} = \frac{.2997}{.86} = .3485$$

- c. There is a higher probability that the younger consumer, age 18 to 24, will use plastic when making a purchase. The probability that the 18 to 24 year old consumer uses plastic is .5021 and the probability that the older than 24 year old consumer uses plastic is .3485. Note that there is greater than .50 probability that the 18 to 24 years old consumer will use plastic.
- d. Companies such as Visa, Mastercard and Discovery want their cards in the hands of consumers who will have a high probability of using the card. So yes, these companies should get their cards in the hands of young consumers even before these consumers have established a credit history. The companies should place a low limit of the amount of credit charges until the young consumer has demonstrated the responsibility to handle higher credit limits.
- 38. Let M = event consumer is a man Let W = event consumer is a woman Let B = event preferred plain bottled water Let S = event preferred sports drink
 - a. P(B) = 280/400 = .70
 - b. Sports drink: 80 + 40 = 120

P(S) = 120/400 = .30

c. $P(\mathbf{M}|\mathbf{S}) = 80/120 = .67$

P(W|S) = 40/120 = .33

d. P(S)P(M|S) = .30(.67) = .20

P(S)P(W|S) = .30(.33) = .10

4 - 14

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.

STUDENTS-HUB.com



e. *P*(M) =.50

$$P(S|M) = \frac{P(M \cap S)}{P(M)} = \frac{.20}{.50} = .40$$

f. P(W) = .50

$$P(S|W) = \frac{P(W \cap S)}{P(W)} = \frac{.10}{.50} = .20$$

g. No;

$$P(S|M) \neq P(S)$$

$$P(S|W) \neq P(S)$$

- 39. a. Yes, since $P(A_1 \cap A_2) = 0$
 - b. $P(A_1 \cap B) = P(A_1)P(B \mid A_1) = .40(.20) = .08$ $P(A_2 \cap B) = P(A_2)P(B \mid A_2) = .60(.05) = .03$
 - c. $P(B) = P(A_1 \cap B) + P(A_2 \cap B) = .08 + .03 = .11$

d.
$$P(A_1|B) = \frac{.08}{.11} = .7273$$

$$P(A_2|B) = \frac{.03}{.11} = .2727$$

40. a.
$$P(B \cap A_1) = P(A_1)P(B \mid A_1) = (.20)(.50) = .10$$

 $P(B \cap A_2) = P(A_2)P(B \mid A_2) = (.50)(.40) = .20$
 $P(B \cap A_3) = P(A_3)P(B \mid A_3) = (.30)(.30) = .09$

b.
$$P(A_2|B) = \frac{.20}{.10 + .20 + .09} = .51$$

с.

	Events	$P(\mathbf{A}_i)$	$P(\mathbf{B} \mid \mathbf{A}_i)$	$P(\mathbf{A}_i \cap \mathbf{B})$	$P(\mathbf{A}_i \mid \mathbf{B})$
-	A_1	.20	.50	.10	.26
	A_2	.50	.40	.20	.51
	A_3	<u>.30</u>	.30	<u>.09</u>	.23
		1.00		.39	1.00

41. S_1 = successful, S_2 = not successful and B = request received for additional information.

a. $P(S_1) = .50$

b. $P(B | S_1) = .75$

4 - 15

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.

STUDENTS-HUB.com

Uploaded By: 1210711@student.birzeit.edu

Introduction to Probability

SOUTH-WESTERN CENGAGE Learning

Chapter 4

c.
$$P(S_1|B) = \frac{(.50)(.75)}{(.50)(.75) + (.50)(.40)} = \frac{.375}{.575} = .65$$

42. M = missed payment $D_1 = customer defaults$ $D_2 = customer does not default$

$$P(D_1) = .05$$
 $P(D_2) = .95$ $P(M \mid D_2) = .2$ $P(M \mid D_1) = 1$

.

a.
$$P(D_1|M) = \frac{P(D_1)P(M|D_1)}{P(D_1)P(M|D_1) + P(D_2)P(M|D_2)} = \frac{(.05)(1)}{(.05)(1) + (.95)(.2)} = \frac{.05}{.24} = .21$$

- b. Yes, the probability of default is greater than .20.
- 43. Let: S = small car

_

- S^{c} = other type of vehicle
- F = accident leads to fatality for vehicle occupant

We have P(S) = .18, so $P(S^c) = .82$. Also P(F | S) = .128 and $P(F | S^c) = .05$. Using the tabular form of Bayes Theorem provides:

	Prior	Conditional	Joint	Posterior
Events	Probabilities	Probabilities	Probabilities	Probabilities
S	.18	.128	.023	.36
S ^c	.82	.050	.041	.64
	1.00		.064	1.00

From the posterior probability column, we have P(S | F) = .36. So, if an accident leads to a fatality, the probability a small car was involved is .36.

44. a.
$$P(A_1) = .47$$
 $P(W | A_1) = .50$

 $P(A_2) = .53$ $P(W \mid A_2) = .45$

b. Using tabular approach

	Prior	Conditional	Joint	Posterior
	Probabilities	Probability	Probability	Probability
Events	$P(A_i)$	$P(W \mid A_i)$	$P(A_i \cap W)$	$P(\mathbf{A}_{i} \mid \mathbf{W})$
Graduate	.47	.50	.2350	.4963
Not a Graduate	.53	.45	.2385	.5037
		$P(\mathbf{W}) =$.4735	1.0000

$$P(A_i | W) = .4963$$

c.

	Prior Drobabilition	Conditional	Joint Drobability	Posterior
	Probabilities	Probability	Probability	Probability
Events	$P(A_i)$	$P(\mathbf{M} \mid \mathbf{A}_i)$	$P(A_i \cap W)$	$P(\mathbf{A}_i \mid \mathbf{M})$
Graduate	.47	.50	.2350	.4463
Not a Graduate	.53	.45	.2915	.5537
		$P(\mathbf{M}) =$.5265	1.0000

$P(A_i \mid M) = .4463$

4 - 16

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.

STUDENTS-HUB.com



Introduction to Probability

About a .05 higher probability a woman student will graduate compared to a man.

d. P(W) = .4735

P(M) = .5265

Approximately 47% women and 53% men.

45. a. Let S = person is age 65 or older

$$P(S) = \frac{34,991,753}{281,421,906} = .12$$

- b. Let D = takes prescription drugs regularly
 - $P(D) = P(D \cap S) + P(D \cap S^{c}) = P(D \mid S)P(S) + P(D \mid S^{c})P(S^{c})$ = .82(.12) + .49(.88) = .53
- c. Let $D_5 =$ takes 5 or more prescriptions

 $P(D_5 \cap S) = P(D_5 | S)P(S) = .40(.12) = .048$

d.
$$P(S \mid D_5) = \frac{P(S \cap D_5)}{P(D_5)}$$

 $P(D_5) = P(S \cap D_5) + P(S^c \cap D_5) = P(D_5 | S)P(S) + P(D_5 | S^c)P(S^c)$ = .40(.12) + (.28)(.88) = .048 + .246 = .294

$$P(S \mid D_5) = \frac{.048}{.294} = .16$$

46. a. P(24 to 26 | Yes) = .1482/.4005 = .3700

- b. P(Yes | 36 and over) = .0253/.1090 = .2321
- c. .1026 + .1482 + .1878 + .0917 + .0327 + .0253 = .5883
- d. P(31 or more | No) = (.0956 + .0837)/.5995 = .2991
- e. No, because the conditional probabilities do not all equal the marginal probabilities. For instance,

$$P(24 \text{ to } 26 \mid \text{Yes}) = .3700 \neq P(24 \text{ to } 26) = .3360$$

47. a. Let F = female. Using past history as a guide, P(F) = .40.

b. Let D = Dillard's

$$P(\mathbf{F}|\mathbf{D}) = \frac{.40\left(\frac{3}{4}\right)}{.40\left(\frac{3}{4}\right) + .60\left(\frac{1}{4}\right)} = \frac{.30}{.30 + .15} = .67$$

The revised (posterior) probability that the visitor is female is .67.

4 - 17

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.

STUDENTS-HUB.com



Chapter 4

We should display the offer that appeals to female visitors.

48. a.

	Yes	No	Total
23 and Under	.1026	.0996	.2022
24 - 26	.1482	.1878	.3360
27 - 30	.0917	.1328	.2245
31 - 35	.0327	.0956	.1283
36 and Over	.0253	.0837	.1090
Total	.4005	.5995	1.0000

.2022 b.

- .2245 + .1283 + .1090 = .4618с.
- .4005 d.
- P(Oil) = .50 + .20 = .7049. a.
 - b. Let S = Soil test results

Events	$P(A_i)$	$P(S \mid A_i)$	$P(A_i \cap S)$	$P(A_i S)$
High Quality (A ₁)	.50	.20	.10	.31
Medium Quality (A ₂)	.20	.80	.16	.50
No Oil (A ₃)	.30	.20	.06	<u>.19</u>
	1.00	P((S) = .32	1.00

P(Oil) = .81 which is good; however, probabilities now favor medium quality rather than high quality oil.

50. a. The first probability table is given.

	Household Income (\$1000s)					
Education Level	Under 24.9	25.0-49.9	50.0-74.9	75.0-99.9	100 or more	Total
Not H.S. Graduate	0.093	0.041	0.016	0.005	0.004	0.159
H.S. Graduate	0.101	0.098	0.060	0.027	0.020	0.308
Some College	0.060	0.082	0.058	0.032	0.031	0.264
Bachelor's Degree	0.021	0.040	0.040	0.027	0.047	0.175
Beyond Bachelor's Degree	0.008	0.015	0.018	0.016	0.038	0.095
Total	0.284	0.276	0.192	0.108	0.140	1.000

4 - 18

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.





b. This is a marginal probability.

P(Not H.S. graduate) = .159

c. This is the sum of 2 marginal probabilities.

 $P(Bachelor's Degree \cup Beyond Bachelor's Degree) = .175 + .095 = .27$

d. This is a conditional probability.

 $P(100 \text{ or more} | \text{Bachelor's Degree}) = \frac{.047}{.175} = .269$

e. This is a marginal probability.

P(Under 25) = P(Under 24.9) = .284

f. This is a conditional probability.

 $P(\text{Under } 25 \mid \text{Bachelor's Degree}) = \frac{.021}{.175} = .12$

- g. No. Note that the probabilities in parts (e) and (f) are not equal.
- 51. Let: B = blogger $B^c = non blogger$ Y = young adult (18-29) $Y^c = older adult$

Given: $P(B) = .08 P(Y | B) = .54 P(Y | B^{c}) = .24$

$$P(\mathbf{Y} \mid \mathbf{B}) = \frac{P(\mathbf{Y} \cap \mathbf{B})}{P(\mathbf{B})}$$

:. $P(Y \cap B) = P(Y | B)P(B) = (.54)(.08) = .0432$

$$P(\mathbf{Y} \mid \mathbf{B}^{c}) = \frac{P(\mathbf{Y} \cap \mathbf{B}^{c})}{P(\mathbf{B}^{c})}$$

:. $P(Y \cap B^{c}) = P(Y | B^{c})P(B^{c}) = (.24)(.92) = .2208$

	Young Adult	Older Adult	Total
Blogger	.0432	.0368	.08
Non Blogger	.2208	.6992	.92
Total:	.2640	.7360	1.00

b. $P(Y) = P(B \cap Y) + P(B^{c} \cap Y) = .0432 + .2208 = .2640$

c.
$$P(Y \cap C) = .0432$$

d.
$$P(B | Y) = \frac{P(B \cap Y)}{P(Y)} = \frac{.0432}{.2640} = .1636$$

4 - 19

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.

STUDENTS-HUB.com

Uploaded By: 1210711@student.birzeit.edu

Introduction to Probability



- 52. a. Probability of the event = P(average) + P(above average) + P(excellent)= $\frac{11}{50} + \frac{14}{50} + \frac{13}{50} = .22 + .28 + .26 = .76$
 - b. Probability of the event = P(poor) + P(below average)= $\frac{4}{50} + \frac{8}{50} = .24$
- 53. Let A = lost time accident in current yearB = lost time accident previous year

Given: P(B) = .06, P(A) = .05, P(A | B) = .15

- a. $P(A \cap B) = P(A \mid B)P(B) = .15(.06) = .009$
- b. $P(A \cup B) = P(A) + P(B) P(A \cap B) = .06 + .05 .009 = .101 \text{ or } 10.1\%$
- 54. Let I = treatment-caused injury
 - D = death from injury
 - N = injury caused by negligence
 - M = malpractice claim filed
 - \$ = payment made in claim

We are given P(I) = 0.04, P(N | I) = 0.25, P(D | I) = 1/7, P(M | N) = 1/7.5 = 0.1333, and P(\$ | M) = 0.50

- a. $P(N) = P(N | I) P(I) + P(N | I^c) P(I^c)$ = (0.25)(0.04) + (0)(0.96) = 0.01
- b. $P(D) = P(D | I) P(I) + P(D | I^c) P(I^c)$ = (1/7)(0.04) + (0)(0.96) = 0.006
- c. $P(M) = P(M | N) P(N) + P(M | N^c) P(N^c)$ = (0.1333)(0.01) + (0)(0.99) = 0.001333
 - P(\$) = P(\$ | M) P(M) + P(\$ | M^c) P(M^c)= (0.5)(0.001333) + (0)(0.9987) = 0.00067
- 55. a. P(A) = 200/800 = .25
 - b. P(B) = 100/800 = .125
 - c. $P(A \cap B) = 10/800 = .0125$
 - d. $P(A | B) = P(A \cap B)/P(B) = .0125/.125 = .10$
 - e. No, $P(A | B) \neq P(A) = .25$
- 56. a. Number favoring elimination = $.47(671) \approx 315$
 - b. Let F = in favor of proposal D = Democrat

P(F | D) = .29

4 - 20

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.

STUDENTS-HUB.com



c. P(F) = .47 and P(F | D) = .29

Introduction to Probability

Since $P(F) \neq P(F \mid D)$ they are not independent.

d. I would expect Republicans to benefit most because they are the ones who had the most people in favor of the proposal.

57. a.
$$P(\mathbf{B}|\mathbf{S}) = \frac{P(\mathbf{B} \cap S)}{P(\mathbf{S})} = \frac{.12}{.40} = .30$$

We have P(B | S) > P(B).

Yes, continue the ad since it increases the probability of a purchase.

b. Estimate the company's market share at 20%. Continuing the advertisement should increase the market share since P(B | S) = .30.

c.
$$P(\mathbf{B}|\mathbf{S}) = \frac{P(\mathbf{B} \cap S)}{P(\mathbf{S})} = \frac{.10}{.30} = .333$$

The second ad has a bigger effect.

58. a.
$$(2)(2) = 4$$

b. Let S = successful U = unsuccessful



This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.

STUDENTS-HUB.com



- d. $O \cup M = \{E_1, E_2, E_3\}$
- e. $O \cap M = \{E_1\}$
- f. No; since $O \cap M$ has a sample point.
- 59. Let I = important or very important M= male F = female
 - a. P(I) = .49 (a marginal probability)
 - b. $P(I \mid M) = .22/.50 = .44$ (a conditional probability)
 - c. P(I | F) = .27/.50 = .54 (a conditional probability)
 - d. It is not independent

 $P(I) = .49 \neq P(I \mid M) = .44$

$$P(I) = .49 \neq P(I \mid F) = .54$$

- e. Since level of importance is dependent on gender, we conclude that male and female respondents have different attitudes toward risk.
- 60. a. P(Excellent) = .18

and

P(Pretty Good) = .50

 $P(\text{Pretty Good} \cup \text{Excellent}) = .18 + .50 = .68$

Note: Events are mutually exclusive since a person may only choose one rating.

b. 1035(.05) = 51.75

We estimate 52 respondents rated US companies poor.

c. 1035(.01) = 10.35

We estimate 10 respondents did not know or did not answer.

This edition is intended for use outside of the U.S. only, with content that may be different from the U.S. Edition. This may not be resold, copied, or distributed without the prior consent of the publisher.

Uploaded By: 1210711@student.birzeit.edu