

Lehninger


Principles of Biochemistry

David L. Nelson | Michael M. Cox



Lipids: Structurally Diverse Class





Organic molecules that are characterized by low solubility in water, that is, are relatively hydrophobic.

STUDENTS-HUB.com Uploaded By: Rawan Rous

group

Lehninger Principles of Biochemistry, Sixth Edition © 2013 W. H. Freeman and Company

Biological Functions of Lipids

Storage of energy

- Reduced compounds: lots of available energy
- Hydrophobic nature: good packing

Insulation from environment


- Low thermal conductivity
- High heat capacity (can "absorb" heat)
- Mechanical protection (can absorb shocks)

Water repellant

- Hydrophobic nature: keeps surface of the organism dry
 - Prevents excessive wetting (birds)
 - Prevents loss of water via evaporation

Buoyancy control and acoustics in marine mammals

- Increased density while diving deep helps sinking (just a hypothesis)
- Spermaceti organ may focus sound energy: sound stun gun?



Box 10-1 *Lehninger Principles of Biochemistry, Fifth Edition*© 2008 W. H. Freeman and Company

- Feeds on squid (in very deep water >1,000m deep)
- Rests quietly waiting (without constant swimming effort)
- In order to do so, its density must be equal to the surrounding water → changes in buoyancy
- Deep sea water is colder than the surface, spermaceti oil freezes (becomes denser) → matching the density of water
- When the whale returns to the surface, the oil warms up and melts decreasing the density to match surface water density

STUDENTS-HUB.com Uploaded By: Rawan Rous

More Functions

Membrane structure

Main structure of cell membranes

Cofactors for enzymes

- Vitamin K: blood clot formation
- Coenzyme Q: ATP synthesis in mitochondria

Signaling molecules

- Paracrine hormones (act locally)
- Steroid hormones (act body-wide)
- Growth factors
- Vitamins A and D (hormone precursors)

Pigments

Color of tomatoes, carrots, pumpkins, some birds

Antioxidants

Vitamin E

Lipids can provide pigment





- ✓ Minor differences in the chemistry of these compounds produce pigments of strikingly different colors
- ✓ Birds obtain the pigments that color their feathers red or yellow by eating plant materials that contain carotenoid pigments (e.g., canthaxanthin and zeaxanthin)
- ✓ The differences in pigmentation between male and female birds are the result of differences in intestinal uptake and processing of carotenoids.

Classification of Lipids

- Based on the structure and function
 - Lipids that do not contain fatty acids: cholesterol, terpenes, ...
 - Lipids that contain fatty acids (complex lipids)
 - can be further separated into:
 - storage lipids and membrane lipids




Figure 10-7
Lehninger Principles of Biochemistry, Sixth Edition
© 2013 W. H. Freeman and Company

10.1 Storage Lipids

Derivatives of fatty acids

Fatty acids are hydrocarbon derivatives

Low oxidation state (highly reduced)

Cellular oxidation is highly exergonic

Fatty Acids

- Carboxylic acids with hydrocarbon chains containing between 4 to 36 carbons
- Almost all natural fatty acids have an even number of carbons
- Most natural fatty acids are unbranched
- Saturated: no double bonds between carbons in the chain
- Monounsaturated: one double bond between carbons in the alkyl chain
- Polyunsaturated: more than one double bond in the alkyl chain


Fatty Acid Nomenclature

Carbon skeleton	Structure*	Systematic name [†]	Common name (derivation)	Melting point (°C)	Solubility at 30 °C (mg/g solvent)	
					Water	Benzene
12:0	CH ₃ (CH ₂) ₁₀ COOH	<i>n</i> -Dodecanoic acid	Lauric acid (Latin <i>laurus,</i> "laurel plant")	44.2	0.063	2,600
14:0	CH ₃ (CH ₂) ₁₂ COOH	n-Tetradecanoic acid	Myristic acid (Latin <i>Myristica</i> , nutmeg genus)	53.9	0.024	874
16:0	CH ₃ (CH ₂) ₁₄ COOH	n-Hexadecanoic acid	Palmitic acid (Latin <i>palma,</i> "palm tree")	63.1	0.0083	348
18:0	CH ₃ (CH ₂) ₁₆ COOH	n-Octadecanoic acid	Stearic acid (Greek stear, "hard fat")	69.6	0.0034	124
20:0	CH ₃ (CH ₂) ₁₈ COOH	n-Eicosanoic acid	Arachidic acid (Latin <i>Arachis,</i> legume genus)	76.5		
24:0	CH ₃ (CH ₂) ₂₂ COOH	n-Tetracosanoic acid	Lignoceric acid (Latin <i>lignum,</i> "wood" + <i>cera,</i> "wax")	86.0		
16:1(Δ°)	$CH_3(CH_2)_5CH = $ $CH(CH_2)_7COOH$	cis-9-Hexadecenoic acid	Palmitoleic acid	1 to -0.5		
18:1(Δ°)	$CH_3(CH_2)_7CH = $ $CH(CH_2)_7COOH$	cis-9-Octadecenoic acid	Oleic acid (Latin oleum, "oil")	13.4		
18:2 (Δ ^{9,12})	CH ₃ (CH ₂) ₄ CH= CHCH ₂ CH= CH(CH ₂) ₇ COOH	cis-,cis-9,12- Octadecadienoic acid	Linoleic acid (Greek <i>linon,</i> "flax")	1–5		
18:3 (Δ ^{9,12,15})	$CH_3CH_2CH=$ $CHCH_3CH=$ $CHCH_2CH=$ $CH(CH_2)_7COOH$	cis-,cis-,cis-9,12,15- Octadecatrienoic acid	lpha-Linolenic acid	-11		
20:4 (Δ ^{5,8,11,14})	CH ₃ (CH ₂) ₄ CH= CHCH ₂ CH= CHCH ₂ CH= CHCH ₂ CH= CH(CH ₃) ₄ COOH	cis-,cis-,cis-, cis-5,8,11,14- Icosatetraenoic acid	Arachidonic acid	-49.5		


^{*}All acids are shown in their nonionized form. At pH 7, all free fatty acids have an ionized carboxylate. Note that numbering of carbon atoms begins at the carboxyl carbon.

[†]The prefix *n*- indicates the "normal" unbranched structure. For instance, "dodecanoic" simply indicates 12 carbon atoms, which could be arranged in a variety of branched forms; "*n*-dodecanoic" specifies the linear, unbranched form. For unsaturated fatty acids, the configuration of each double bond is indicated; in biological fatty acids the configuration is almost always cis.

Fatty Acid Nomenclature



(a) 18:1(Δ^9) cis-9-Octadecenoic acid



(b) 20:5($\Delta^{5,8,11,14,17}$) Eicosapentaenoic acid (EPA), an omega-3 fatty acid

- (a)Standard nomenclature # 1 to the carboxyl carbon (C-1), and α to the carbon next to it. The position of any double bond(s) is indicated by Δ followed by a superscript number indicating the lower-numbered carbon in the double bond.
- (b)Polyunsaturated fatty acids (PUFAs) nomenclature # 1 to the methyl carbon at the other end of the chain (also designated ω; the last letter in the Greek alphabet). The STEDENTICAL double bonds are indicated relative to the ω carbon Uploaded By: Rawan Rous

Omega-3 PUFA

- Omega-3 fatty acids are essential nutrients
 - Humans need them but cannot synthesize one
 - No synthetic pathway in humans to make the ω-3 PUFA α-linolenic acid; 18:3(Δ^{9,12,15}) ALA

- Must be obtained in the diet
 - Including ALA, DHA, and EPA
 - Although DHA and EPA can be synthesized from ALA

• Imbalance in ω -6 and ω -3 leads to an

Solubility and Melting Point of Fatty Acids

Solubility

decreases as the chain length increases

Melting Point

- decreases as the chain length decreases
- decreases as the number of double bonds increases At room temperature, saturated fatty acids (from 12:0 to 24:0) have a waxy consistency, whereas unsaturated fatty acids of the same lengths are oily liquids

Conformation of Fatty Acids

- The saturated chain tends to adopt extended conformations
- The double bonds in natural unsaturated fatty acids are commonly in cis configuration, which kinks the chain





Figure 10-2ab

Lehninger Principles of Biochemistry, Sixth Edition
© 2013 W. H. Freeman and Company

Melting Point and Double Bonds

- Saturated fatty acids pack in a fairly orderly way
 - extensive favorable interactions
- Unsaturated cis fatty acid pack less orderly due to the kink
 - less-extensive favorable interactions
- It takes less thermal energy to disrupt disordered packing of unsaturated fatty acids:
 - unsaturated cis fatty acids have a lower melting point



Trans Fatty Acids

- Trans fatty acids form by partial dehydrogenation of unsaturated fatty acids
 - Done to increase shelf life or stability at high temperature of oils used in cooking (especially deep frying)
- A trans double bond allows a given fatty acid to adopt an extended conformation
- Trans fatty acids can pack more regularly and show higher melting points than cis forms
- Consuming trans fats increases risk of cardiovascular disease
 - Avoid deep-frying partially hydrogenated vegetable oils
 - Current trend: reduce trans fats in foods (Wendy's, KFC)

TABLE 10-2

Trans Fatty Acids in Some Typical Fast Foods and Snacks

	Trans fatty acid content		
	In a typical serving (g)	As % of total fatty acids	
French fries	4.7–6.1	28-36	
Breaded fish burger	5.6	28	
Breaded chicken			
nuggets	5.0	25	
Pizza	1.1	9	
Corn tortilla chips	1.6	22	
Doughnut	2.7	25	
Muffin	0.7	14	
Chocolate bar	0.2	2	

Source: Adapted from Table 1 in Mozaffarian, D., Katan, M.B., Ascherio, P.H., Stampfer, M.J., & Willet, W.C. (2006). Trans fatty acids and cardiovascular disease. *N. Engl. J. Med.* **354**, 1604–1605.

Note: All data for foods prepared with partially hydrogenated vegetable oil in the United States in 2002.


Table 10-2

Lehninger Principles of Biochemistry, Sixth Edition


Triacylglycerols (Nonpolar)

- Majority of fatty acids in biological systems are found in the form of triacylglycerols
- Solid ones are called fats
- Liquid ones are called oils
- The primary storage form of lipids (body fat)
- Less soluble in water than fatty acids due to the lack of charged carboxylate group
- Less dense than water: fats and oils float

Triacylglycerols



Glycerol




1-Stearoyl, 2-linoleoyl, 3-palmitoyl glycerol, a mixed triacylglycerol

Fats Provide Efficient Fuel Storage

- The advantage of fats over polysaccharides:
 - Fatty acids carry more energy per carbon because they are more reduced
 - Fatty acids carry less water per gram because they are nonpolar

- Glucose and glycogen are for short-term energy needs, quick delivery
- Fats are for long-term (months) energy needs, good storage, slow delivery


Fats Provide Efficient Fuel Storage



 $125\,\mu\mathrm{m}$

Waxes

- Waxes are esters of long-chain saturated and unsaturated fatty acids with long-chain alcohols
- Insoluble and have high melting points ~ 60 − 100 °C
- Variety of functions:
 - Storage of metabolic fuel in plankton
 - Protection and pliability for hair and skin in vertebrates
 - Waterproofing of feathers in birds
 - Protection from evaporation in tropical plants and ivy
 - Used by people in lotions, ointments, and polishes



Wax: The Material of the Honeycomb

Beeswax is a mixture of a large number of lipids, including esters of triacontanol, and a long-chain alkane hentiacontane




Structural Lipids in Membranes (Polar)

- Contain polar head groups and nonpolar tails (usually attached fatty acids)
- Diversification can come from:
 - modifying a different backbone
 - changing the fatty acids
 - modifying the head groups
- The properties of head groups determine the surface properties of membranes
- Different organisms have different membrane lipid head group compositions
- Different tissues have different membrane lipid head group compositions

Glycerophospholipids

- Primary constituents of cell membranes
- Two fatty acids form ester linkages with the first and second hydroxyl groups of L-glycerol-3-phosphate
- Head group is charged at physiological pH



L-Glycerol 3-phosphate (sn-glycerol 3-phosphate)

Figure 10-8

Lehninger Principles of Biochemistry, Sixth Edition © 2013 W. H. Freeman and Company

Glycerol itself is not chiral.

However, glycerol is **prochiral** – it can be converted to a chiral compound by adding a substituent such as phosphate to either of the $-CH_2OH$ groups. Uploaded By: Rawan Rous

STUDENTS-HUB.com

General Structure of Glycerophospholipids

- Unsaturated fatty acids are commonly found connected to C2
- The highly polar phosphate group may be further esterified by an alcohol; such substituent groups are called the head groups




Figure 10-9
Lehninger Principles of Biochemistry, Sixth Edition
© 2013 W. H. Freeman and Company


Examples of Glycerophospholipids

Name of glycerophospholipid	Name of X — O	Formula of X	Net charge (at pH 7)
Phosphatidic acid	_	— H	-2
Phosphatidylethanolamine	Ethanolamine	NH ₃	0
Phosphatidylcholine	Choline	N -	0
Phosphatidylserine	Serine	H O-NH3	-1
Phosphatidylglycerol	Glycerol	но н	-1
Phosphatidylinositol 4,5-bisphosphate	<i>myo-</i> lnositol 4,5- bisphosphate	OH OPO3- OPO3- OH	-4 *
Cardiolipin	Phosphatidyl- glycerol	HO HO H	-2

Figure 10-9
Lehninger Principles of Biochemistry, Sixth Edition
© 2013 W. H. Freeman and Company

Phosphatidylcholine

- Phosphatidylcholine is the major component of most eukaryotic cell membranes
- Many prokaryotes, including E. coli, cannot synthesize this lipid; their membranes do not contain phosphatidylcholine



Ether Lipids: Plasmalogen

- Vinyl ether analog of phosphatidylethanolamine
- Common in vertebrate heart tissue
- Also found in some protozoa and anaerobic bacteria
- Function is not well understood
 - Resistant to cleavage by common lipases but cleaved by few specific lipases
 - Increase membrane rigidity?
 - Sources of signaling lipids?
 - May be antioxidants?

Ether Lipids: Plasmalogen




Figure 10-10

Lehninger Principles of Biochemistry, Sixth Edition
© 2013 W. H. Freeman and Company

Ether Lipids: Platelets-Activating Factor

- Aliphatic ether analog of phosphatidylcholine
- Acetic acid has esterified position C2
- First signaling lipid to be identified
- Stimulates aggregation of blood platelets
- Plays role in mediation of inflammation




Figure 10-10
Lehninger Principles of Biochemistry, Sixth Edition
© 2013 W. H. Freeman and Company

Sphingolipids

- The backbone of sphingolipids is NOT glycerol
- The backbone of sphingolipids is a long-chain amino alcohol sphingosine
- A fatty acid is joined to sphingosine via an amide linkage rather than an ester linkage as usually seen in lipids
- A polar head group is connected to sphingosine by a glycosidic or phosphodiester linkage
- The sugar-containing glycosphingolipids are found largely in the outer face of plasma membranes

Sphingolipids

Ceramide is the parent compound for this group (structurally similar to diacylglycerol).





Figure 10-13
Lehninger Principles of Biochemistry, Sixth Edition
© 2013 W. H. Freeman and Company



glycerol

sphingosine



Substituent (R)	Sphingolipid
H	Ceramide
Phosphocholine	Sphingomyelin
Sugar(s)	Glycosphingolipid

- 2. **J.** Stiban, L. Caputo and M. Colombini (2008) Ceramide synthesis in the endoplasmic reticulum can permeabilize mitochondria to proapoptotic proteins. J. Lipid Res. 49(3): 625-634.
- 3. J. Stiban, L.C. Silva and A.H. Futerman (2008) Ceramide-containing membranes: the interface between biophysics and biology. Trends Glycosci. Glycotech. 20(116): 297–313.

4. J. Stiban, R. Tidhar and A.H. Futerman (2009) Ceramide Synthases: Roles in Cell

10902-10910.

1. J. Stiban, D. Fistere and M. Colombini (2006) Dihydroceramide hinders ceramide

channel formation: Implications on apoptosis. Apoptosis. 11(5): 773-780.

- Physiology and Signaling, *In:* 'Sphingolipids as Signaling and Regulatory Molecules' Landes Bioscience

 5. Y. Pewzner-Jung Y, H. Park, E.L. Laviad, L.C. Silva, S. Lahiri, J. Stiban, et. al. A critical role for ceramide synthase 2 in liver homeostasis: I. alterations in lipid metabolic pathways.(2010) *J. Biol. Chem.* 285(14):
- 6. S. Samanta, J. Stiban, T.K. Maugel and M. Colombini (2011) Visualization of ceramide channels by transmission electron microscopy. Biochim. Biophys. Acta Biomembranes 1808(4):1196-1201.
- 7. *J. Stiban and M. Perera* (2015) Very long chain <u>ceramides</u> interfere with C16-ceramide channel formation: A plausible mechanism for regulating the initiation of intrinsic apoptosis. *Biochim. Biophys. Acta* 1848(2): 561-567.
- 8. M. Abou-Ghali and J. Stiban (2015) Regulation of ceramide channel formation and disassembly: Insights of the first of the properties. Saudi J. Biol. Sci. 22(6): 760-772


 Uploaded By: Rawan Rous

Sphingomyelin

- Ceramide (sphingosine + amide-linked fatty acid) + phosphocholine attached to the alcohol
- Sphingomyelin is abundant in myelin sheath that surrounds some nerve cells in animals

Figure 10-14
Lehninger Principles of Biochemistry, Sixth Edition
© 2013 W. H. Freeman and Company

Sphingomyelin is structurally similar to phosphatidylcholine






Figure 10-14


Lehninger Principles of Biochemistry, Sixth Edition
© 2013 W. H. Freeman and Company

STUDENTS-HUB.com

Glycosphingolipids and Blood Groups

- The blood groups are determined in part by the type of sugars located on the head groups in glycosphingolipids.
- The structure of sugar is determined by an expression of specific glycosyltransferases
 - Individuals with no active glycosyltransferase will have the O antigen
 - Individuals with a glycosyltransferase that transfers an N-acetylgalactosamine group have A blood group
 - Individuals with a glycosyltransferase that transfers a galactose group have B blood group

Glycosphingolipids determine blood groups






Glc

Gal

GlcNAc

GalNAc

Defects in the turnover of membrane lipids lead to a number of diseases




Box 10-1 figure 1

Lehninger Principles of Biochemistry, Sixth Edition
© 2013 W. H. Freeman and Company

Sterols and Cholesterol

- Sterol
 - Steroid nucleus: four fused rings
 - Hydroxyl group (polar head) in the A-ring
 - Various nonpolar side chains
- The steroid nucleus is almost planar



Physiological Role of Sterols

- Cholesterol and related sterols are present in the membranes of most eukaryotic cells
 - Modulate fluidity and permeability
 - Thicken the plasma membrane
 - Most bacteria lack sterols
- Mammals obtain cholesterol from food or synthesize it de novo in the liver
- Cholesterol, bound to proteins, is transported to tissues via blood vessels
 - Cholesterol in low-density lipoproteins tends to deposit and clog arteries
- Many hormones are derivatives of sterols

Membrane microdomains (rafts) Raft, enriched in sphingolipids, cholesterol **Cholesterol GPI-linked** protein **Outside Inside Doubly Acyl groups** acylated (palmitoyl, **Prenylated** protein Caveolin myristoyl) protein

Synthesis of cholesteryl esters

Esterification converts cholesterol to an even more hydrophobic form for storage and transport.




Figure 21-38
Lehninger Principles of Biochemistry, Fifth Edition
© 2008 W. H. Freeman and Company



Blood plasma Blood plasma after fast after meal


Figure 21-40b

Lehninger Principles of Biochemistry, Fifth Edition
© 2008 W. H. Freeman and Company


TABLE 21–1

Major Classes of Human Plasma Lipoproteins: Some Properties


		Composition (wt %)				
Lipoprotein	Density (g/mL)	Protein	Phospholipids	Free cholesterol	Cholesteryl esters	Triacylglycerols
Chylomicrons	<1.006	2	9	1	3	85
VLDL	0.95-1.006	10	18	7	12	50
LDL	1.006-1.063	23	20	8	37	10
HDL	1.063-1.210	55	24	2	15	4




- ☐ Structure of an LDL molecule
- \square Apolipoprotein B-100 (apoB-100) is one of the largest single polypeptide chains known (4,636 aa; M_r 513,000)
- ☐ One particle of LDL contains a core with about 1,500 molecules of cholesteryl esters, surrounded by a shell composed of about 500 more molecules of cholesterol, 800 molecules of phospholipids, and one molecule of apoB-100




Chylomicrons (×60,000)




VLDL (×180,000)



LDL (×180,000)



HDL (×180,000)



Mammary, muscle, or adipose tissue

- Dietary lipids are packaged into chylomicrons
- ➤ Much of their triacylglycerol content is released by lipoprotein lipase to adipose and muscle tissues during transport through capillaries
- ➤ Remaining chylomicrons (containing largely protein and cholesterol) are taken up by the liver
- ➤ Endogenous lipids and cholesterol from the liver are delivered to adipose and muscle tissue by **VLDL**
- ➤ Extraction of lipid from VLDL (along with loss of some apolipoproteins) gradually converts some of it to LDL, which delivers cholesterol to extrahepatic tissues or returns to the liver
- ➤ The liver takes up LDL, VLDL remnants, and chylomicron remnants by receptor mediated endocytosis
- Excess cholesterol in extrahepatic tissues is transported back to the liver as HDL
- In the liver, some cholesterol is converted to bile salts
 Uploaded By: Rawan Rous





Figure 21-42
Lehninger Principles of Biochemistry, Fifth Edition

Steroid Hormones

- Steroids are oxidized derivatives of sterols
- Steroids have the sterol nucleus, but lack the alkyl chain found in cholesterol
- More polar than cholesterol
- Steroid hormones are synthesized from cholesterol in gonads and adrenal glands
- They are carried through the body in the bloodstream, usually attached to carrier proteins
- Many of the steroid hormones are male and female sex hormones

Steroid Hormones



Biologically Active Lipids

- Are present in much smaller amounts than storage or structural lipids
- Play vital roles as signaling molecules between nearby cells
- Lipid soluble vitamins (A, D, E, and K)

Signaling Lipids

- Paracrine lipid hormones are present in small amounts but play vital roles as signaling molecules between nearby cells
- Enzymatic oxidation of arachidonic acid yields
 - prostaglandins,
 - thromboxanes,
 - leukotrienes
- Arachidonic acid: 20:4(Δ^{5,8,11,14})

Arachidonic Acid Derivatives as Signaling Lipids

- Variety of functions:
 - Inflammation and fever, increase in body temperature (prostaglandins)
 - Formation of blood clots (thromboxanes)
 - Smooth muscle contraction in lungs; overproduction causes asthmatic attacks (leukotrienes)
 - Smooth muscle contraction in uterus during labor and menstruation (prostaglandins)

Nonsteroidal antiinflammatory drugs (NSAIDs) such as **aspirin** and **ibuprofen** block the formation of prostaglandins and thromboxanes from arachidonate by inhibiting the enzyme cyclooxygenase (prostaglandin H2 synthase).




Figure 10-18
Lehninger Principles of Biochemistry, Fifth Edition
© 2008 W. H. Freeman and Company

Vitamin D regulates calcium uptake




Figure 10-20a
Lehninger Principles of Biochemistry, Sixth Edition
© 2013 W. H. Freeman and Company

Vitamin A (Retinol)

- Involved in visual pigment
- Precursor for other hormones involved in signaling




Figure 10-21
Lehninger Principles of Biochemistry, Sixth Edition
© 2013 W. H. Freeman and Company

Vitamin E, K, and other lipid quinones are antioxidants

(a)
Vitamin E: an antioxidant

$$\begin{array}{c} \text{CH}_{3} \\ \text{HO} \\ \text{CH}_{3} \\ \text{CH}_{2} \\ \text{CH}_{3} \\ \text{CH}_{3} \\ \text{CH}_{3} \\ \text{CH}_{3} \\ \text{CH}_{3} \\ \text{CH}_{2} \\ \text{CH}_{2} \\ \text{CH}_{2} \\ \text{CH}_{2} \\ \text{CH}_{2} \\ \text{CH}_{3} \\ \text{CH}_{4} \\ \text{CH}_{3} \\ \text{CH}_{4} \\ \text{CH}_{5} \\ \text{C$$

(b)
Vitamin K₁: a blood-clotting cofactor (phylloquinone)

(c)
Warfarin: a blood
anticoagulant

(d)

Ubiquinone: a mitochondrial electron carrier (coenzyme Q) (n = 4 to 8)

$$\begin{array}{c} \mathsf{CH_3O} \\ \mathsf{CH_3} \\ \mathsf{CH_3O} \\ \mathsf{CH_2-CH=C-CH_2} \\ \mathsf{CH_2-CH=C-CH_2} \\ \mathsf{CH_2-CH=C-CH_2} \\ \mathsf{CH_2-CH=C-CH_3} \\ \end{array}$$

(e)

Plastoquinone: a chloroplast electron carrier (n = 4 to 8)

$$\begin{array}{c} CH_{3} \\ CH_{3} \\ CH_{2}-CH= \\ C-CH_{2} + \\ \\ CH_{2}-CH= \\ \\ C-CH_{2} \\ \\ \end{array} \\ \begin{array}{c} CH_{3} \\ CH_{3} \\ CH_{2}-CH= \\ \\ C-CH= \\ \\ \end{array} \\ \begin{array}{c} CH_{3} \\ CH_{3} \\ CH_{2}-CH= \\ \\ C-CH= \\ \end{array}$$

(**f**)

Dolichol: a sugar carrier (n = 9 to 22)

$$\begin{array}{c} \mathsf{CH_3} & \mathsf{CH_3} & \mathsf{CH_3} \\ \mathsf{I} & \mathsf{I} & \mathsf{I} \\ \mathsf{HO} + \mathsf{CH_2} - \mathsf{CH_2} - \mathsf{CH_2} + (\mathsf{CH_2} - \mathsf{CH} = \mathsf{C} - \mathsf{CH_2})_n + \mathsf{CH_2} - \mathsf{CH} = \mathsf{C} - \mathsf{CH_3} \end{array}$$

Figure 10-22

Lehninger Principles of Biochemistry, Sixth Edition

Polyketides are biologically active lipids with medicinal uses

Erythromycin (antibiotic)

Amphotericin B (antifungal)

Lovastatin (statin)