## **Gastrointestinal Tract**

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Organ Mouth and pharynx Salivary glands Esophagus Stomach Pancreas Liver Gallbladder Small intestine Large intestine Uploaded By: anonymous

STUDENTS-HUB.com

# **Organs of the Digestive system**

 Alimentary canal: performs the digestive functions (ingests, digests, absorbs, and defecates)
 -mouth, pharynx, esophagus, stomach, small intestine, And large intestine.

 Accessory digestive organs: assist the process of digestive breakdown in various ways
 -teeth, tongue, and several large digestive glands

# Mouth

- Oral or buccal cavity:
  - Is bounded by lips, cheeks, palate, and tongue
    - Has the oral orifice as its anterior opening
    - Is continuous with the oropharynx posteriorly
- To withstand abrasions:

 The mouth is lined with stratified squamous epithelium
 The gums, hard palate, and dorsum of the tongue are slightly keratinized

#### **Oral Cavity and Pharynx: Anterior View**



STUDENTS-HUB.com

#### Uploaded By: anonymous

#### Anatomy of the Oral Cavity: Mouth





STUDENTS-HUB.com

 $\label{eq:copyright} @ 2006 \ Pearson \ Education, \ Inc., \ publishing \ as \ Benjamin \ Cummings.$ 

Uploaded By: anonymous

# Tongue

- Intrinsic muscles change the shape of the tongue
- Extrinsic muscles alter the tongue's position
- Lingual frenulum secures the tongue to the floor of the mouth

#### Tongue



STUDEN SOBUERCO Folucation, Inc., publishing as Benjamin Cummings

Uploaded By: arguny 123081s

# **Salivary Glands**

- Produce and secrete saliva that:
  - Cleanses the mouth
  - Moistens and dissolves food chemicals
  - Aids in bolus formation
  - Contains enzymes that break down starch
- Three pairs of extrinsic glands parotid, submandibular, and sublingual
- Intrinsic salivary glands (buccal glands) scattered throughout the oral mucosa

STUDENTS-HUB.com

# **Salivary Glands**

- Parotid lies anterior to the ear between the masseter muscle and skin
- Parotid duct opens into the vestibule next to second upper molar
- Submandibular lies along the medial aspect of the mandibular body

Its ducts open at the base of the lingual frenulum

• Sublingual – lies anterior to the submandibular gland under the tongue

It opens via 10-12 ducts into the floor of the STUDENTS-HIMPOUTH Uploaded By: anonymous

#### **Salivary Glands**



# **Control of Salivation**

- Intrinsic glands keep the mouth moist
- Extrinsic salivary glands secrete serous, enzyme rich saliva in response to:
  - -Ingested food which stimulates chemoreceptors and pressoreceptors
  - -The thought of food
- Strong sympathetic stimulation inhibits salivation and results in dry mouth

# Pharynx

- From the mouth, the oro- and laryngopharynx allow passage of:
  - Food and fluids to the esophagus
  - Air to the trachea
- Lined with stratified squamous epithelium and mucus glands
- Has two skeletal muscle layers
  - Inner longitudinal
  - Outer pharyngeal constrictors

# **Esophagus**

- Muscular tube going from the laryngopharynx to the stomach
- Travels through the mediastinum and pierces the diaphragm
- Joins the stomach at the cardiac orifice

# **Esophagus**



STUDENTS-HUB.com

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings.

Uploaded By: anonymous

# **Esophageal Characteristics**

- Esophageal mucosa nonkeratinized stratified squamous epithelium
- The empty esophagus is folded longitudinally and flattens when food is present
- Glands secrete mucus as a bolus moves through the esophagus
- Muscularis changes from skeletal (superiorly) to smooth muscle (inferiorly)



Mosby items and derived items @ 2007, 2003 by Mosby, Inc.

#### Uploaded By: anonymous

STUDENTS-HUB.com

## Mucosa

- Moist epithelial layer that lines the lumen of the
- alimentary canal
- Three major functions:
- Secretion of mucus
- Absorption of end products of digestion
- Protection against infectious disease
- Consists of three layers: a lining epithelium, lamina propria, and muscularis mucosae

# **Mucosa: Epithelial Lining**

- Simple columnar epithelium and mucus-secreting
- goblet cells
- Mucus secretions:

-Protect digestive organs from digesting themselves

- Ease food along the tract
- Stomach and small intestine mucosa contain:
  - Enzyme-secreting cells
  - Hormone-secreting cells (making them endocrine

and digestive organs)

## Mucosa: Lamina Propria and Muscularis Mucosae

#### Lamina Propria

- Loose areolar and reticular connective tissue
- Nourishes the epithelium and absorbs nutrients
- Contains lymph nodes (part of MALT, mucosa associated lymphatic tissue) important in defense against bacteria
- Muscularis mucosae smooth muscle cells that produce local movements of mucosa

# **Mucosa: Other Sublayers**

- Submucosa dense connective tissue containing elastic fibers, blood and lymphatic vessels, lymph nodes, and nerves
- Muscularis externa responsible for segmentation and peristalsis
- Serosa the protective visceral peritoneum
  - Replaced by the fibrous adventitia in the esophagus
  - Retroperitoneal organs have both an adventitia and serosa

# **Enteric Nervous System**

- Composed of two major intrinsic nerve plexuses:
  - Submucosal nerve plexus regulates glands and smooth muscle in the mucosa
  - Myenteric nerve plexus Major nerve supply that controls GI tract mobility
- Segmentation and peristalsis are largely automatic involving local reflex arcs
- Linked to the CNS via long autonomic reflex arc

# **Digestive Processes in the Mouth**

- Food is ingested
- Mechanical digestion begins (chewing)
- Propulsion is initiated by swallowing
- Salivary amylase begins chemical breakdown of starch
- The pharynx and esophagus serve as conduits to pass food from the mouth to the stomach

# **Deglutition (Swallowing)**

- Coordinated activity of the tongue, soft palate, pharynx, esophagus, and 22 separate muscle groups
- Buccal phase bolus is forced into the oropharynx

# **Deglutition (Swallowing)**

Pharyngeal-esophageal phase – controlled by the medulla and lower pons

- All routes except into the digestive tract are sealed off
- Peristalsis moves food through the pharynx to the esophagus

Tongue pushes bolus against soft palate and back of mouth, triggering swallowing reflex. Upper esophageal sphincter relaxes while epiglottis closes to keep swallowed material out of the airways. Food moves downward into the esophagus, propelled by peristaltic waves and aided by gravity.







Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

# Stomach

- Chemical breakdown of proteins begins and food is converted to chyme
- Cardiac region surrounds the cardiac orifice
- Fundus dome-shaped region beneath the diaphragm
- Body midportion of the stomach
- Pyloric region made up of the antrum and canal which terminates at the pylorus
- The pylorus is continuous with the duodenum through the pyloric sphincter

# Stomach

- Greater curvature entire extent of the convex lateral surface
- Lesser curvature concave medial surface
- Lesser omentum runs from the liver to the lesser curvature
- Greater omentum drapes inferiorly from the greater curvature to the small intestine

# Stomach

- Nerve supply sympathetic and parasympathetic fibers of the autonomic nervous system
- Blood supply celiac trunk, and corresponding veins (part of the hepatic portal system)



STUDENTS-HUB.com

- Muscularis has an additional oblique layer that:
  - Allows the stomach to churn, mix, and pummel
  - food physically
- Breaks down food into smaller fragments

- Epithelial lining is composed of:
  Goblet cells that produce a coat of alkaline mucus
  - The mucous surface layer traps a bicarbonate-rich fluid beneath it
- Gastric pits contain gastric glands that secrete gastric juice, mucus, and gastrin



(a)

STUDENTS-HUB.com Copyright © 2

Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cuppingsled By: anonymous



STUDENTS-HUB.com

Uploaded B**y garren 23 m 5** bus



STUDENTS-HUB.com

Uploaded By igner 23mbc is

#### **Glands of the Stomach Fundus and Body**

- Gastric glands of the fundus and body have a variety of secretory cells
  - Mucous neck cells secrete acid mucus
- Parietal cells secrete HCI and intrinsic factor
- Chief cells produce pepsinogen
  -Pepsinogen is activated to pepsin by:
  - HCI in the stomach
  - Pepsin itself via a positive feedback mechanism
- Enteroendocrine cells secrete gastrin, histamine,endorphins, serotonin, cholecystokinin (CCK), and somatostatin into the lamina prop

STUDENTS-HUB.com



Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

STUDENTS-HUB.com
## Regulation and Mechanism of HCl Secretion

- HCl secretion is stimulated by ACh, histamine, and gastrin through second-messenger systems
- Release of hydrochloric acid:
  - Is low if only one ligand binds to parietal cells
  - Is high if all three ligands bind to parietal cells
- Antihistamines block H receptors and

2

decrease HCl release

### **Response of the Stomach to Filling**

- Stomach pressure remains constant until about 1L of food is ingested
- Relative unchanging pressure results from reflex-mediated relaxation and plasticity
- Reflex-mediated events include:
  - Receptive relaxation as food travels in the esophagus, stomach muscles relax
  - Adaptive relaxation the stomach dilates in response to gastric filling
- Plasticity intrinsic ability of smooth muscle to exhibit the stress-relaxation response

STUDENTS-HUB.com

Uploaded By: anonymous

## **Gastric Contractile Activity**

- Peristaltic waves move toward the pylorus at
- the rate of 3 per minute
- This basic electrical rhythm (BER) is initiated by pacemaker cells (cells of Cajal)
- Most vigorous peristalsis and mixing occurs near the pylorus
- Chyme is either:
  - delivered in small amounts to the duodenum or
- Forced backward into the stomach for further mixing STUDENTS-HUB.com



#### **Propulsion:**

peristaltic wave moves from the fundus to the pylorus: 30 ml of chyme is held in the stomach each contraction of the stomach muscle squirts

#### (b)

<u>Grinding:</u> most vigorous peristalsis and mixing occur close to the pylorus, small amount of chyme is delivered in to the duodenum

**Retropulsion:** peristaltic wave closes the pyloric valve forcing most of the contents of the pylorus back in to the stomach

#### Regulation of Gastric secretion

- Neural and hormonal mechanisms regulate the release of gastric juice
- Stimulatory and inhibitory events occur in three phases:

|                  | Excitatory  | Inhibitory  |
|------------------|---|---|
| Cephalic phase   | Sight, taste, smell thinking about food   | Loss of appetite or depression  |
| Gastric phase    | Release of gastrin to the<br>blood, activation of<br>stretch receptors,<br>chemoreceptors by<br>peptides, caffeine &<br>rising PH | A PH lower than 2,<br>emotional upset that<br>override the<br>Parasympathetic<br>activation |
| Intestinal phase | Low PH, partially<br>digested food enter the<br>duodenum & encourages<br>gastric gland activity                                   | Distention of the<br>duodenum, presence of<br>fatty, acidic or hypertonic<br>chyme          |
|                  |   |   |

### **Regulation of Gastric Emptying**

- Gastric emptying is regulated by:

   The neural enterogastric reflex
   Hormonal (enterogastrone) mechanisms
- These mechanisms inhibit gastric secretion and duodenal filling

### **Regulation of Gastric Emptying**

- Carbohydrate-rich chyme quickly moves through the duodenum
- Fat-laden chyme is digested more slowly causing food to remain in the stomach longer

### **Small Intestine: Gross Anatomy**

- Runs from pyloric sphincter to the ileocecal Valve, about 2 to 4 miters long
- Has three subdivisions: duodenum (10%), jejunum (40%), and ileum (55%)

### **Small Intestine: Gross Anatomy**

- The bile duct and main pancreatic duct:
  - Join the duodenum at the hepatopancreatic ampulla
  - Are controlled by the sphincter of Oddi
- The jejunum extends from the duodenum to the ileum
- The ileum joins the large intestine at the ileocecal valve

#### **Small Intestine: Microscopic Anatomy**

- Structural modifications of the small intestine wall increase surface area
  - Plicae circulares: deep circular folds of the mucosa and submucosa
    - Villi fingerlike extensions of the mucosa
  - Microvilli tiny projections of absorptive mucosal cells' plasma membranes

### **Duodenum and Related Organs**



#### STUDENTS-HUB.com

#### Uploaded By: anonymous

### **Small Intestine: Microscopic Anatomy**



Uploaded By Fanon 230 210



Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

STUDENTS-HUB.com

Uploaded By: anonymous

### **Small Intestine: Histology of the Wall**

- The epithelium of the mucosa is made up of:
  - Absorptive cells and goblet cells
  - Enteroendocrine cells
  - Interspersed T cells called intraepithelial lymphocytes (IELs)
  - -Cells of intestinal crypts secrete intestinal juice -Peyer 's patches are lymphatic tissues found in the submucosa, increase in numbers toward the end of the small intestine

-Brunner's glands in the duodenum secrete

alkaline mucus

## **Intestinal Juice**

- Secreted by intestinal glands in response to distension or irritation of the mucosa
- Slightly alkaline and isotonic with blood plasma
- Largely water, enzyme-poor, but contains mucus



© 2016 Columbia University. All Rights Reserved. Uploaded By: anonymous

STUDENTS-HUB.com

### Liver

- The largest gland in the body
- Superficially has four lobes right, left, caudate, and quadrate
- The falciform ligament:
  - Separates the right and left lobes anteriorly
  - Suspends the liver from the diaphragm and anterior abdominal wall

### **Liver: Associated Structures**

- Bile leaves the liver via:
- right and left hepatic ducts, which fuse into the common hepatic duct
- The common hepatic duct fuses with the cystic duct to form the common Bile duct

## Liver: Microscopic Anatomy

- Hexagonal-shaped liver lobules are the structural and functional units of the liver
- Composed of hepatocyte (liver cell) plates radiating outward from a central vein
- Portal triads are found at each of the six corners of each liver lobule



## Liver: Microscopic Anatomy

- Portal triads consist of a bile duct and
  - Hepatic artery supplies oxygen-rich blood to the liver
  - Hepatic portal vein carries venous blood with nutrients from digestive viscera
- Liver sinusoids enlarged, leaky capillaries located between hepatic plates
- Kupffer cells hepatic macrophages found in liver sinusoids

## Liver: Microscopic Anatomy

- Hepatocytes' functions include:
  - Production of bile
  - Processing bloodborne nutrients
  - Storage of fat-soluble vitamins
  - Detoxification
- Secreted bile flows between hepatocytes toward the bile ducts in the portal triads

### **Microscopic Anatomy of the Liver**



STUDENTS-HUB.com Copyright © 2006 Pearson Education, Inc., publishing as Benjamin Cummings ploaded By: anonymous

## **Composition of Bile**

- A yellow-green, alkaline solution containing bile salts, bile pigments, cholesterol, neutral fats, phospholipids, and electrolytes
  - Bile salts are cholesterol derivatives that:
  - Emulsify fat
  - Facilitate fat and cholesterol absorption
  - Help solubilize cholesterol
- Enterohepatic circulation recycles bile salts
- The chief bile pigment is bilirubin, a waste product of heme

### The Gallbladder

- Thin-walled, green muscular sac on the ventral surface of the liver
- Stores and concentrates bile by absorbing its water and ions
- Releases bile via the cystic duct, which flows into the bile duct

### **Regulation of Bile Release**



STUDENTS-HUB.com

#### Uploaded By Fanon 230 251s

### **Pancreas**

- Location
  - Lies deep to the greater curvature of the stomach
  - The head is encircled by the duodenum and the tail abuts the spleen

### **Pancreas**

- Exocrine function
  - Secretes pancreatic juice which breaks down all categories of foodstuff
  - Acini (clusters of secretory cells) contain zymogen granules with digestive enzymes
- The pancreas also has an endocrine function release of insulin and glucagon

### **Acinus of the Pancreas**



(a)

STUDENTS-HUB.com

### **Pancreatic Activation**



Uploaded By Fanon 230 2018

# Composition and Function of **Pancreatic Juice**

- Water solution of enzymes and electrolytes (primarily HCO<sub>3</sub><sup>-</sup>)
  - Neutralizes acid chyme
  - Provides optimal environment for pancreatic enzymes
- Enzymes are released in inactive form and activated in the duodenum

# Composition and Function of **Pancreatic Juice**

- Examples include
  - Trypsinogen is activated to trypsin
  - Procarboxypeptidase is activated to carboxypeptidase
- Active enzymes secreted
  - Amylase, lipases, and nucleases
  - These enzymes require ions or bile for optimal activity

### **Regulation of Pancreatic Secretion**

- Secretin and CCK are released when fatty or acidic chyme enters the duodenum
- CCK and secretin enter the bloodstream
- Upon reaching the pancreas:
  - CCK induces the secretion of enzyme-rich pancreatic juice
  - Secretin causes secretion of bicarbonate-rich pancreatic juice
- Vagal stimulation also causes release of pancreatic juice

#### **Regulation of Pancreatic Secretion**

During cephalic and gastric phases, stimulation by vagal nerve fibers causes release of pancreatic juice and weak contractions of the gallbladder.

(1) Acidic chyme entering duodenum causes the enteroendocrine cells of the duodenal wall to release secretin, whereas fatty, protein-rich chyme induces release of cholecystokinin.

> (2) Cholecystokinin and secretin enter bloodstream.

3 Upon reaching the pancreas, cholecystokinin induces the secretion of enzyme-rich pancreatic juice; secretin causes copious secretion of bicarbonate-rich pancreatic juice.

#### STUDENTS-HUB.com

#### Uploaded By Fanoe 2308 Is

0

0

### **Large Intestine**

- Has three unique features:
  - Teniae coli three bands of longitudinal smooth muscle in its muscularis
  - Haustra pocketlike sacs caused by the tone of the teniae coli
  - Epiploic appendages fat-filled pouches of visceral peritoneum

### Large Intestine

- Is subdivided into the cecum, appendix, colon, rectum, and anal canal
- The saclike cecum:
  - Lies below the ileocecal valve in the right iliac fossa
  - Contains a wormlike vermiform appendix

### **Large Intestine**


## Colon

- Has distinct regions: ascending colon, hepatic flexure, transverse colon, splenic flexure, descending colon, and sigmoid colon
- The transverse and sigmoid portions are anchored via mesenteries called mesocolons
- The sigmoid colon joins the rectum
- The anal canal, the last segment of the large intestine, opens to the exterior at the anus

### **Bacterial Flora**

- The bacterial flora of the large intestine consist of:
  - Bacteria surviving the small intestine that enter the cecum and
    - Those entering via the anus
- These bacteria:
  - Colonize the colon
  - Ferment indigestible carbohydrates
  - Release irritating acids and gases (flatus)
  - Synthesize B complex vitamins and vitamin K

## **Functions of the Large Intestine**

- Other than digestion of enteric bacteria, no further digestion takes place
- Vitamins, water, and electrolytes are reclaimed
- Its major function is propulsion of fecal material toward the anus

#### TABLE 21-1 The Digestive Hormones

|                                     | SECRETED BY   | TARGET(S)   | EFFECTS ON<br>ENDOCRINE<br>SECRETION                           | EFFECTS ON<br>EXOCRINE<br>SECRETION  |
|-------------------------------------|---|---|--|--|
| Gastrin                             | G cells in stomach<br>antrum  | ECL cells;<br>parietal cells                          | None   | Stimulates gastric<br>acid   |
| Cholecystokinin<br>(CCK)            | Endocrine<br>cells of small<br>intestine; neurons<br>of brain and gut | Gallbladder,<br>pancreas,<br>gastric smooth<br>muscle | None   | Stimulates pancreatic<br>enzyme secretion;<br>potentiates bicarbonate<br>secretion; inhibits<br>acid secretion |
| Secretin                            | Endocrine<br>cells in small<br>intestine                              | Pancreas,<br>stomach                                  | None   | Stimulates bicarbonate<br>secretion; inhibits gastric<br>acid and gastrin                                      |
| Gastric inhibitory<br>peptide (GIP) | Endocrine<br>cells in small<br>intestine                              | Beta cells of<br>endocrine<br>pancreas                | Stimulates<br>insulin release<br>(feedforward<br>mechanism)    | Inhibits acid<br>secretion   |
| Motilin                             | Endocrine cells in small intestine                                    | Smooth muscle<br>of antrum and<br>duodenum            | None   | None   |
| Glucagon-like<br>peptide 1          | Endocrine cells in small intestine                                    | Endocrine<br>pancreas                                 | Stimulates<br>insulin release;<br>inhibits glucagon<br>release | Possibly inhibits<br>acid secretion  |

Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

STUDENTS-HUB.com

### TABLE 21-1 The Digestive Hormones

|                                     | EFFECTS ON<br>MOTILITY   | OTHER EFFECTS                      | STIMULUS FOR<br>RELEASE   | RELEASE<br>INHIBITED BY | OTHER<br>INFORMATION   |
|-------------------------------------|--|------------------------------------|---|-------------------------|--|
| Gastrin                             | None   | Enhanced mucosal<br>cell growth    | Peptides and amino<br>acids in lumen; gastrin<br>releasing peptide and<br>ACh in nervous reflexes | pH < 1.5; somatostatin  | _  |
| Cholecystokinin<br>(CCK)            | Stimulates gallbladder<br>contraction for bile<br>release; inhibits gastric<br>emptying; promotes<br>intestinal motility | Stimulates satiety                 | Fatty acids and some amino acids  | Somatostatin            | Some effects may<br>be due to CCK as a<br>neuropeptide rather<br>than as a hormone           |
| Secretin                            | Inhibits gastric<br>emptying   | None                               | Acid in small intestine   | Somatostatin            | _  |
| Gastric inhibitory<br>peptide (GIP) | None   | Satiety and lipid<br>metabolism(?) | Glucose, fatty acids,<br>and amino acids in<br>small intestine                                    | NA                      | Acid inhibition<br>questionable<br>at physiological<br>concentrations                        |
| Motilin                             | Stimulates<br>migrating<br>motor complex   | Action in brain(?)                 | Fasting: periodic<br>release every<br>1.5–2 hours by<br>neural stimulus                           | NA                      | Changes associated<br>with both constipation<br>and diarrhea, but<br>relationship is unclear |
| Glucagon-like<br>peptide 1          | Slows gastric<br>emptying  | Satiety                            | Mixed meal that<br>includes carbohydrates<br>or fats in the lumen                                 | NA                      | Related to but not<br>identical to pancreatic<br>glucagon; acts in<br>concert with GIP       |

STUDENTS-HUB.com Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings. Uploaded By: anonymous



Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

STUDENTS-HUB.com



STUDENTS-HUB.com

Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.



Glucose enters the cell with Na<sup>+</sup> on the SGLT symporter and exits on GLUT2. Fructose enters on GLUT5 and exits on GLUT 2.

Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.



Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

#### STUDENTS-HUB.com





(b) Micelles are small disks with bile salts, phospholipids, fatty acids, cholesterol, and mono- and diglycerides.



Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

STUDENTS-HUB.com



Copyright © 2007 Pearson Education, Inc., publishing as Benjamin Cummings.

STUDENTS-HUB.com

# **Chemical Digestion: Nucleic Acids**

- Absorption: active transport via membrane carriers
- Absorbed in villi and transported to liver via hepatic portal vein
- Enzymes used: pancreatic ribonucleases and deoxyribonuclease in the small intestines

### **Electrolyte Absorption**

- Most ions are actively absorbed along the length of small intestine
  - Na is coupled with absorption of glucose and amino acids
  - Ionic iron is transported into mucosal cells where it binds to ferritin
- Anions passively follow the electrical potential established by Na<sup>+</sup>

### **Electrolyte Absorption**

- K +diffuses across the intestinal mucosa in response to osmotic gradients
- Ca<sup>2+</sup> absorption:
  - Is related to blood levels of ionic calcium
  - Is regulated by vitamin D and parathyroid hormone (PTH)

### Water Absorption

- 95% of water is absorbed in the small intestines by osmosis
- Water moves in both directions across intestinal mucosa
- Net osmosis occurs whenever a concentration gradient is established by active transport of solutes into the mucosal cells
- Water uptake is coupled with solute uptake, and as water moves into mucosal cells, substances follow along their concentration gradients

STUDENTS-HUB.com

### **Malabsorption of Nutrients**

- Results from anything that interferes with delivery of bile or pancreatic juice
- Factors that damage the intestinal mucosa (e.g., bacterial infection)
- Gluten enteropathy (adult celiac disease) gluten damages the intestinal villi and reduces the length of microvilli
  - Treated by eliminating gluten from the diet (all grains but rice and corn)