

56 Film Coated Tablets



SitaGlu-Met®

Sitagliptin / Metformin

50/850



Pharmacare PLC

Each film coated tablet contains 50 mg Sitagliptin (as phosphate monohydrate) and 850 mg Metformin HCl.
Store tablets at temperature not exceeding 25°C.
Keep out of reach of children.
Please read the enclosed leaflet.

PHARMACARE PLC
Betunia - Industrial Zone
PO Box 877 - Ramallah - Palestine
شركة دار الشفاء
الصيدلانية - المنطقة الصناعية
الرام الله - 877 - صندوق بريد

Patient Pack
Prescription Only
Carefully read the entire leaflet

SITAGLU-MET® 50/500, 50/850

COMPOSITION:

SITAGLU-MET 50/500: Each tablet contains 50 mg Sitagliptin (as phosphate monohydrate) and 500 mg Metformin HCl.

SITAGLU-MET 50/850: Each tablet contains 50 mg Sitagliptin (as phosphate monohydrate) and 850 mg Metformin HCl.

SITAGLU-MET 50/1000: Each tablet contains 50 mg Sitagliptin (as phosphate monohydrate) and 1000 mg Metformin HCl.

THERAPEUTIC ACTIVITY: Sitagliptin belongs to DPP-4 (Dipeptidyl peptidase 4) enzyme inhibitor and Metformin belongs to Biguanide group.

APPROVED INDICATIONS: Sitaglu-Met is a medicine contains 2 active ingredients: Sitagliptin and Metformin, which lower blood sugar. Together to control blood sugar levels in patients with type 2 diabetes in whom this combination is appropriate. Sitaglu-Met along with recommended diet and exercise program is intended to lower blood sugar levels in patients with type 2 diabetes.

WHEN SHOULD THE PREPARATION NOT BE USED?

- If you have type 1 diabetes mellitus.
- If you have kidneys which are not working properly.
- If you have diabetic ketoacidosis (increased ketone levels in the urine; diabetes complication which includes high blood sugar, rapid weight loss, nausea or vomiting).
- If you are going to get an injection of dye or contrast agents for a procedure, Sitaglu-Met will need to be stopped for a short time, talk to your doctor about when you should stop Sitaglu-Met and when you should start again.
- If you are allergic to any of the active ingredients. Symptoms of allergic reaction may include: rash, raised red patches on your skin, swelling of the face, lips, tongue, and throat that may cause difficulty breathing or swallowing.
- Sitaglu-met is not intended for administration to children under 18 years of age.