


6.4 Area of Surfaces of Revolution

Note Title

٢٢/٠١/٢٢



DEFINITION If the function $f(x) \geq 0$ is continuously differentiable on $[a, b]$, the **area of the surface** generated by revolving the graph of $y = f(x)$ about the x-axis is

$$S = \int_a^b 2\pi y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx = \int_a^b 2\pi f(x) \sqrt{1 + (f'(x))^2} dx. \quad (3)$$

Surface Area for Revolution About the y-Axis

If $x = g(y) \geq 0$ is continuously differentiable on $[c, d]$, the area of the surface generated by revolving the graph of $x = g(y)$ about the y-axis is


$$S = \int_c^d 2\pi x \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy = \int_c^d 2\pi g(y) \sqrt{1 + (g'(y))^2} dy. \quad (4)$$

EXAMPLE 1 Find the area of the surface generated by revolving the curve $y = 2\sqrt{x}$, $1 \leq x \leq 2$, about the x-axis

Sol: $f(x) = 2\sqrt{x} \Rightarrow f' = \frac{1}{\sqrt{x}}$ which is cont. on $[1, 2]$. Moreover $f(x) \geq 0 \forall x \in [1, 2]$. So,

$$\begin{aligned} S &= 2\pi \int_1^2 f(x) \sqrt{1 + (f')^2} dx = 2\pi \int_1^2 2\sqrt{x} \sqrt{1 + \frac{1}{x}} dx \\ &= 4\pi \int_1^2 \sqrt{x+1} dx = 4\pi \left[\frac{2}{3} (x+1)^{3/2} \right]_1^2 = \frac{8\pi}{3} (3\sqrt{3} - 2\sqrt{2}). \end{aligned}$$

الممكن $y = 2\sqrt{x}$ على الفترة $[1, 2]$ ، (الجسم الكروي) الذي أوجدنا مساحة سطحه موضح في الرسمة التالية .



EXAMPLE 2 The line segment $x = 1 - y$, $0 \leq y \leq 1$, is revolved about the y -axis to generate the cone in Figure 6.35. Find its lateral surface area (which excludes the base area).

in book

sol:

clearly $x = f(y) = 1 - y \geq 0$ on $[0, 1]$. Moreover

$\frac{dx}{dy} = -1$ is const. on $[0, 1]$. So the surface

area of the cone is

$$S = 2\pi \int_0^1 f(y) \sqrt{1 + f'(y)^2} dy = 2\pi \int_0^1 (1 - y) \sqrt{1 + (-1)^2} dy$$

$$= 2\sqrt{2}\pi \left(y - \frac{y^2}{2} \right) \Big|_0^1 = \boxed{\sqrt{2}\pi}$$

(الرسمة التالية توضح الجسم الكروي) الذي نتج عن دوران القطعة المستقيمة $x = 1 - y$ حول محور y .

