

# 1.1 Review of Calculus

1

## Limits and Continuity

Assume  $f(x)$  is defined on an open interval containing  $x_0$ .

Then [1]  $f$  has limit  $L$  at  $x_0$  if  $\lim_{x \rightarrow x_0} f(x) = L$

[2]  $f$  is continuous at  $x_0$  if  $\lim_{x \rightarrow x_0} f(x) = f(x_0)$

[3]  $f$  is continuous on a set  $S$  if  $f$  is continuous at each point  $x \in S$ .

•  $C^n(S)$ : is the set of all functions  $f$  s.t  $f$  and its first  $n$  derivatives are continuous on  $S$ .

Exp:  $f(x) = x^{4/3}$  is  $C^1[-1, 1]$

$f(x)$  and  $f'(x) = \frac{4}{3} x^{1/3}$  are continuous on  $[-1, 1]$

but  $f'' = \frac{4}{9} x^{-2/3}$  is not continuous at  $x=0$

## Convergent sequence

The sequence  $\{x_n\}_{n=1}^{\infty}$  converges to a limit  $L$  if

STUDENTS-HUB.com

$$\lim_{n \rightarrow \infty} x_n = L$$

(or we write  $x_n \rightarrow L$  as  $n \rightarrow \infty$ )

Uploaded By: anonymous

## Error Sequence

$\{E_n\}_{n=1}^{\infty} = \{x_n - L\}_{n=1}^{\infty}$  is called an error sequence

$\forall$ : for all  
 $\exists$ : there exists

s.t: such that  
 $f^{(n)}(x)$ :  $n^{\text{th}}$  derivative of  $f$

$f \in C[a, b]$ :  $f$  is continuous on  $[a, b]$   
 $f \in C^1[a, b]$ :  $f, f'$  are continuous on  $[a, b]$   
 $f \in C^n[a, b]$ :  $f, f', \dots, f^{(n)}$  are cont. on  $[a, b]$

Th • Assume  $f(x)$  is defined on the set  $S$ .

2

• Let  $x_0 \in S$ . Then the following are equivalent:

$f$  is cont. at  $x_0 \iff$  if  $\lim_{n \rightarrow \infty} x_n = x_0$  then  $\lim_{n \rightarrow \infty} f(x_n) = f(x_0)$

Th (Intermediate Value Theorem)

• Assume  $f \in C[a, b]$  and  $f(a) < L < f(b)$ .


• Then  $\exists c \in (a, b)$  s.t.  $f(c) = L$

Exp •  $f(x) = x^2$  is cont. on  $[0, 4]$

• Take  $L = 9 \in (f(0), f(4))$

• The solution of  $f(c) = 9$  is

$$c^2 = 9 \iff c = 3 \in (0, 4)$$



Th (Extreme Value Theorem)

• Assume that  $f \in C[a, b]$ , then  $f$  has abs. max

$f(b) = M = \max_{a \leq x \leq b} \{f(x)\}$  and abs. min  $m = \min_{a \leq x \leq b} \{f(x)\} = f(a)$

• Differentiation

•  $f$  is diff at  $x_0$  if  $f'(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0+h) - f(x_0)}{h}$  exists

•  $f$  is differentiable on set  $S$  if  $f$  has derivative at each point in  $S$ .

STUDENTS-HUB.COM

Uploaded By: anonymous

• If  $f(x)$  is diff at  $x_0$ , then  $f$  is cont. at  $x_0$ .


•  $m = f'(x_0)$  is the slope of the tangent line to the graph  $y = f(x)$  at the point  $(x_0, f(x_0))$ :

$$y - f(x_0) = m(x - x_0) \quad \text{tangent line}$$

### Th (Rolle's Theorem)

3


- Assume that  $f \in C[a, b]$  and  $f$  is diff on  $(a, b)$ .
- If  $f(a) = f(b)$ , then  $\exists$  a number  $c \in (a, b)$  s.t.  $f'(c) = 0$



Belzani here

### Th (Mean Value Theorem)

- Assume that  $f \in C[a, b]$  and  $f$  is Diff on  $(a, b)$ .
- Then,  $\exists$  a number  $c \in (a, b)$  s.t.  $f'(c) = \frac{f(b) - f(a)}{b - a}$



### Th (First Fundamental Theorem)

- Assume  $f \in C[a, b]$  and  $F$  is any antiderivative of  $f$  on  $[a, b]$ .
- Then,  $\int_a^b f(x) dx = F(b) - F(a)$  where  $F'(x) = f(x)$ .

### Th (Second Fundamental Theorem)

Assume  $f \in C[a, b]$ . Then  $\frac{d}{dx} \int_a^x f(t) dt = f(x) \quad \forall x \in (a, b)$

Exp  $\frac{d}{dx} \int_0^{x^2} \cos t dt = 2x \cos x^2$


STUDENTS-HUB.com

Exp  $f(x) = x^2$  on  $[0, 2]$

$av(f) = \frac{1}{2} \int_0^2 x^2 dx$

$= \frac{1}{2} \cdot \frac{8}{3}$

$= \frac{4}{3} = f(c) = c^2 \Leftrightarrow c = \frac{2}{\sqrt{3}}$



Uploaded By: anonymous

### Th (Mean Value Theorem for Integrals)

Assume  $f \in C[a, b]$ . Then  $\exists$  a number  $c \in (a, b)$  s.t.  $\frac{1}{b-a} \int_a^b f(x) dx = f(c)$

$f(c)$  is the average value of  $f$  over the interval  $[a, b]$

Area of rectangle is  $(b-a)f(c) = 2 \left( \frac{4}{3} \right) = \frac{8}{3}$  which is the red area

## Taylor Series Expansion

3.1

def Assume  $f(x)$  is infinitely many differentiable at  $x_0$ .

Then the Taylor series of  $f(x)$  at  $x_0$  is

$$f(x_0) + f'(x_0)(x-x_0) + \frac{f''(x_0)}{2!}(x-x_0)^2 + \frac{f'''(x_0)}{3!}(x-x_0)^3 + \dots$$

Exp Find the Maclaurin Series of  $e^x$ ,  $\cos x$ ,  $\sin x$   
 $x_0 = 0$

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots = \sum_{k=0}^{\infty} \frac{x^k}{k!}$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots = \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k}}{(2k)!}$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots = \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k+1}}{(2k+1)!}$$



## • Series

4

Given an infinite series  $a_1 + a_2 + \dots + a_n + \dots = \sum_{n=0}^{\infty} a_n$

- The  $n^{\text{th}}$  partial sum is  $S_n = a_1 + a_2 + \dots + a_n$
- The infinite series **converges** iff  $S_n$  converges ( $\lim_{n \rightarrow \infty} S_n = L$ ).  
otherwise, it diverges.

Exp  $\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = \sum_{n=1}^{\infty} \left( \frac{1}{n} - \frac{1}{n+1} \right)$ ,  $\frac{1}{n(n+1)} = \frac{A}{n} + \frac{B}{n+1}$  where  $A=1$   
 $B=-1$

$$S_n = \left(1 - \frac{1}{2}\right) + \left(\frac{1}{2} - \frac{1}{3}\right) + \left(\frac{1}{3} - \frac{1}{4}\right) + \dots + \left(\frac{1}{n} - \frac{1}{n+1}\right) = 1 - \frac{1}{n+1}$$

Hence,  $\lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \left(1 - \frac{1}{n+1}\right) = 1$  so the series converges to 1.

## Th (Taylor's Theorem)

Assume  $f \in C^{n+1}[a, b]$ . Let  $x_0 \in [a, b]$ . Then  $\exists$  a number  $c \in (x_0, x)$

s.t. the Taylor formula  $\overset{\text{True value}}{f(x)} = \overset{\text{approximated value about } x_0}{P_n(x)} + R_n(x)$  holds where

$P_n(x)$  is polynomial of degree  $n$  used to approximate  $f(x)$  with error (or remainder)  $R_n(x)$  given by:

$$P_n(x) = \sum_{k=0}^n \frac{f^{(k)}(x_0)}{k!} (x-x_0)^k \quad \text{and} \quad R_n(x) = \frac{f^{(n+1)}(c)}{(n+1)!} (x-x_0)^{n+1}$$


Exp  $f(x) = \sin x$  with  $x_0 = 0$ . Then

$f(x) = \sin x = P_9(x) + R_9(x)$  where

$$P_9(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \frac{x^9}{9!}$$

$$R_9(x) = \frac{f^{(10)}(c)}{(10)!} (x-0)^{10} \leq \frac{|x|^{10}}{(10)!}, \quad c \in (0, x)$$

$M=1$



Uploaded By: anonymous

$$\text{Error} = |R_n(x)| = \left| \frac{f^{(n+1)}(c)}{(n+1)!} \right| \frac{|x-x_0|^{n+1}}{(n+1)!} \leq M \frac{|x-x_0|^{n+1}}{(n+1)!}$$

## Linear Estimation:

5

$$f(x) = P_1(x) + R_1(x)$$

$$= f(x_0) + f'(x_0)(x-x_0) + \frac{f''(c)}{2!}(x-x_0)^2, \quad c \in (x_0, x)$$

Exp •  $f(x) = e^x$  with  $x_0 = 0$

$$e^x = P_1(x) + R_1(x)$$

$$= f(0) + f'(0)(x-0) + \frac{f''(c)}{2!}(x-0)^2$$

$$= 1 + x + \frac{e^c x^2}{2!}, \quad c \in (0, x)$$

Hence,  $e^x \approx 1+x$  with error  $= \left| \frac{e^c x^2}{2!} \right| = \frac{e^c x^2}{2!}$

$$e \approx 1+1 \text{ with error} = \frac{e^c}{2!} < \frac{3}{2}$$

$$c \in (0, 1)$$

$$0 < c < 1$$

$$e^0 < e^c < e$$

$$1 < e^c < 3 \Leftrightarrow 1 < e^c < e \Leftrightarrow$$

•  $e^x = P_2(x) + R_2(x)$

$$= f(0) + f'(0)(x-0) + \frac{f''(0)}{2!}(x-0)^2 + \frac{f'''(c)}{3!}(x-0)^3$$

$$= 1 + x + \frac{x^2}{2!} + \frac{e^c x^3}{3!}$$

$$0 < c < 0.1$$

$$1 < e^c < e^{0.1} < 2$$

Hence,  $e^x \approx 1+x+\frac{x^2}{2!}$  with error  $= \frac{e^c}{3!}|x^3|$

STUDENTS-HUB.com

$$e \approx 1 + 0.1 + \frac{(0.1)^2}{2!} \text{ with error} = \frac{e^c}{3!}(0.1)^3 < \frac{2}{3!}(0.001) < 10^{-3}$$


Uploaded By: anonymous

## Th (Bolzano)

Assume  $f(x) \in C[a, b]$  with  $f(a)f(b) < 0$

Then  $\exists$  a number  $c \in (a, b)$  s.t.  $f(c) = 0$

means:  $c$  is root for  $f(x)$   
 $c$  is zero for  $f(x)$   
 $c$  is solution for  $f(x) = 0$   
 $f$  crosses  $x$ -axis at  $x=c$   
 $x=c$  is the  $x$ -intercept



go to page 4



Def Assume  $\tilde{p}$  is approximation to  $p$ , where  $p \neq 0$ . Then,

- The (absolute) error is  $E_p = |p - \tilde{p}|$  and
- the relative error is  $R_p = \frac{|p - \tilde{p}|}{|p|}$  which expresses the error as percentage of the true value.

Exp Find the error and relative error for the following cases:

[1]  $x = 3.141592$  and  $\tilde{x} = 3.14$

$$\text{Error} = E_x = |x - \tilde{x}| = |3.141592 - 3.14| = 0.001592$$

$$\text{Relative Error} = R_x = \frac{|x - \tilde{x}|}{|x|} = \frac{0.001592}{3.141592} = 0.000507$$

$\tilde{x}$  is good approx. of  $x$

[2]  $y = 1000000$  and  $\tilde{y} = 999996$

$$E_y = |y - \tilde{y}| = |1000000 - 999996| = 4$$

$$R_y = \frac{|y - \tilde{y}|}{|y|} = \frac{4}{1000000} = 4 \times 10^{-6} = 0.000004$$

$\tilde{y}$  is good approx. of  $y$

[3]  $z = 0.000012$  and  $\tilde{z} = 0.000009$

$$E_z = |z - \tilde{z}| = |0.000012 - 0.000009| = 0.000003$$

$$R_z = \frac{|z - \tilde{z}|}{|z|} = \frac{0.000003}{0.000012} = 0.25$$

$\tilde{z}$  is bad approx. of  $z$

Remarks

[1]  $\tilde{x}$  is a good estimate for  $x$  since there is no much difference between  $E_x$  and  $R_x$  and so any of them could be used to determine the accuracy of  $\tilde{x}$ .

[2]  $\tilde{y}$  is good estimate for  $y$  since  $R_y$  is small (even if  $E_y$  is large since  $y$  is of magnitude  $10^6$ )

[3]  $\tilde{z}$  is bad approximation for  $z$  even that  $E_z$  is the smallest of the three cases. This because  $R_z$  is the largest.

Exp  $0.000004321$

$3.10045$

$2 \times 10^{-4} = 0.0002$

4 significant digits

6 significant digits

1 significant digit



Def The number  $\tilde{p}$  approximates  $p$  to  $d$  significant digits if  $d$  is the largest non-negative integer s.t. 7

$$R_p < 5 \times 10^{-d}$$

Exp [1]  $x = 3.141592$  and  $\tilde{x} = \underline{3.14}$

$$R_x = 0.000507 < 0.005 = 5 \times 10^{-3} \Leftrightarrow d=3$$

[2]  $y = 1000\ 000$  and  $\tilde{y} = \underline{999\ 996}$

$$R_y = 0.000\ 004 < 0.000\ 005 = 5 \times 10^{-6} \Leftrightarrow d=6$$

[3]  $z = 0.000\ 012$  and  $\tilde{z} = \underline{0.000\ 009}$

$$R_z = 0.25 < 0.5 = 5 \times 10^{-1} \Leftrightarrow d=1$$



- Exp II Describe the error in their sum

Hence, the error of the sum is the sum of the errors.

Assume  $\tilde{p}$  and  $\tilde{q}$  are good approximations for  $p$  and  $q$ . Show that the relative error in the product  $pq$  is approximately the sum of the relative errors in the approximation  $\tilde{p}$  and  $\tilde{q}$ .

$$p\tilde{q} - \tilde{p}\tilde{q} = \tilde{p} \in \mathfrak{g} + \tilde{q} \in \mathfrak{p} + \epsilon_p \in \mathfrak{g}$$

$$R_{pq} = \frac{p_q - \tilde{p}\tilde{q}}{p_q} = \frac{\tilde{p} \epsilon_q}{p_q} + \frac{\hat{q} \epsilon_p}{p_q} + \frac{\epsilon_p \epsilon_q}{p_q}$$

$$\approx \frac{G_f}{f} + \frac{G_p}{p} + 0$$

$$= R_f + R_p$$

- This is because  $\tilde{p}$  and  $\tilde{q}$  are good approximation for  $p$  and  $q$   
 $\Rightarrow \frac{\hat{p}}{p} \approx 1$  and  $\frac{\tilde{q}}{q} \approx 1$

## Normalized decimal form

7.2

Any real number  $p$  can be written in normalized decimal form:

$$p = \pm 0.d_1 d_2 d_3 \dots d_k d_{k+1} \dots \times 10^n$$

where  $d_1 \neq 0$  and  $d_j \in \{0, 1, 2, \dots, 9\}$  for  $j > 1$ .

Exp.  $0.01234 = 0.1234 \times 10^{-1}$

$12.034 = 0.12034 \times 10^2$

$0.000101 = 0.101 \times 10^{-3}$

## Source of Error

Truncation Error

↓  
Error results from  
estimating a formula  
by a formula

↓  
TE is the difference between  
a truncated value  $\tilde{p}$  and the  
actual value  $p$  arises from  
executing a finite number of  
steps to approximate an  
infinite process.

Exp.  $e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$   
 $\tilde{e} \approx 1 + x + \frac{x^2}{2!}$

TE = Error

$$= \left| e^x - \left( 1 + x + \frac{x^2}{2!} \right) \right|$$

Round-off Errors

↓  
Error results from  
estimating a number  
by a number

Round-off Errors  
Two Types

↓  
Rounding

↓  
 $fl(P)$   
round

↓  
rounded  
floating  
point  
representation

↓  
Chopping

↓  
 $fl(P)$   
chop

↓  
chopped  
floating  
point  
representation


Uploaded By: anonymous



Exp Assume the truncated Taylor series  $P_8(x)$  is used 7.3  
to approximate  $p = \int_0^{\frac{1}{2}} e^{x^2} dx = 0.544\ 987\ 104\ 184$ .  
Determine the accuracy and TE.

$$\tilde{p} = \int_0^{\frac{1}{2}} \left( 1 + x^2 + \frac{x^4}{2!} + \frac{x^6}{3!} + \frac{x^8}{4!} \right) dx = \left( x + \frac{x^3}{3} + \frac{x^5}{10} + \frac{x^7}{42} + \frac{x^9}{216} \right) \Big|_0^{\frac{1}{2}}$$

$$\begin{aligned} Z_p &= \frac{|p - \tilde{p}|}{|p|} = \frac{0.000\ 000\ 383\ 367}{0.544\ 987\ 104\ 184} \\ &= 7.03442 \times 10^{-7} \\ &= 0.000\ 000\ 703\ 442 < 0.000\ 005 = 5 \times 10^{-6} \end{aligned}$$



Exp TE  $p = \frac{22}{7} = 3.14\ 285\ 714\ 285\ 714\ 285\ 7 \dots$  computer works with finite digits  
Find the 6<sup>th</sup> digits representation of  $p$  in chopping and 16 rounding.  
 $f_l(p) = 3.14285 = \underline{0.314285 \times 10}$   
normalized  
chop

$$f_l(p) = 3.14286 = \underline{0.314286 \times 10}$$

normalized

round

Exp Find the 4<sup>th</sup> digits chopping and rounding of

[2]  $p = 0.123\ 444\ 445$

$$f_l(p) = 0.1234 = 0.1234 \times 10^0$$

chop

$$f_l(p) = 0.1235 = 0.1235 \times 10^0$$

round

STUDENTS HUB.com

[3]  $y = 2.00475$

$$f_l(y) = 2.004$$

chop

$$f_l(y) = 2.005$$

round

[4]  $x = 0.000\ 182\ 79$

$$f_l(x) = 0.000\ 1827$$

chop

$$f_l(x) = 0.000\ 1828$$

round

Uploaded By: anonymous

Q. How does computer approximate operations?

8

A. Apriority to 1) Brackets

2) Powers

3)  $\times, \div$  from left to right

4)  $+, -$  from left to right

Exp Use 4-digits rounding to find  $f(0.3456)$  if

$$f(x) = \frac{x - \sin \sqrt{x}}{2x^2 + x \cos x}$$

$$f(0.3456) = \frac{0.3456 - \sin(\sqrt{0.3456})}{2(0.3456)^2 + (0.3456) \cos(0.3456)}$$

$$= \frac{0.3456 - \sin(0.5879)}{2(0.1194) + (0.3456)(0.9409)}$$

$$= \frac{0.3456 - 0.5546}{0.2388 + 0.3252}$$

$$= \frac{-0.2090}{0.5640}$$

$$= -0.3706$$



Exp Determine the proper answer of  $\frac{3}{7} + \frac{5}{8} + \frac{11}{5}$  using four significant digits of accuracy. 9

$$\begin{aligned} \frac{3}{7} &= 0.428571... \approx 0.4286 \\ \frac{5}{8} &= 0.625 = 0.6250 \\ \frac{11}{5} &= 2.2 = 2.200 \end{aligned}$$

$$\left( \frac{3}{7} + \frac{5}{8} \right) + \frac{11}{5} = \frac{0.4286 + 0.6250 + 2.200}{21} = \frac{1.0536 + 2.200}{21} = \frac{1.054 + 2.200}{21} = \frac{3.254}{21} = 0.1550$$

$\swarrow$   
 $0.15495$

loss of Significance

- Let  $p = 3.1415926536$  and  $q = 3.1415957341$  with 11 decimal digits
- Note that  $p - q = -0.0000030805$  has 5 decimal digits
- We have loss of 6 digits (which are the first 6 digits in  $p$  and  $q$ )
- This is called loss of significance or subtractive cancellation.

$p - q$

Exp Let  $f(x) = x(\sqrt{x+1} - \sqrt{x})$  and  $g(x) = \frac{x}{\sqrt{x+1} + \sqrt{x}}$   
Use 6 digits and rounding to compare  $f(500)$  with  $g(500)$ .

$$\begin{aligned} f(500) &= 500(\sqrt{501} - \sqrt{500}) = 500(22.3830 - 22.3607) \\ &= 500(0.0223) \quad \text{loss of 3 digits} \\ &= 11.1500 \end{aligned}$$

$$g(500) = \frac{500}{\sqrt{501} + \sqrt{500}} = \frac{500}{22.3830 + 22.3607} = \frac{500}{44.7437} = 11.1748$$

- True value is  $11.1747553...$   $E_f = 0.0247$  and  $E_g = 0$
- Note that  $g(500)$  involves less error and becomes true value to the 6 digits. so  $g$  is a better approximation than  $f$

although  $f(x) = x(\sqrt{x+1} - \sqrt{x})$

$$\frac{\sqrt{x+1} + \sqrt{x}}{\sqrt{x+1} + \sqrt{x}} = \frac{x(x+1 - x)}{\sqrt{x+1} + \sqrt{x}} = \frac{x}{\sqrt{x+1} + \sqrt{x}} = g(x)$$

هذا بسبب عملية الطرح  
we solve this problem by finding  $g(x)$

How to solve this function to avoid loss of significant figures:

9.1

$$\textcircled{1} f(x) = \ln x - \ln(x+1)$$

$$P(x) = \ln \left( \frac{x}{x+1} \right)$$

•  $\textcircled{2} f(x) = \frac{x - \sin x}{\ln(x+2)}$ . Find  $f(\frac{7}{12})$  using 6 digits rounding.

$$\begin{aligned} \bullet f(\frac{7}{12}) &= f(0.583333) = \frac{0.583333 - \sin(0.583333)}{\ln(0.583333 + 2)} \\ &= \frac{0.583333 - 0.550809}{0.949080} \\ &= \frac{0.0325240}{0.949080} = 0.0342690 \end{aligned}$$

$$\bullet P(x) = \frac{x - \sin x}{\ln(x+2)} \cdot \frac{x + \sin x}{x + \sin x} = \frac{x^2 - \sin^2 x}{[\ln(x+2)][x + \sin x]}$$

$$P(0.583333) = \frac{(0.583333)^2 - \sin^2(0.583333)}{[\ln(0.583333+2)][0.583333 + \sin(0.583333)]}$$

STUDENTS-HUB.com

$$\begin{aligned} &= \frac{0.340277 - 0.303391}{(0.949080)(1.13414)} \\ &= \frac{0.0368860}{1.07639} \end{aligned}$$

$$= 0.0342682$$

Uploaded By: anonymous

we compare  
with the  
true value



Exp Compare the results of calculating  $f(0.01)$  and  $P(0.01)$  10 using 6 digits rounding arithmetic for

$$f(x) = \frac{e^x - 1 - x}{x^2} \quad \text{and} \quad P(x) = \frac{1}{2} + \frac{x}{6} + \frac{x^2}{24}$$

loss 1 digit      loss 2 digits

$$\bullet f(0.01) = \frac{e^{0.01} - 1 - 0.01}{(0.01)^2} = \frac{1.01005 - 1 - 0.01}{0.0001} = \frac{0.01005 - 0.01}{0.0001}$$

$$= \frac{0.00005}{0.0001} = 0.5 \Rightarrow E_f = 0.001671$$

$$\bullet P(0.01) = \frac{1}{2} + \frac{0.01}{6} + \frac{(0.01)^2}{24}$$

P solves the problem and it is easy to find it  
 $E_P = 0$

$$= 0.5 + 0.00166667 + 0.00000416670 = 0.501671$$

• Note that  $P(x)$  is Taylor polynomial of degree 2 for  $f(x)$  at  $x=0$ .

That is,  $f(x) = P_2(x) + R_2(x)$ .

• Now  $P(0.01)$  contains less error and becomes same as true answer 0.50167084168057542... when rounding

### Order of Approximation $O(h^n)$

Def • Assume  $f(h)$  is approximated by the function  $p(h)$ .

• Assume  $\exists$  a real constant  $M > 0$

and  $\exists$  a positive integer  $n$  so that

$$\frac{|f(h) - p(h)|}{|h^n|} \leq M \quad \text{for sufficiently small } h.$$

STUDENTS-HUB.COM

Uploaded By: anonymous

• In this case, we say  $p(h)$  approximates  $f(h)$  with order of approximation  $O(h^n)$  and we write this as

$$f(h) = p(h) + O(h^n)$$

Note: if we write \* as  $|f(h) - p(h)| \leq M |h^n|$ , then we see that  $O(h^n)$  stands in place of the error bound  $M |h^n|$ .

Th • Assume  $f(h) = p(h) + o(h^n)$  and  $g(h) = q(h) + o(h^m)$  where  $n, m$  are positive integers. 11

• Then  $f(h) + g(h) = p(h) + q(h) + o(h^r)$

and  $f(h)g(h) = p(h)q(h) + o(h^r)$

and  $\frac{f(h)}{g(h)} = \frac{p(h)}{q(h)} + o(h^r)$ ,  $g(h) \neq 0$  and  $q(h) \neq 0$

where  $r = \min\{n, m\}$

Exp If  $f(h) = p(h) + o(h^5)$  and  $g(h) = q(h) + o(h^3)$ , then  $f(h)g(h) = p(h)q(h) + o(h^3)$

Remark • If  $p(x)$  is the  $n^{\text{th}}$  Taylor polynomial approximation of  $f(x)$ , then by Taylor formula

$$f(x) = p(x) + R(x)$$

Truncation Error Term  $\rightarrow$  the remainder  $R(x)$  is simply  $o(h^{n+1})$ . That is

$$E = o(h^{n+1}) \approx M h^{n+1} \approx \frac{f^{(n+1)}(c)}{(n+1)!} h^{n+1}, \quad h \text{ small}, \quad c \in (x_0, x)$$

Th (Taylor's Th) Assume  $f \in C^{n+1}[a, b]$ . Then for  $x_0, x \in [a, b] \Rightarrow$

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x-x_0)^n + o(h^{n+1})$$

Uploaded By: anonymous


Remark ① If  $k \geq n$  then  $h^k + o(h^n) = o(h^n)$

Exp  $h^3 + o(h^3) = o(h^3)$  since  $h^3 + o(h^3) = h^3 + ch^3 = (1+c)h^3 = ch^3 = o(h^3)$

Exp  $h^4 + o(h^3) = o(h^3)$

② If  $f(h) = p(h) + o(h^n)$  with  $n > m$ , then  $p(h)$  is a better approximation for  $f(h)$ .



Exp 
$$e^h = 1 + h + \frac{h^2}{2!} + \frac{h^3}{3!} + \frac{h^4}{4!} + \dots$$
  12

•  $e^h \approx 1 + h + o(h^2)$  where  $E = o(h^2) \approx \frac{h^2}{2!} = \text{Error}$

$e^{0.1} = 1.105170918$  true value ↑  
order of approximation

$e^{0.1} \approx 1 + 0.1 = 1.1$  with error  $= M h^2 = M (0.1)^2 = M (0.01) < 10^{-2}$   
 where  $M = \frac{f^{(n+1)}(c)}{(n+1)!} = \frac{e^c}{2!}$ ,  $c \in (0, 0.1)$   
 $\Rightarrow 0 < c < 0.1 \Rightarrow 1 < e^c < e^{0.1} < 2 \Rightarrow M < 1$

•  $e^h = 1 + h + \frac{h^2}{2} + o(h^3) \Rightarrow \text{Error} = o(h^3) = M h^3$

$e^{0.1} = 1 + 0.1 + \frac{0.01}{2} = 1.105 \Rightarrow \text{Error} \approx M (0.1)^3 = M (0.001) < 10^{-3}$   
 since  $M < 1$

Exp  $\sinh h = h - \frac{h^3}{3!} + \frac{h^5}{5!} - \dots$

$\sinh h \approx h$  with error  $= o(h^3)$

$\Leftrightarrow \sin(0.1) \approx 0.1$

$\sinh h \approx h - \frac{h^3}{3!}$  with error  $= o(h^5)$

$\Leftrightarrow \sin(0.1) \approx 0.1 - \frac{(0.1)^3}{3!}$   
 $\approx 0.0998$

Exp Suppose  $e^h = 1 + h$  (Error  $= o(h^2)$ )

and  $\sinh h = h - \frac{h^3}{3!}$  (Error  $= o(h^5)$ )

Then 
$$e^h + \sinh h = 1 + 2h - \frac{h^3}{3!} + o(h^2) + o(h^5)$$
  

$$\approx 1 + 2h + o(h^2)$$

Exp  $\cosh h = 1 + \frac{h^2}{2!} + \frac{h^4}{4!} + \frac{h^6}{6!} + \frac{h^8}{8!} + \dots$

$\cosh h \approx 1 + \frac{h^2}{2!} + \frac{h^4}{4!}$  with  $E = o(h^6) = \text{constant} \cdot h^6$

Exp Consider the Taylor Polynomial expansions

13

$$e^h = 1 + h + \frac{h^2}{2!} + \frac{h^3}{3!} + o(h^4) \quad \text{and}$$

$$\cosh h = 1 - \frac{h^2}{2!} + \frac{h^4}{4!} + o(h^6).$$

Determine the order of approximation for their sum and product.

$$e^h + \cosh h = 1 + h + \cancel{\frac{h^2}{2!}} + \frac{h^3}{3!} + o(h^4) + 1 - \cancel{\frac{h^2}{2!}} + \frac{h^4}{4!} + o(h^6)$$

$$= 2 + h + \frac{h^3}{3!} + o(h^4) + \frac{h^4}{4!} + o(h^6)$$

But  $o(h^4) + \frac{h^4}{4!} = o(h^4)$  and

$o(h^4) + o(h^6) = o(h^4)$

$$= 2 + h + \frac{h^3}{3!} + o(h^4) \quad \text{with order of approximation } o(h^4).$$

$$e^h \cosh h = \left(1 + h + \frac{h^2}{2!} + \frac{h^3}{3!} + o(h^4)\right) \left(1 - \frac{h^2}{2!} + \frac{h^4}{4!} + o(h^6)\right)$$

or  $1 - \cancel{\frac{h^2}{2!}} + h - \frac{h^3}{2!} + \cancel{\frac{h^2}{2!}} + \frac{h^3}{3!} + o(h^4)$

$$= \left(1 + h + \frac{h^2}{2!} + \frac{h^3}{3!}\right) \left(1 - \frac{h^2}{2!} + \frac{h^4}{4!}\right) + \left(1 + h + \frac{h^2}{2!} + \frac{h^3}{3!}\right) o(h^6)$$

$$+ o(h^4) o(h^6) + \left(1 - \frac{h^2}{2!} + \frac{h^4}{4!}\right) o(h^4) \quad \text{error term}$$

$$= 1 + h - \frac{h^3}{3} - \frac{5h^4}{24} - \frac{h^5}{24} + \frac{h^6}{48} + \frac{h^7}{144} + o(h^6)$$

STUDENTS-HUB.com

Uploaded By: anonymous

$$+ o(h^4) o(h^6) + o(h^4)$$

But  $o(h^4) o(h^6) = o(h^{10})$  and so

$$-\frac{5}{24}h^4 - \frac{h^5}{24} + \frac{h^6}{48} + \frac{h^7}{144} + o(h^6) + o(h^{10}) + o(h^4) = o(h^4)$$

Hence,  $e^h \cosh h = 1 + h - \frac{h^3}{3} + o(h^4)$  and the order of approximation is  $o(h^4)$ .



## Def (order of Convergence of a sequence)

14

- Suppose that  $\lim_{n \rightarrow \infty} x_n = x$  and  $\lim_{n \rightarrow \infty} r_n = 0$
- We say  $x_n$  converges to  $x$  with order of convergence  $O(r_n)$  if  $\exists$  a constant  $K > 0$  s.t

$$\frac{|x_n - x|}{|r_n|} \leq K \quad \text{for } n \text{ sufficiently large}$$

and we write  $x_n = x + O(r_n)$

Exp show that [1]  $x_n = \frac{\cos n}{n^2}$  converges to 0 with rate of convergence  $O(\frac{1}{n^2})$ .

$$\frac{|x_n - x|}{|r_n|} = \frac{|\frac{\cos n}{n^2}|}{|\frac{1}{n^2}|} = |\cos n| \leq 1 \quad \text{for all } n$$

[2]  $p(h) = 1+h$  estimate  $f(h) = e^h$  with order  $O(h^2)$

$$\frac{|f(h) - p(h)|}{|r_h|} = \frac{|e^h - (1+h)|}{h^2} = \frac{\cancel{1} + \cancel{h} + \frac{h^2}{2!} + \frac{h^3}{3!} + \dots - (\cancel{1} + \cancel{h})}{h^2}$$

$$= \frac{1}{2!} + \frac{h}{3!} + \frac{h^2}{4!} + \frac{h^3}{5!} + \dots = \sum_{n=2}^{\infty} \frac{h^{n-2}}{n!}$$

Apply Ratio Test to see  $\sum_{n=2}^{\infty} \frac{h^{n-2}}{n!}$  converges to some  $K$

STUDENTS-HUB.com

Uploaded By: anonymous

$$\lim_{n \rightarrow \infty} \frac{\frac{h^{n-1}}{(n+1)!}}{\frac{h^{n-2}}{n!}} = \lim_{n \rightarrow \infty} \frac{h}{n+1} = 0 < 1 \quad \text{for all } h$$

[3]  $\sinh = h - \frac{h^3}{3!} + O(h^5)$  Exercise