

5.2 Logarithmic Functions and Their Properties

94

The logarithmic function is defined by

$$y = \log_a x = \frac{\ln x}{\ln a}, \quad \underbrace{x > 0}_{\text{domain}}, \quad \underbrace{a > 0, a \neq 1}_{\text{base}}$$

$$\begin{array}{ccc} y = \log_a x & \Leftrightarrow & \frac{y}{a} = x \\ \text{logarithmic form} & & \text{exponential form} \end{array}$$

Exp Write the following logarithmic equations using exponential form:

$$\textcircled{1} \quad 4 = \log_2 16 \quad \Leftrightarrow \quad 2^4 = 16$$

$$\textcircled{2} \quad \frac{1}{2} = \log_9 3 \quad \Leftrightarrow \quad 9^{\frac{1}{2}} = 3$$

$$\textcircled{3} \quad -2 = \log_5 \left(\frac{1}{25}\right) \quad \Leftrightarrow \quad 5^{-2} = \frac{1}{25}$$

Natural logarithmic function is $y = \log_e x = \frac{\ln x}{\ln e} = \ln x$
when the base $a = e \Rightarrow \ln e = 1$

• Graph $y = \ln x$, domain = $(0, \infty)$

x	y = ln x
0.1	-2.3
0.5	-0.7
1	0
2	0.7
3	1.1
10	2.3
e	1


Using Calculator:

$\log x$ means $\log_{10} x$

$\ln x$ means $\log_e x$

Exp $\log 2 \approx 0.301$

$$\ln 2 \approx 0.693$$



Uploaded By: Jibreel Bornat

negative y-axis is the vertical asymptote

$$e \approx 2.718$$

Exp Solve for x

95

$$\textcircled{1} \log_3 x = 4 \Rightarrow x = 3^4 = 3 \times 3 \times 3 \times 3 = 81$$

$$\textcircled{2} \log_{16} x = -\frac{1}{2} \Rightarrow x = (16)^{-\frac{1}{2}} = \frac{1}{\sqrt{16}} = \frac{1}{4}$$

$$\begin{aligned} \textcircled{3} \log(4x+20) &= 2 \Rightarrow \log_{10}(4x+20) = 2 \Rightarrow 4x+20 = 10^2 \\ &\Rightarrow 4x+20 = 100 \Rightarrow 4x = 80 \Rightarrow \boxed{x=20} \end{aligned}$$

$$\begin{aligned} \textcircled{4} \ln(3x-17) &= 0 \Rightarrow \log_e(3x-17) = 0 \Rightarrow e^0 = 3x-17 \\ &\Rightarrow 3x-17 = 1 \Rightarrow 3x = 18 \Rightarrow \boxed{x=6} \end{aligned}$$

Exp write the following equation in logarithmic form:

$$\textcircled{1} 2^4 = 16 \Leftrightarrow \log_2 16 = 4$$

$$\textcircled{2} 3^{-1} = \frac{1}{3} \Leftrightarrow \log_3 \left(\frac{1}{3}\right) = -1$$

Exp Graph $y = \log_2 x$

$$y = \log_2 x \Leftrightarrow x = 2^y$$

$$\Rightarrow \begin{array}{c|c|c|c|c|c|c|c|c} y & -3 & -2 & -1 & 0 & 1 & 2 & 3 \\ \hline x & \frac{1}{8} & \frac{1}{4} & \frac{1}{2} & 1 & 2 & 4 & 8 \end{array}$$

Exp Graph $f(x) = 2^x$


STUDENTS-HUB.COM

x	f(x)
-3	$\frac{1}{8}$
-2	$\frac{1}{4}$
-1	$\frac{1}{2}$
0	1
1	2
2	4
3	8

2^x is the inverse function of $\log_2 x$ and

$\log_2 x$ is the inverse function of 2^x

since they are symmetric about $y=x$



Uploaded By: Jibriel Bornat

Exp Graph ① $y = \log_y^x$


② $y = 4^x$

96


③ $y = \log_q^x$

④ $y = 9^x$


⑤ $y = \log_q^{-x}$



y	-1	0	1	2	-2
x	$\frac{1}{4}$	1	4	16	$\frac{1}{16}$




$\log_4^x$ is the inverse function of $y = 4^x$



$y = q^x$


x	-1	0	1	2	-2
y	$\frac{1}{q}$	1	q	q^2	$\frac{1}{q^2}$



$y = 9^x$ is the inverse function of $y = \log_9^x$

⑤ $y = \log_q^{-x}$ compare with $y = \log_q^x$


$\frac{y}{q} = -x \Rightarrow x = -\frac{y}{q}$



STUDENTS-HUB.com

Uploaded By: Jibreel Bornat

x	$-\frac{1}{q}$	$-\frac{1}{q^2}$	-1	-q	-q^2
y	-2	-1	0	1	2



Logarithmic Properties

97

Assume the base $a > 0$ and $a \neq 1$. Then

[1] $\log_a a^x = x$ for any real number x .

since the exponential form gives $a^x = a^x$ ✓

That is, if $a^y = a^x$ then $y = x$

Exp ① $\log_2 2^4 = 4$

② $\ln e^3 = \log_e e^3 = 3$

③ $\log_7 7 = \log_7 7^1 = 1$ ④ $\log_9 1 = \log_9 9^0 = 0$

[2] $\log_a a = 1$ and $\log_a 1 = 0$

Exp $\log_{12} 12 = 1$ and $\log_6 1 = 0$

[3] $a^{\log_a x} = x$ for any positive real number x

Exp ① $3^{\log_3 7} = 7$

② $e^{\ln x} = e^{\log_e x} = x$

check $\log_3 7 = \log_3 7$

check $\log_e x = \log_e x = \frac{\ln x}{\ln e} = \ln x$

[4] $\log_a xy = \log_a x + \log_a y$ for any positive real numbers x and y

STUDENTS HUB.com

Uploaded By: Jibreel Borhat

Exp Find ① $\log_3 27 = \log_3 3^3 = 3$

② $\log_4 \frac{1}{16} = \log_4 4^{-2} = -2$

Exp Assume $\log_a x = 2$ and $\log_a y = 3$. Find $\log_a x^2 y^3$

$$\log_a x^2 y^3 = \log_a x^2 + \log_a y^3 = 2 \log_a x + 3 \log_a y$$

$$= 2(2) + 3(3)$$

$$= 4 + 9 = 13$$

$$\boxed{5} \quad \log_a \frac{x}{y} = \log_a x - \log_a y \quad \text{for any } x > 0, y > 0$$

98

Exp Find ① $\log_3 \frac{9}{27} = \log_3 9 - \log_3 27$

$$= \log_3 3^2 - \log_3 3^3 = 2 - 3 = -1$$

② $\log_2 \frac{1}{\sqrt{2}} = \log_2 1 - \log_2 \sqrt{2} = 0 - \log_2 2^{\frac{1}{2}} = -\frac{1}{2}$

$$\boxed{6} \quad \log_a x^r = r \log_a x \quad \text{for any } x > 0 \text{ and } r \in \mathbb{R}$$

Exp Find ① $\log_2 32 = \log_2 2^5 = 5$

or $\ln x^3 e^2 = \log_e x^3 e^2 = \log_e x^3 + \log_e e^2 = 3 \log_e x + 2$
 $= 3 \frac{\ln x}{\ln e} + 2 = 3 \ln x + 2$

$$\boxed{7} \quad \log_b x = \frac{\log_a x}{\log_a b}$$

convert the base b to the base a

$$\log_b x = \frac{\ln x}{\ln b} = \frac{\log_e x}{\log_e b} \quad \text{convert the base b to the base e}$$

$$\log_b x = \frac{\log_{10} x}{\log_{10} b} = \frac{\log x}{\log b} \quad \text{convert the base b to the base 10}$$

STUDENTS-HUB.COM

Exp Evaluate ① $\log_7 15$ using the change of base e

$$\log_7 15 = \frac{\ln 15}{\ln 7} = \frac{2.71}{1.95} \approx 1.4$$

② $\log_7 15$ using the change of base 10

$$\log_7 15 = \frac{\log 15}{\log 7} = \frac{1.18}{0.845} \approx 1.4$$

Uploaded By: Jibreel Bornat