


6.3

Arc Length

114

Def: If f' is continuous on $[a, b]$, then the arc length of the curve $y=f(x)$ from $(a, f(a))$ to $(b, f(b))$ is

$$L = \int_a^b \sqrt{1 + [f'(x)]^2} dx = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$



• The Length of the curve is approximated by:

$$L = \sum_{k=1}^n L_k = \sum_{k=1}^n \sqrt{(\Delta x_k)^2 + (\Delta y_k)^2}$$

• By Mean Value Theorem, there is a point c_k with $x_{k-1} < c_k < x_k$ such that

$$f'(c_k) = \frac{\Delta y_k}{\Delta x_k} \quad \Leftrightarrow \quad \Delta y_k = f'(c_k) \Delta x_k$$

• Therefore, $L = \sum_{k=1}^n \boxed{\sqrt{1 + [f'(c_k)]^2}} \Delta x_k$ This is continuous on $[a, b]$ $\Rightarrow$

• The approximation L improves as the partition of $[a, b]$ becomes finer:

$$L = \lim_{n \rightarrow \infty} \sum_{k=1}^n \sqrt{1 + [f'(c_k)]^2} \Delta x_k = \int_a^b \sqrt{1 + [f'(x)]^2} dx$$

Example: Find the length of the curve $f(x) = \frac{x^3}{12} + \frac{1}{x}$, $1 \leq x \leq 4$.

$$f'(x) = \frac{x^2}{4} - \frac{1}{x^2} \Rightarrow 1 + [f'(x)]^2 = 1 + \left(\frac{x^2}{4} - \frac{1}{x^2}\right)^2 = 1 + \left(\frac{x^4}{16} - \frac{1}{2} + \frac{1}{x^4}\right)$$

$$= \frac{x^4}{16} + \frac{1}{2} + \frac{1}{x^4} = \left(\frac{x^2}{4} + \frac{1}{x^2}\right)^2$$

$$L = \int_1^4 \sqrt{\left(\frac{x^2}{4} + \frac{1}{x^2}\right)^2} dx = \int_1^4 \left(\frac{x^2}{4} + \frac{1}{x^2}\right) dx = \left[\frac{x^3}{12} - \frac{1}{x}\right]_1^4 = \left(\frac{64}{12} - \frac{1}{4}\right) - \left(\frac{1}{12} - 1\right) = \frac{72}{12} = 6.$$

Example: Find the length of the curve $y = \left(\frac{x}{2}\right)^{\frac{2}{3}}$ from $x=0$ to $x=2$. (115)

$$y' = \frac{2}{3} \left(\frac{x}{2}\right)^{-\frac{1}{3}} \cdot \frac{1}{2} = \frac{1}{3} \left(\frac{2}{x}\right)^{\frac{1}{3}}$$

" y' is not defined at $x=0$ "

$$\Rightarrow y^{\frac{3}{2}} = \frac{x}{2} \Rightarrow x = 2 y^{\frac{3}{2}}$$

$$\Rightarrow x' = 3 y^{\frac{1}{2}}$$

$$\Rightarrow (x')^2 = 9 y$$

$$\Rightarrow L = \int_0^1 \sqrt{1+9y} dy$$

$$= \int_1^{10} \frac{1}{9} u^{\frac{1}{2}} du$$

$$= \frac{1}{9} u^{\frac{3}{2}} \cdot \frac{2}{3} \Big|_1^{10} = \frac{2}{27} \left[(10)^{\frac{3}{2}} - (1)^{\frac{3}{2}} \right] = \frac{2}{27} [10\sqrt{10} - 1]$$

$$u = 1 + 9y$$

$$du = 9 dy$$

$$y=0 \Rightarrow u=1$$

$$y=1 \Rightarrow u=10$$

$$\approx 2.27$$

Example Find the length of the curve $y = \int_0^x \sqrt{\cos 2t} dt$ from $x=0$ to $x = \frac{\pi}{4}$

$$L = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

$$y' = \sqrt{\cos 2x} \Rightarrow (y')^2 = \cos 2x$$

$$= \int_0^{\frac{\pi}{4}} \sqrt{1 + \cos 2x} dx = \int_0^{\frac{\pi}{4}} \sqrt{1 + 2\cos^2 x - 1} dx$$

$$= \int_0^{\frac{\pi}{4}} \sqrt{2\cos^2 x} dx = \sqrt{2} \int_0^{\frac{\pi}{4}} \cos x dx$$

$$= \sqrt{2} \sin x \Big|_0^{\frac{\pi}{4}} = \sqrt{2} \left[\frac{1}{\sqrt{2}} - 0 \right] = 1$$