

GLUCOCARE®

5 mg Glibenclamide

100 Tablets

P S



LOT: EXP: MFG: PAK: 100 TABLETS

Patient Package Insert
Prescription Only Medicine
Carefully read the entire leaflet before starting use
GLUCOCARE® TABLET

COMPOSITION: Each tablet contains 5 mg Glibenclamide.

THERAPEUTIC ACTIVITY: Oral hypoglycemic agent.

APPROVED INDICATIONS: It is indicated for the treatment of non-Insulin dependent diabetes mellitus (TYPE 2).

WHEN SHOULD THE PREPARATION NOT BE USED?

In patients with known hypersensitivity to the active ingredient.

DO NOT USE THIS MEDICINE WITHOUT CONSULTING A PHYSICIAN: Do not take this drug if you are diabetic type I or in case of ketoacidose with or without coma. If you are pregnant or breastfeeding. If you are suffering or have suffered in the past from impaired function of the heart, kidney, and / or vascular system.

WARNINGS: If you are allergic to any type of food or medicine, inform your physician before commencing treatment with this medicine. This medicine may cause hypersensitivity when exposed to sunlight. Therefore, avoid exposure to sunlight. Do not drink alcohol while taking Glucocare. Blood and urine glucose should be monitored periodically.

DRUG INTERACTIONS: If you are taking any additional drugs or if you have just finished a course with another medicine, inform the attending physician, in order to prevent hazards or lack of efficacy arising from drug interactions, especially with medicines belonging to the following: Salicylates group, sulfonamides group, chloramphenicol, non-steroid anti-inflammatory drugs, probenecid, coumarins, and beta-adrenergic blocking agents.

SIDE EFFECTS: In addition to the desired effect of the medicine, adverse reactions may occur during the course of taking this medicine. Major side effects are hypoglycemia, sweating, blurred vision, and skin hypersensitivity. If any side effects not mentioned in this leaflet are experienced or if there has been any change in the general health, consult your physician.

RECOMMENDED DOSAGE: Dosage is according to physician's instructions only. Do not exceed the recommended dosage. This medicine is to be taken at specific time intervals as determined by the physician.

DIRECTIONS FOR USE: