

Uploaded BY AHMAD T JUNDI Upload by :Ahmad T Al-Jundi







Chapter 01

Lecture Outline

See separate *Image PowerPoint* slides for all figures and tables pre-inserted into PowerPoint without notes.

Chapter 1

Keys to the Study of Chemistry



Chapter 1: Keys to the Study of Chemistry

- 1.1 Some Fundamental Definitions
- 1.2 The Scientific Approach: Developing a Model
- 1.3 Chemical Problem Solving
- 1.4 Measurement in Scientific Study
- 1.5 Uncertainty in Measurement: Significant Figures



Chemistry

Chemistry is the study of *matter*, its *properties*,

the *changes* that matter undergoes,

and

the *energy* associated with these changes.



Definitions

Matter

anything that has both mass and volume
- the "stuff" of the universe: books, planets,
trees, professors, students

Composition

the types and amounts of simpler substances that make up a sample of matter

Properties

the characteristics that give each substance a unique identity



Physical Properties

properties a substance shows by itself without interacting with another substance

- color, melting point, boiling point, density

Chemical Properties


properties a substance shows as it interacts with, or transforms into, other substances

- flammability, corrosiveness



Figure 1.1 The distinction between physical and chemical change.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Oxygen gas Hydrogen gas

A Physical change:

Solid form of water becomes liquid form; composition does *not* change because particles are the same.

B Chemical change:


Electric current decomposes water into different substances (hydrogen and oxygen); composition *does* change because particles are different.



Visualizing Change on the Atomic Scale

PROBLEM: The scenes below represent an atomic-scale view of substance A undergoing two different changes. Decide whether each scene shows a physical or a chemical change.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




PLAN:

We need to determine what change is taking place. The numbers and colors of the little spheres that represent each particle tell its "composition". If the composition does not change, the change is physical, whereas a chemical change results in a change of composition.



SOLUTION:

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




Each particle of substance A is composed of one blue and two red spheres.

Sample B is composed of two different types of particles – some have two red spheres while some have one red and one blue.

As A changes to B, the chemical composition has changed.

 $A \rightarrow B$ is a **chemical change**.





Each particle of C is still composed of one blue and two red spheres, but the particles are closer together and are more organized. The composition remains unchanged, but the physical form is different.

 $A \rightarrow C$ is a **physical change**.




The States of Matter

A **solid** has a fixed shape and volume. Solids may be hard or soft, rigid or flexible.

A *liquid* has a varying shape that conforms to the shape of the container, but a fixed volume. A liquid has an *upper* surface.

A *gas* has no fixed shape or volume and therefore does not have a surface.

Figure 1.2 The physical states of matter.





Temperature and Change of State

- A change of state is a *physical* change.
 - Physical form changes, composition does not.
- Changes in physical state are reversible
 - by changing the temperature.
- A chemical change cannot simply be reversed by a change in temperature.



Distinguishing Between Physical and Chemical Change

PROBLEM: Decide whether each of the following processes is primarily a physical or a chemical change, and explain briefly:

- (a) Frost forms as the temperature drops on a humid winter night.
- (b) A cornstalk grows from a seed that is watered and fertilized.
- (c) A match ignites to form ash and a mixture of gases.
- (d) Perspiration evaporates when you relax after jogging.
- (e) A silver fork tarnishes slowly in air.

PLAN: "Does the substance change composition or just change form?"



SOLUTION:

(a) Frost forms as the temperature drops on a humid winter night.

physical change

(b) A cornstalk grows from a seed that is watered and fertilized.

chemical change

(c) A match ignites to form ash and a mixture of gases.

chemical change

(d) Perspiration evaporates when you relax after jogging.

physical change

(e) A silver fork tarnishes slowly in air.

chemical change



Energy in Chemistry

Energy is the ability to do work.

Potential Energy

is energy due to the *position* of an object.

Kinetic Energy is energy due to the movement of an object.

Total Energy = Potential Energy + Kinetic Energy



Energy Changes

Lower energy states are more stable and are favored over higher energy states.


Energy is neither created nor destroyed

- it is conserved
- and can be converted from one form to another.



Figure 1.3A Potential energy is converted to kinetic energy.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




A gravitational system. Potential energy is gained when a weight is lifted. It is converted to kinetic energy as the weight falls.


A gravitational system. The potential energy gained when a lifted weight is converted to kinetic energy as the weight falls.

A lower energy state is more stable.



Figure 1.3B Potential energy is converted to kinetic energy.





B A system of two balls attached by a spring. Potential energy is gained when the spring is stretched. It is converted to the kinetic energy of the moving balls as the spring relaxes.


A system of two balls attached by a spring. The potential energy gained by a stretched spring is converted to kinetic energy when the moving balls are released.

Energy is conserved when it is transformed.



Figure 1.3C Potential energy is converted to kinetic energy.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




C A system of oppositely charged particles. Potential energy is gained when the charges are separated. It is converted to kinetic energy as the attraction pulls the charges together.

A system of oppositely charged particles. The potential energy gained when the charges are separated is converted to kinetic energy as the attraction pulls these charges together.



Figure 1.3D Potential energy is converted to kinetic energy.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



D A system of fuel and exhaust. A fuel is higher in chemical potential energy than the exhaust. As the fuel burns, some of its potential energy is converted to the kinetic energy of the moving car.

A system of fuel and exhaust. A fuel is higher in chemical potential energy than the exhaust. As the fuel burns, some of its potential energy is converted to the kinetic energy of the moving car.



Figure 1.4 The scientific approach to understanding nature.

Observations

Hypothesis is revised if experimental results do not support it.

Hypothesis

Experiment

Model (Theory)

Model is altered if predicted events do not support it.

Further Experiment

Natural phenomena and measured events; can be stated as a *natural law* if universally consistent.

Tentative proposal that explains observations.

Procedure to test hypothesis; measures one variable at a time.

Set of conceptual assumptions that explains data from accumulated experiments; predicts related phenomena.

Tests predictions based on model



Chemical Problem Solving

- All measured quantities consist of
 - a number and a unit.
- Units are manipulated like numbers:

$$-3 \text{ ft } \times 4 \text{ ft} = 12 \text{ ft}^2$$

$$\frac{350 \text{ mi}}{7 \text{ h}} = \frac{50 \text{ mi}}{1 \text{ h}}$$
 or 50 mi.h



Conversion Factors

A *conversion factor* is a ratio of equivalent quantities used to express a quantity in different units.

The relationship 1 mi = 5280 ft gives us the conversion factor:

$$\frac{1 \text{ mi}}{5280 \text{ ft}} = \frac{5280 \text{ ft}}{5280 \text{ ft}} = 1$$



A conversion factor is chosen and set up so that all units cancel except those required for the answer.

PROBLEM: The height of the Angel Falls is 3212 ft. Express this quantity in miles (mi) if 1 mi = 5280 ft.

PLAN: Set up the conversion factor so that ft will cancel and the answer will be in mi.

SOLUTION:
$$3212 \text{ ft } \times \frac{1 \text{ mi}}{5280 \text{ ft}} = 0.6083 \text{ mi}$$



Systematic Approach to Solving Chemistry Problems

- State Problem
 - Clarify the known and unknown.
- Plan Suggest steps from known to unknown.

Prepare a visual summary of steps that includes conversion factors, equations, known variables.


- Solution
- Check
- Comment
- Follow-up Problem



Converting Units of Length

PROBLEM: To wire your stereo equipment, you need 325 centimeters (cm) of speaker wire that sells for \$0.15/ft. What is the price of the wire?

PLAN: We know the length (in cm) of wire and cost per length (\$/ft). We have to convert cm to inches and inches to feet. Then we can find the cost for the length in feet.





SOLUTION:

Length (in) = length (cm) x conversion factor

$$= 325 \text{ cm} \times \frac{1 \text{ in}}{2.54 \text{ cm}} = 128 \text{ in}$$

Length (ft) = length (in) x conversion factor

$$= 128 \frac{\text{in}}{\text{12 in}} = 10.7 \text{ ft}$$

Price (\$) = length (ft) x conversion factor

$$= 10.7 \text{ ft} \times \frac{\$ \ 0.15}{1 \text{ ft}} = \$ \ 1.60$$

1.4 Measurements in scientific study (SI Units)

Seven fundamental units (base Units)

Table 1.1 SI Base Units					
Physical Quantity (Dimension)	Unit Name	Unit Abbreviation			
Mass	kilogram	kg			
Length	meter	m			
Time	second	S			
Temperature	kelvin	K			
Electric current	ampere	A			
Amount of substance	mole	mol			
Luminous intensity	candela	cd			

All others are derived units and used as conversion factors (speed: m/s)



Table 1.2 Common Decimal Prefixes Used with SI Units

Prefix*	Prefix Symbol	Word	Conventional Notation	Exponential Notation
tera	Т	trillion	1,000,000,000,000	1×10 ¹²
giga	\mathbf{G}	billion	1,000,000,000	1×10^{9}
mega	M	million	1,000,000	1×10^{6}
kilo	k	thousand	1,000	1×10^{3}
hecto	h	hundred	100	1×10^{2}
deka	da	ten	10	1×10^{1}
	80 - 1 8	one	1	1×10^{0}
deci	d	tenth	0.1	1×10^{-1}
centi	c	hundredth	0.01	1×10^{-2}
milli	m	thousandth	0.001	1×10^{-3}
micro	μ	millionth	0.000001	1×10^{-6}
nano	n	billionth	0.000000001	1×10^{-9}
pico	p	trillionth	0.000000000001	1×10^{-12}
femto	f	quadrillionth	0.00000000000001	1×10^{-15}

^{*}The prefixes most frequently used by chemists appear in bold type.




Table 1.3 Common SI-English Equivalent Quantities


Quantity	SI to English Equivalent	English to SI Equivalent
Length	1 km = 0.6214 mile 1 m = 1.094 yd 1 m = 39.37 inches 1 cm = 0.3937 inch	1 mi = 1.609 km 1 yd = 0.9144 m 1 ft = 0.3048 m 1 in = 2.54 cm
Volume	1 cubic meter (m³) = 35.31 ft³ 1 dm³ = 0.2642 gal 1 dm³ = 1.057 qt 1 cm³ = 0.03381 fluid ounce	1 ft ³ = 0.02832 m ³ 1 gal = 3.785 dm ³ 1 qt = 0.9464 dm ³ 1 qt = 946.4 cm ³
Mass	1 kg = 2.205 lb 1 g = 0.03527 ounce (oz)	1 fluid ounce = 29.57 cm ³ 1 lb = 0.4536 kg 1 oz = 28.35 g



Figure 1.5 Common laboratory volumetric glassware.

Copyright @The McGraw-Hill Companies, Inc. Permission required for reproduction or display.








Converting Units of Volume

PROBLEM: A graduated cylinder contains 19.9 mL of water. When a small piece of galena, an ore of lead, is added, it sinks and the volume increases to 24.5 mL. What is the volume of the piece of galena in cm³ and in L?

PLAN: The volume of the galena is equal to the difference in the volume of the water before and after the addition.





SOLUTION:

(24.5 - 19.9) mL = volume of galena = 4.6 mL

4.6 mL x
$$\frac{1 \text{ cm}^3}{1 \text{ mL}}$$
 = 4.6 cm³

4.6 mL x
$$\frac{10^{-3} L}{1 mL}$$
 = 4.6 x 10⁻³ L

MENTS-HUB.com

Converting Units of Mass

PROBLEM: Many international computer communications are carried out by optical fibers in cables laid along the ocean floor. If one strand of optical fiber weighs 1.19 x 10⁻³ lb/m, what is the mass (in kg) of a cable made of six strands of optical fiber, each long enough to link New York and Paris (8.94 x 10³ km)?

PLAN: The sequence of steps may vary but essentially we need to find the length of the entire cable and convert it to mass.

```
length (km) of fiber

1 km = 10<sup>3</sup> m

length (m) of fiber

1 m = 1.19 x 10<sup>-3</sup> lb

mass (lb) of fiber

6 fibers = 1 cable

2.205 lb = 1 kg

mass (kg) of cable
```

SOLUTION:

$$8.84 \times 10^3 \frac{\text{km}}{1 \frac{\text{km}}{\text{m}}} = 8.84 \times 10^6 \text{ m}$$

$$8.84 \times 10^6 \,\text{m} \times \, \frac{1.19 \times 10^{-3} \,\text{lb}}{1 \,\text{m}} = 1.05 \times 10^4 \,\text{lb}$$

$$\frac{1.05 \times 10^4 \text{ lb}}{1 \text{ fiber}} \times \frac{6 \text{ fibers}}{1 \text{ cable}} = 6.30 \times 10^4 \text{ lb/cable}$$

$$\frac{6.30 \times 10^4 \text{ lb}}{1 \text{ cable}} \times \frac{1 \text{ kg}}{2.205 \text{ lb}} = 2.86 \times 10^4 \text{ kg/cable}$$

Density

density =
$$\frac{\text{mass}}{\text{volume}}$$

At a given temperature and pressure, the density of a substance is a characteristic physical property and has a specific value.

Substance	Physical State	Density (g/cm³)
Hydrogen	gas	0.0000899
Oxygen	gas	0.00133
Grain alcohol	liquid	0.789
Water	liquid	0.998
Table salt	solid	2.16
Aluminum	solid	2.70
Lead	solid	11.3
Gold	solid	19.3


^{*}At room temperature (20°C) and normal atmospheric pressure (1 atm).



Calculating Density from Mass and Length

PROBLEM: Lithium, a soft, gray solid with the lowest density of any metal, is a key component of advanced batteries. A slab of lithium weighs 1.49x10³ mg and has sides that are 20.9 mm by 11.1 mm by 11.9 mm. Find the density of lithium in g/cm³.

PLAN: Density is expressed in g/cm³ so we need the mass in g and the volume in cm³.





SOLUTION:

$$1.49 \times 10^3 \, \text{mg} \times \, \frac{1 \, \text{g}}{10^3 \, \text{mg}} = 1.49 \, \text{g}$$

$$20.9 \text{ mm x } \frac{1 \text{ cm}}{10 \text{ mm}} = 2.09 \text{ cm}$$

Similarly the other sides will be 1.11 cm and 1.19 cm, respectively.

Volume =
$$2.09 \times 1.11 \times 1.19 = 2.76 \text{ cm}^3$$

density of Li =
$$\frac{1.49 \text{ g}}{2.76 \text{ cm}^3}$$
 = **0.540 g/cm**³

Temperature (T): is a measure of how hot or cold one object is relative to another. (Thermometer)

Heat: is energy that flows from the object of higher (T) to object of lower (T).


Kelvin (K) - The "absolute temperature scale" begins at absolute zero and has only positive values. Note that the kelvin is not used with the degree sign (°).

Celsius (°C) - The Celsius scale is based on the freezing and boiling points of water. This is the temperature scale used most commonly around the world. The Celsius and Kelvin scales use the **same size degree** although their starting points differ.

Fahrenheit (°F) — The Fahrenheit scale is commonly used in the US. The Fahrenheit scale has a different degree size and different zero points than both the Celsius and Kelvin scales.



Figure 1.6 Freezing and boiling points of water in the Celsius, Kelvin (absolute) and Fahrenheit scales.





Temperature Conversions

T (in K) = T (in
$$^{\circ}$$
C) + 273.15
T (in $^{\circ}$ C) = T (in K) - 273.15

T (in °F) =
$$\frac{9}{5}$$
 T (in °C) + 32

T (in °C) = [T (in °F) – 32]
$$\frac{5}{9}$$



Converting Units of Temperature

PROBLEM: A child has a body temperature of 38.7°C, and normal body temperature is 98.6°F. Does the child have a fever? What is the child's temperature in kelvins?

PLAN: We have to convert °C to °F to find out if the child has a fever. We can then use the °C to Kelvin relationship to find the temperature in Kelvin.

SOLUTION:

$$\frac{9}{5}$$
(38.7 °C) + 32 = **101.7** °F

Yes, the child has a fever.

Converting from °C to K



Significant Figures


Every measurement includes some *uncertainty*. The *rightmost* digit of any quantity is always *estimated*.

The recorded digits, both certain and uncertain, are called *significant figures*.

The greater the number of significant figures in a quantity, the greater its certainty.



Figure 1.7 The number of significant figures in a measurement.





Determining Which Digits are Significant

All digits are significant

- except zeros that are used <u>only</u> to position the decimal point.
- Make sure the measured quantity has a decimal point.
- Start at the left and move right until you reach the first nonzero digit.
- Count that digit and every digit to its right as significant.



- Zeros that end a number are significant as long as a decimal point is present.
- 1.030 mL has 4 significant figures.
- 5300. L has 4 significant figures.
- If there is no decimal point is present
 - zeros at the end of the number are <u>not</u> significant.
 unless exponential notation clarify the quantity
- 5300 L has only 2 significant figures
- 5.300 *10³ has 4 significant figures

Determining the Number of Significant Figures

PROBLEM:

For each of the following quantities, underline the zeros that are significant figures (sf), and determine the number of significant figures in each quantity. For (d) to (f), express each in exponential notation first.

- (a) 0.0030 L
- **(b)** 0.1044 g **(c)** 53,069 mL

- (d) 0.00004715 m (e) 57,600. s (f) 0.0000007160 cm³

PLAN:

We determine the number of significant figures by counting digits, paying particular attention to the position of zeros in relation to the decimal point, and underline zeros that are significant.



SOLUTION:

- (a) 0.003<u>0</u> L has **2 sf**
- **(b)** 0.1<u>0</u>44 g has **4 sf**
- (c) 53,<u>0</u>69 mL has **5 sf**
- (d) $0.00004715 \text{ m} = 4.715 \times 10^{-5} \text{ m}$ has 4 sf
- (e) 57,600. $s = 5.7600 \times 10^4 \text{ s}$ has 5 sf
- (f) $0.0000007160 \text{ cm}^3 = 7.160 \times 10^{-7} \text{ cm}^3 \text{ has 4 sf}$

Rules for Significant Figures in Calculations

1. *For multiplication and division*. The answer contains the same number of significant figures as there are in the measurement with the **fewest significant figures**.

Multiply the following numbers:

9.2 cm x 6.8 cm x 0.3744 cm = 23.4225 cm³ = 23 cm³



Rules for Significant Figures in Calculations

2. *For addition and subtraction*. The answer has the same number of decimal places as there are in the measurement with the fewest decimal places.

Example: adding two volumes 83.5 mL + 23.28 mL

106.78 mL = 106.8 mL

Example: subtracting two volumes 865.9 mL

- 2.8121 mL

863.0879 mL = **863.1** mL



Rules for Rounding Off Numbers

- 1. If the <u>digit removed</u> is *more than 5*, the preceding number increases by 1.
- 5.379 rounds to 5.38 if 3 significant figures are retained.

- 2. If the <u>digit removed</u> is *less than 5*, the preceding number is unchanged.
- 0.2413 rounds to 0.241 if 3 significant figures are retained.



3. If the <u>digit removed</u> is 5 followed by <u>zeros or</u> <u>with no following digits</u>, the preceding number <u>increases by</u> 1 if it is <u>odd</u> and remains <u>unchanged</u> if it is <u>even</u>.

17.75 rounds to 17.8, but 17.65 rounds to 17.6.

If the *5 is followed by <u>other nonzero digits</u>*, rule 1 is followed:

17.6500 rounds to 17.6, but 17.6513 rounds to 17.7

4. Be sure to carry two or more additional significant figures through a multistep calculation and round off the *final* answer *only*.



Exact Numbers

Exact numbers have no uncertainty associated with them.

Numbers may be exact by definition:

```
1000 \text{ mg} = 1 \text{ g}
```

60 min = 1 hr

2.54 cm = 1 in

Numbers may be exact by count: exactly 26 letters in the alphabet

Exact numbers do not limit the number of significant digits in a calculation.



Significant Figures and Rounding

PROBLEM: Perform the following calculations and round each answer to the correct number of significant figures:

(a)
$$\frac{16.3521 \text{ cm}^2 - 1.448 \text{ cm}^2}{7.085 \text{ cm}}$$
 (b) $\frac{4.80 \times 10^4 \text{ mg}}{1000 \text{ mg}}$ $\frac{1 \text{ g}}{1000 \text{ mg}}$

PLAN: We use the rules for rounding presented in the text: **(a)** We subtract before we divide. **(b)** We note that the unit conversion involves an exact number.

SOLUTION:

(a)
$$\frac{16.3521 \text{ cm}^2 - 1.448 \text{ cm}^2}{7.085 \text{ cm}} = \frac{14.904 \text{ cm}^2}{7.085 \text{ cm}} = 2.104 \text{ cm}$$

(b)
$$\frac{4.80 \times 10^4 \text{ mg}}{11.55 \text{ cm}^3} = \frac{48.0 \text{ g}}{11.55 \text{ cm}^3} = 4.16 \text{ g/ cm}^3$$

Precision, Accuracy, and Error

Precision refers to how close the measurements in a series are to each other.

Accuracy refers to how close each measurement is to the actual value.

Systematic error produces values that are either all higher or all lower than the actual value.

This error is part of the experimental system.

Random error produces values that are both higher and lower than the actual value.



Figure 1.9 Precision and accuracy in a laboratory calibration.








Figure 1.9 Precision and accuracy in the laboratory. continued





UPLOADED BY AHMAD T JUNDI

Upload By: Ahmad T Al -Jundi







Chapter 2

The Periodic Table and Some
Properties
of the Elements





UPLO Definitions for Components of Matter

Matter can be classified in to three types based on it composition: <u>elements</u>, <u>compounds</u> and <u>mixtures</u>.

Substance: is matter whose composition is fixed. Mixtures?

Element - the simplest type of substance with unique physical and chemical properties. *An element consists of only one type of atom.* It cannot be broken down into any simpler substances by physical or chemical means. Cu, Ag, N_2 , O_2





Figure 2.1

Molecule - a structure that consists of two or more atoms that are chemically bound together and thus behaves as an independent unit.



 $H_2O, O_3, CO_2, NH_3, O_2$








Definitions for Components of Matter

Compound - a substance composed of two or more elements which are chemically combined.

-Elements are present in fixed mass ratio.


NH₃ 14 g N with 3 g H

-Compound properties are different from its elements NaCl



H₂O + NaCl

- -Components of mixtures can vary in their parts.
- Mixtures retains many of the properties of its components
 - -Can be separated to its components by physical **D**



Uploaded By: mariam Qadah



Mixtures

A *heterogeneous mixture* has one or more visible boundaries between the components.


A *homogeneous mixture* has no visible boundaries because the components are mixed as individual atoms, ions, and molecules.

A homogeneous mixture is also called a *solution*. Solutions in water are called *aqueous solutions*.






Figure 2.17 The distinction between mixtures and compounds.



A physical mixture of Fe and S_8 can be separated using a magnet.



Fe and S have reacted chemically to form the compound FeS. The elements cannot be separated by physical means.





Table 2.1 Some Properties of Sodium, Chlorine, and Sodium **Chloride**

Property	Sodium +	Chlorine →	Sodium Chloride
Melting point	97.8°C	-101°C	801°C
Boiling point	881.4°C	-34°C	1413°C
Color	Silvery	Yellow-green	Colorless (white)
Density	0.97 g/cm ³	0.0032 g/cm ³	2.16 g/cm ³
Behavior in water	Reacts	Dissolves slightly	Dissolves freely












Distinguishing Elements, Compounds, and Mixtures at the Atomic Scale

PROBLEM:

The following scenes represent an atomic-scale view of three samples of matter. Describe each sample as an element, compound, or mixture.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



PLAN:


A sample that contains only one type of particle is either an element or a compound. The particles of an element consist of only one type of atom whereas the particles of a compound have two or more types of atom bonded together.





SOLUTION:

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Sample (a) contains three different types of particles and is therefore a **mixture**.

Sample (b) contains only one type of particle and each particle has only one atom. This is an **element**.

Sample (c) contains only one type of particle, each of which contains two different types of atoms. This is a **compound**.





Observations that lead to an atomic view of matter


- 1- Law of mass conservation.
- 2- Law of definite (constant) composition.
- 3- Law of multiple proportion.





Law of Mass Conservation

The total mass of substances present does not change during a chemical reaction. (Matter cannot be created or destroyed).








Law of Definite (or Constant) Composition

No matter the source, a particular compound is composed of the same elements in the same parts (fractions) by mass.(mass fraction).







Calcium carbonate

Analysis	by	Mas	S
(grams	/20 .	.0 g)	

8.0 g calcium

2.4 g carbon

9.6 g oxygen

20.0 g

Mass Fraction (parts/1.00 part)

0.40 calcium

0.12 carbon

0.48 oxygen

1.00 part by mass

Percent by Mass (parts/100 parts)

40% calcium

12% carbon

48% oxygen

100% by mass

Mass of elements in sample = mass of compound in sample \boldsymbol{x}

(mass of element in compound)

mass of compound





Calculating the Mass of an Element in a Compound

PROBLEM: Analysis of 84.2 g of the uranium containing compound pitchblende shows it is composed of 71.4 g of uranium, with oxygen as the only other element. How many grams of uranium can be obtained from 102 kg of pitchblende?

PLAN: The mass ratio of uranium/pitchblende is the same no matter the source. We can use the ratio to find the answer.

mass (kg) of pitchblende

mass ratio of U in pitchblende

mass (kg) of uranium

1 kg = 1000 g

mass (g) of uranium





SOLUTION:

mass (kg) of uranium =

mass (kg) pitchblende x mass (kg) uranium in pitchblende mass (kg) pitchblende

= 102 kg pitchblende x 71.4 kg uranium = 86.5 kg uranium 84.2 kg pitchblende

86.5 kg uranium x
$$\frac{1000 \text{ g}}{1 \text{ kg}}$$
 = 8.65 x 10⁴ g uranium







Law of Multiple Proportions

If elements A and B react to form two compounds, the different masses of B that combine with a fixed mass of A can be expressed as a ratio of small whole numbers.

Example: Carbon Oxides A & B

Carbon Oxide I: 57.1% oxygen and 42.9% carbon





Carbon Oxide II: 72.7% oxygen and 27.3% carbon





Assume that you have 100 g of each compound.


In 100 g of each compound:

g C = 42.9 g for oxide I & 27.3 g for oxide II

For oxide I:
$$\frac{g O}{g C} = \frac{57.1}{42.9} = 1.33$$

For oxide II:
$$\frac{g O}{g C} = \frac{72.7}{27.3} = 2.66$$

$$\frac{2.66 \text{ g O/g C in II}}{1.33 \text{ g O/g C in I}} = \frac{2}{1}$$



In two compounds of same the elements, the mass fraction of one element relative to the other element changes in increments based on ratios of small whole number





Dalton postulated that:

- All matter consists of atoms; tiny indivisible particles of an element that cannot be created or destroyed.
- 2. Atoms of one element *cannot* be converted into atoms of another element.
- Atoms of an element are *identical* in mass and other properties and are different from the atoms of any other element.
- 4. Compounds result from the chemical combination of a specific ratio of atoms of different elements.





explains the mass laws

Mass conservation

Atoms cannot be created or destroyed **postulate 1**

or converted into other types of atoms. postulate 2

Since every atom has a fixed mass, postulate 3

during a chemical reaction the same atoms are present but in different combinations; therefore there is no mass change overall.



explains the mass laws

Definite composition

Atoms are combined in compounds in **postulate 4** specific ratios and each atom has a specific mass. **postulate 3**

Each element constitutes a fixed fraction of the total mass in a compound.



explains the mass laws

Multiple proportions

Atoms of an element have the same mass *postulate 3* and atoms are indivisible. *postulate 1*

When different numbers of atoms of elements combine, they must do so in ratios of small, whole numbers.








Visualizing the Mass Laws

PROBLEM:

The following scene represents an atomic-scale view of a chemical reaction. Which of the mass laws (mass conservation, definite composition, or multiple proportions) is (are) illustrated?






PLAN:

Note the numbers, types and combination of atoms before and after the reaction.



SOLUTION:



There are 7 purple and 9 green atoms both before and after the reaction. Mass is therefore conserved.

After the reaction some purple atoms remain unreacted, but some have combined with green atoms to form a compound. Each particle of this compound contains 1 purple and 2 green atoms – the composition is constant, illustrating the law of definite composition.


Only one compound forms, law of multiple proportions does not pertain.





Observations that established the properties of cathode rays.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



J. J Thomson, (1897) was able to measure the electron (Mass/charge) ratio = -5.686x10⁻¹² Kg/C

Note: Dalton atoms are not divisible??

Observation	Conclusion
Ray bends in magnetic field.	Ray consists of charged particles.
Ray bends toward positive plate in electric field.	Ray consists of negative particles.
Ray is identical for any cathode.	These particles are found in ALL matter.






Figure 2.4

Millikan's oil-drop experiment for measuring an electron's charge. (1909)

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Electron charge = $-1.60218 \times 10^{-19} \text{ C}$ (coulomb)





Millikan's findings were used to calculate the mass of an electron.

determined by J.J. Thomson and others

=
$$(-5.686 \times 10^{-12} \text{ kg/C}) \times (-1.602 \times 10^{-19} \text{ C})$$


$$= 9.109 \times 10^{-31} \text{ kg} = 9.109 \times 10^{-28} \text{ g}$$





Figure 2.5 of the atomic nucleus. (1910)

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Rutherford: Atoms are empty space occupied by electrons, but in the center a tiny region he called nucleus contains all the positive particles and called them protons.


In 1932 James Chadwick discovered the Neutrons.





Figure 2.6 General features of the atom.

The atom is an electrically neutral, spherical entity composed of a positively charged central nucleus surrounded by one or more negatively charged electrons.



The atomic nucleus consists of protons and neutrons.





Table 2.2 Properties of the Three Key Subatomic Properties

	Ch	arge	Mas	s	
Name (Symbol)	Relative	Absolute (C)*	Relative (amu) [†]	Absolute (g)	Location in Atom
Proton (p+)	1+	+1.60218x10 ⁻¹⁹	1.00727	1.67262x10 ⁻²⁴	Nucleus
Neutron (n ⁰)	0	0	1.00866	1.67493x10 ⁻²⁴	Nucleus
Electron (e ⁻)	1-	-1.60218x10 ⁻¹⁹	0.00054858	9.10939x10 ⁻²⁸	Outside nucleus

^{*}The coulomb (C) is the SI unit of charge.

[†] The atomic mass unit (amu) equals 1.66054x10⁻²⁴ g.






Atomic Number, Mass Number, and Atomic Symbol


Figure 2.7

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




X = Atomic symbol of the element














Isotopes

Isotopes are atoms of an element with the same number of *protons*, but a different number of *neutrons*.

Isotopes have the same atomic number, but a different mass number.



98.8% from C atoms have 6 n⁰ 1.11% have 7 n⁰ Less than 0.001% have 8 n⁰




Figure 2.7





Determining the Number of Subatomic Particles in the Isotopes of an Element

PROBLEM: Silicon (Si) has three naturally occurring isotopes: ²⁸Si, ²⁹Si, and ³⁰Si. Determine the number of protons, neutrons, and electrons in each silicon isotope.

PLAN: The mass number (A) is given for each isotope and is equal to the number of protons + neutrons. The atomic number Z, found on the periodic table, equals the number of protons. The number of neutrons = A - Z, and the number of electrons equals the number of protons for a neutral atom.

SOLUTION: The atomic number of silicon is 14; therefore

²⁸Si has 14p⁺, 14e⁻ and 14n⁰ (28-14)

²⁹Si has 14p⁺, 14e⁻ and 15n⁰ (29-14)


³⁰Si has 14p⁺, 14e⁻ and 16n⁰ (30-14)





The Mass Spectrometer and Its Data

Mass Spectrometer measures the mass ratio and mass to charge ratio





Atomic Symbol

C Carbon

Na (Natrium) Sodium

Note: The chemical properties of an element are determined by the number of electrons or number of protons. So having Isotopes of an element will not change its chemical properties **much.**

Mass of an atom

Atomic mass unit (amu) or Dalton (Da)

Today (u) only

$$amu = \frac{1}{12} \quad mass of ^{12}C$$

So ¹²C has a mass of 12 u





Calculating the Atomic Mass of an Element

PROBLEM: Silver (Ag, Z = 47) has two naturally occurring isotopes,

¹⁰⁷Ag and ¹⁰⁹Ag. From the mass spectrometric data

provided, calculate the atomic mass of Ag.

<u>lsotope</u>	<u>Mass (amu)</u>	<u> Abundance (%)</u>
¹⁰⁷ Ag	106.90509	51.84
¹⁰⁹ Ag	108.90476	48.16

PLAN: Find the weighted average of the isotopic masses.

mass (g) of each isotope

multiply by fractional abundance of each isotope

portion of atomic mass from each isotope

add isotopic portions

atomic mass





SOLUTION:

```
mass portion from ^{107}Ag = 106.90509 amu x 0.5184 = 55.42 amu
```

mass portion from
109
Ag = 108.90476 amu x $0.4816 = 52.45$ amu

atomic mass of Ag = 55.42amu + 52.45amu

= 107.87amu

The atomic mass is average value

No individual atom has a mass of 107.87 amu





ELEMENTS: A FIRST LOOK AT THE PERIODIC TABLE.

1870 65 elements were known

Today more than 116 elements


Dmitri Mendeleev and Meyer discovered the periodic table separately in 1869

Dmitri Mendeleev organized elements, he listed the elements by increasing atomic mass.





Figure 2.9 The modern periodic table.



- **7** Periods-(horizontal rows)
- 8 groups(A and B)- (vertical columns)
- -Metal (blue) about three quarters of elements are metal, generally they are shiny solid at room temperature (mercury is liquid)
- -Nonmetals (yellow) generally gases, brittle solid at room temperature (bromine is only liquid).
- Metalloids-semimetals(green)
- -Elements in a group have similar chemical properties and elements in a period have different chemical properties.
- There is gradation in properties from left to right and top to bottom.





UPLOADED BY AHMAD TUND UCTION to bonding





Figure 2.11 Factors that influence the strength of ionic bonding.



Predicting the Ion an Element Forms

PROBLEM: Predict the monoatomic ion formed by each of the following elements:

(a) Iodine (Z = 53) (b) Calcium (Z = 20) (c) Aluminum (Z = 13)

PLAN: Use Z to find the element on the periodic table and see where it lies relative to its **nearest** noble gas.

SOLUTION:

- (a) Iodine is a nonmetal in Group 7A(17). It gains one electron to have the same number of electrons as ₅₄Xe.

 The ion is I
- (b) Calcium is a metal in Group 2A(2). It loses two electrons to have the same number of electrons as ₁₈Ar.

 The ion is Ca²⁺
- (c) Aluminum is a metal in Group 3A(13). It loses three electrons to have the same number of electrons as ₁₀Ne. The ion is AI³⁺






Formation of a covalent bond between two H atoms.

Figure 2.12


Covalent bonds form when elements share electrons, which usually occurs between nonmetals.



Atoms far apart: No interactions.



Atoms closer: Attractions (green arrows) between nucleus of one atom and electron of the other increase. Repulsions between nuclei and between electrons are very weak.



Optimum distance: H_2 molecule forms because attractions (green arrows) balance repulsions (red arrows).





Wolecules and lons

Covalent V.S ionic substance

- -Most covalent substances consist of molecules. Covalent bonding involves the mutual attraction between two positively charged (nuclei) and two negatively charges (electrons).
- -lonic bonding involves he mutual attraction among positive and negative ions.
- **Molecule** the basic unit of an element or covalent compound, consisting of two or more atoms bonded by the sharing of electrons.(H₂O)
 - **Ion** a single atom or covalently bonded group of atoms that has an overall electrical charge.
 - There are *no molecules* in an ionic compound.



Figure 2.13 Elements that occur as molecules.









Figure 2.14 The carbonate ion in calcium carbonate.



A polyatomic ion consists of two or more atoms covalently bonded together and has an overall charge.

In many reactions the polyatomic ion will remain together as a unit.



Chemical Formulas

- A chemical formula consists of
 - element symbols with
 - numerical subscripts.
- The chemical formula indicates the
 - type and number of each atom present
 - in the smallest unit of a substance.



Naming Binary Ionic Compounds

For all ionic compounds, the name and formula lists the cation first and the anion second.

In a **binary ionic** compound, both the cation and the anion are monatomic.

The name of the cation is the **same** as the name of the metal. Many metal names end in -ium.

The anion is named by adding the suffix *-ide* to the root of the nonmetal name.

Calcium and bromine form calcium bromide.





Table 2.3 Common Monatomic Ions*

Charge	Cations Formula	Name
+1	H ⁺ Li ⁺ Na ⁺ K ⁺ Cs ⁺ Ag ⁺	hydrogen lithium sodium potassium cesium silver
+2	Mg ²⁺ Ca ²⁺ Sr ²⁺ Ba ²⁺ Zn ²⁺ Cd ²⁺	magnesium calcium strontium barium zinc cadmium
+3	Al ³⁺	aluminum

	Anions	
Charge	Formula	Name
-1	H ⁻ F ⁻ CI ⁻ Br ⁻	hydride fluoride chloride bromide iodide
-2	O ²⁻ S ²⁻	oxide sulfide
-3	N ³⁻	nitride

^{*}Listed by charge; those in boldface are most common.





Figure 2.15 Some common monatomic ions of the elements.

		1A (1)																7A (17)	8A (18)
	1	H+	2A (2)											3A (13)	4A (14)	5A (15)	6A (16)	Н-	
	2	Li+														N ³⁻	O ²⁻	F	
	3	Na+	Mg ²⁺	3B (3)	4B (4)	5B (5)	6B (6)	7B (7)	(8)	— 8B — (9)	(10)	1B (11)	2B (12)	Al ³⁺			S ²⁻	CI-	
Period	4	K ⁺	Ca ²⁺				Cr ²⁺ Cr ³⁺	Mn ²⁺	Fe ²⁺	Co ²⁺		Cu ⁺	Zn ²⁺					Br-	
	5	Rb ⁺	Sr ²⁺									Ag+	Cd ²⁺		Sn ²⁺ Sn ⁴⁺			г	
	6	Cs ⁺	Ba ²⁺										Hg ₂ ²⁺ Hg ²⁺		Pb ²⁺ Pb ⁴⁺				
	7																		

Most main-group elements form one monatomic ion. Most transition elements form two monatomic ions.



Naming Binary Ionic Compounds

PROBLEM: Name the ionic compound formed from each of the following pairs of elements:

(a) magnesium and nitrogen (b) iodine and cadmium

(c) strontium and fluorine (d) sulfur and cesium

PLAN: Use the periodic table to decide which element is the metal and which the nonmetal. The metal (cation) is named first and the suffix-*ide* is added to the root of the non-metal name.

SOLUTION:

(a) magnesium nitride (b) cadmium iodide

(c) strontium fluoride (d) cesium sulfide





Determining Formulas of Binary Ionic Compounds

PROBLEM: Write empirical formulas for each of the compounds named in Sample Problem 2.7.

(a) magnesium nitride (b) cadmium iodide

(c) strontium fluoride (d) cesium sulfide

PLAN: A compound is neutral. We find the smallest number of each ion that will produce a neutral formula. These numbers appear as *right subscripts* to the relevant element symbol.

SOLUTION:

(a) Mg^{2+} and N^{3-} ; three $Mg^{2+}(6+)$ and two $N^{3-}(6^{-})$; Mg_3N_2

(b) Cd^{2+} and I^{-} ; one $Cd^{2+}(2+)$ and two $I^{-}(2^{-})$; Cdl_2

(c) Sr^{2+} and F^{-} ; one $Sr^{2+}(2+)$ and two $F^{-}(2^{-})$; SrF_{2}

(d) Cs⁺ and S²⁻; two Cs⁺(2+) and one S²⁻(2⁻); Cs₂S





S2 ⊌5 2NTS-HUB.com

Table 2.4 Some Metals That Form More Than One Monatomic Ion*

Element	Ion Formula	Systematic Name	Common Name
Chromium	Cr ²⁺	chromium(II)	chromous
	Cr ³⁺	chromium(III)	chromic
Cobalt	Co ²⁺	cobalt(II)	
	Co ³⁺	cobalt(III)	
Copper	Cu⁺	copper(I)	cuprous
	Cu ²⁺	copper(II)	cupric
Iron	Fe ²⁺	iron(II)	ferrous
	Fe ³⁺	iron(III)	ferric
Lead	Pb ²⁺	lead(II)	
	Pb ⁴⁺	lead(IV)	
Mercury	Hg_2^{2+}	mercury (I)	mercurous
	Hg ²⁺	mercury (II)	mercuric
Tin	Sn ²⁺	tin(II)	stannous
	Sn ⁴⁺	tin(IV)	stannic

^{*}Listed alphabetically by metal name; the ions in boldface are most common.

Uploaded By: mariam Qadah

Determining Names and Formulas of lonic Compounds of Elements That Form More Than One Ion

PROBLEM: Give the systematic name for each formula or the formula for each name for the following compounds:

(a) tin(II) fluoride (b) Crl₃ (c) ferric oxide (d) CoS

PLAN: Find the smallest number of each ion that will produce a neutral formula.

SOLUTION:

- (a) Tin(II) is Sn^{2+} ; fluoride is F^- ; so the formula is SnF_2 .
- (b) The anion I⁻ is iodide; 3I⁻ means that Cr (chromium) is +3. CrI₃ is chromium(III) iodide.
- (c) Ferric is a common name for Fe³⁺; oxide is O²⁻; therefore the formula is **Fe₂O₃**.
- (d) Co is cobalt; the anion S²⁻ is sulfide; the compound is **cobalt(II)** sulfide.



Table 2.5	Common	Polyat	tomic	lons*

Formula	Name	Formula	Name			
Cations						
NH ₄ ⁺	ammonium	H ₃ O ⁺	hydronium			
Common Anions						
CH ₃ COO-	acetate	CO ₃ 2-	carbonate			
CN-	cyanide	HCO ₃ -	bicarbonate			
OH-	hydroxide	CrO ₄ 2-	chromate			
CIO-	hypochlorite	Cr ₂ O ₇ ²⁻	dichromate			
CIO ₂ -	chlorite	O ₂ ² -	peroxide			
CIO ₃ -	chlorate	PO ₄ 3-	phosphate			
NO_2^{-}	nitrite	HPO ₄ 2-	hydrogen phosphate			
NO ₃ -	nitrate	SO ₃ ²⁻	sulfite			
MnO ₄ -	permanganate	SO ₄ 2-	sulfate			
*Bold face ions are most common. (partial table)						



Figure 2.16 Naming oxoanions

	Prefix	Root	Suffix	Exa	mple
atoms	per	root	ate	CIO ₄	perchlorate
0		root	ate	CIO ₃	chlorate
No. of		root	ite	CIO ₂ -	chlorite
	hypo	root	ite	CIO ⁻	hypochlorite



Table 2.6 Numerical Prefixes* for Hydrates and Binary Covalent Compounds

Number	Prefix	Number	Prefix	Number	Prefix
1	mono-	4	tetra-	8	octa-
2	di-	5	penta-	9	nona-
3	tri-	6	hexa-	10	deca-
		7	hepta-		



Determining Names and Formulas of Ionic Compounds Containing Polyatomic Ions

PROBLEM: Give the systematic name for each formula or the formula for each name for the following compounds:

- (a) $Fe(ClO_4)_2$ (b) sodium sulfite (c) $Ba(OH)_2 \cdot 8H_2O$

Remember to use parentheses when more than one unit PLAN: of a particular polyatomic ion is present in the compound.

SOLUTION: (a) ClO₄⁻ is perchlorate; Fe must have a 2+ charge since there are 2 ClO_4^- ions. This is **iron(II) perchlorate**.

- **(b)** The anion sulfite is SO₃²⁻; therefore you need 2 Na⁺ for each sulfite. The formula is Na₂SO₃.
- (c) The ionic compound is barium hydroxide. When water is included in the formula, we use the term "hydrate" and a prefix that indicates the number of molecules of H₂O. This compound is barium hydroxide octahydrate.





Recognizing Incorrect Names and Formulas of Ionic Compounds

PROBLEM: There is an error in the second part of each statement. Provide the correct name or formula in each case.

- (a) Ba($C_2H_3O_2$)₂ is called barium diacetate.
- **(b)** Sodium sulfide has the formula (Na)₂SO₃.
- (c) Iron(II) sulfate has the formula $Fe_2(SO_4)_3$.
- (d) Cesium carbonate has the formula $Cs_2(CO_3)$.

SOLUTION:

- (a) The charge of Ba²⁺ *must* be balanced by *two* C₂H₃O₂⁻ ions. The prefix "di" is not required and is not used in this way when naming ionic compounds. The correct name is simply **barium acetate**.
- (b) An ion of a single element does not need parentheses, and sulfide is S²⁻, not SO₃²⁻. The correct formula is Na₂S.





- (c) Sulfate or SO₄²⁻ has a 2⁻ charge, and only *one* Fe²⁺ is needed to form a neutral compound. The formula should be **FeSO₄**.
- (d) The parentheses are unnecessary, since only *one* CO_3^{2-} ion is present. The correct formula is Cs_2CO_3 .





Naming Acids

1) *Binary acid* solutions form when certain gaseous compounds dissolve in water.

For example, when gaseous hydrogen chloride (HCI) dissolves in water, it forms a solution called hydrochloric acid.

Prefix *hydro-* + anion nonmetal *root* + suffix *-ic* + the word *acid -* hydro + chlor + ic + acid hydrochloric acid

2) Oxoacid names are similar to those of the oxoanions, except for two suffix changes:

-ate in the anion becomes -ic in the acid

-ite in the anion becomes -ous in the acid

The oxoanion prefixes *hypo-* and *per-* are retained. Thus,

BrO₄ is *perbromate*, and HBrO₄ is *perbromic* acid;

 IO_2^- is *iodite*, and HIO_2 is *iodous acid*.





Determining Names and Formulas of Anions and Acids

PROBLEM: Name the following anions and give the name and formula of the acid derived from each:

(a) Br⁻ (b) IO_3^- (c) CN^- (d) SO_4^{2-} (e) NO_2^-

SOLUTION:

- (a) The anion is **bromide**; the acid is **hydrobromic acid**, **HBr**.
- (b) The anion is **iodate**; the acid is **iodic acid**, HIO₃.
- (c) The anion is cyanide; the acid is hydrocyanic acid, HCN.
- (d) The anion is **sulfate**; the acid is **sulfuric acid**, H₂SO₄.
- (e) The anion is **nitrite**; the acid is **nitrous acid**, **HNO**₂.





Naming Binary Covalent Compounds

A binary covalent compound is typically formed by the combination of two non-metals.

Some of these compounds are very common and have *trivial* names, eg., H₂O is water.

For a binary covalent compound, the element with the *lower* group number in the periodic table is *first* in the name and formula. Its name remains unchanged.

The element that is second is named using the root with the suffix *-ide*. Numerical prefixes indicate the number of atoms of each element present.





Determining Names and Formulas of Binary Covalent Compounds

- **PROBLEM:** (a) What is the formula of carbon disulfide?
 - (b) What is the name of PCl₅?
 - (c) Give the name and formula of the compound whose molecules each consist of two N atoms and four O atoms.

SOLUTION:

- (a) Carbon is C, sulfide is sulfur S and di-means two; the formula is CS₂.
- **(b)** P is phosphorous, Cl is chloride, the prefix for 5 is penta. This is **phosphorous pentachloride**.
- (c) N is nitrogen and is in a lower group number than O (oxygen). The compound formula is N₂O₄ and the name is **dinitrogen** tetraoxide.





Recognizing Incorrect Names and Formulas of Binary Covalent Compounds

PROBLEM:

Explain what is wrong with the name of formula in the second part of each statement and correct it:

- (a) SF₄ is monosulfur pentafluoride.
- (b) Dichlorine heptaoxide is Cl₂O₆.
- (c) N_2O_3 is dinitrotrioxide.

SOLUTION:

- (a) The prefix mono- is not needed if there is only one atom of the first element, and the prefix for four is tetra-. So the name is sulfur tetrafluoride.
- **(b)** Hepta- means 7; the formula should be $\mathbf{Cl_2O_7}$.
- (c) The first element is given its elemental name so this is dinitrogen trioxide.





Naming Straight-Chain Alkanes

Hydrocarbons are compounds that contain only carbon and hydrogen atoms.

Alkanes are the simplest type of hydrocarbon.

Alkanes are named using a root name followed by the suffix *-ane*.





Table 2.7 The First 10 Straight-Chain Alkanes

Name (Formula)	Model
Methane (CH ₄)	
Ethane (C_2H_6)	
Propane (C_3H_8)	
Butane (C_4H_{10})	
Pentane (C_5H_{12})	
Hexane (C_6H_{14})	
Heptane (C_7H_{16})	
Octane (C_8H_{18})	
Nonane (C_9H_{20})	
Decane $(C_{10}H_{22})$	

Molecular Masses from Chemical Formulas

Molecular mass = sum of atomic masses

For the H₂O molecule:

molecular mass =

(2 x atomic mass of H) + (1 x atomic mass of O)

 $= (2 \times 1.008 \text{ amu}) + (1 \times 16.00 \text{ amu})$

= 18.02 amu

By convention, we read masses off the periodic table to 4 significant figures.

For **ionic compounds** we refer to a **formula mass** since ionic compounds do not consist of molecules.



Calculating the Molecular Mass of a Compound

PROBLEM: Using the periodic table, calculate the molecular (or formula) mass of:

(a) tetraphosphorous trisulfide (b) ammonium nitrate

PLAN: Write the formula and then multiply the number of atoms by the respective atomic masses. Add the masses for each compound.

SOLUTION:

```
(a) P_4S_3
molecular mass = (4 \times atomic mass of P) + (3 \times atomic mass of S)
= (4 \times 30.97 \text{ amu}) + (3 \times 32.07 \text{ amu}) = 220.09 \text{ amu}
```

(b) NH_4NO_3

formula mass = (2 x atomic mass of N) + (4 x atomic mass of H) + (3 x atomic mass of O)

 $= (2 \times 14.01 \text{ amu}) + (4 \times 1.008 \text{ amu}) + (3 \times 16.00 \text{ amu})$


= 80.05 amu





Using Molecular Depictions to determine Formula, Name, and Mass for a compound

PROBLEM: Each scene represents a binary compound. Determine its formula, name, and molecular (formula) mass.



PLAN: Each compound contains only two elements. Find the simplest whole number ratio of atoms in each compound and use this formula to determine the name and the formula mass.






UPLOADED BY AHMAD T JUNDI

Sample Problem 2.16

SOLUTION:

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



(a) There is 1 brown Na⁺ for every green F⁻, so the formula is **NaF**, an ionic compound, which is named **sodium fluoride**.

Formula mass = $(1 \times atomic mass of Na) + (1 \times atomic mass of F)$ = 22.99 amu + 10.00 amu = 41.99 amu

(b) There are 3 green F for every blue N, so the formula is NF₃, a covalent compound, which is named **nitrogen trifluoride**.

Molecular mass = (1 x atomic mass of N) + (3 x atomic mass of F)= 14.01 amu + (3 x 19.00) = 71.01 amu





Representing Molecules with Formulas and Models


 H_2O

Molecular formula for water.

H:O:H

H-0-H

Structural formulas for water.



Ball-and-stick model for water.




Space-filling model for water.



Representing Molecules with Formulas and Models

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




e.g., lithium, Li



UPLOADED BY AHMAD T JUNDI

Upload By: Ahmad T Al-Jundi







Chapter 3

Stoichiometry of Formulas and Equations





Mole - Mass Relationships in Chemical Systems

- 3.1 The Mole
- 3.2 Determining the Formula of an Unknown Compound
- 3.3 Writing and Balancing Chemical Equations
- 3.4 Calculating Quantities of Reactant and Product
- 3.5 Fundamentals of Solution Stoichiometry





The Mole

The mole (mol) is the amount of a substance that contains the same number of entities as there are atoms in exactly 12 g of carbon-12.

The term "*entities*" refers to atoms, ions, molecules, formula units, or electrons – in fact, any type of particle.

One mole (1 mol) contains 6.022x10²³ entities (to four significant figures).


This number is called Avogadro's number and is abbreviated as **N**.





One mole (6.022x10²³ entities) of some familiar Figure 3.1 substances.







Determining Molar Mass

The molar mass (\mathcal{M}) of a substance is the mass per mole of its entities (atoms, molecules or formula units).

For **monatomic elements**, the molar mass is the same as the atomic mass in grams per mole. The atomic mass is simply read from the Periodic Table.

The molar mass of Ne = 20.18 g/mol.





For **molecular elements** and for **compounds**, the formula is needed to determine the molar mass.

The molar mass of
$$O_2 = 2 \times \mathcal{M}$$
 of $O_2 = 2 \times 16.00$
= 2×16.00
= 32.00 g/mol

The molar mass of
$$SO_2 = 1 \times M$$
 of $S + 2 \times M$ of $O = 32.00 + 2(16.00)$
= 64.00 g/mol





Table 3.1 Information Contained in the Chemical Formula of Glucose, $C_6H_{12}O_6$ ($\mathcal{M}=180.16$ g/mol)

	Carbon (C)	Hydrogen (H)	Oxygen (O)
Atoms/molecule of compound	6 atoms	12 atoms	6 atoms
Moles of atoms/mole of compound	6 mol of atoms	12 mol of atoms	6 mol of atoms
Atoms/mole of compound	6(6.022x10 ²³) atoms	12(6.022x10 ²³) atoms	6(6.022x10 ²³) atoms
Mass/molecule of compound	6(12.01 amu) = 72.06 amu	12(1.008 amu) = 12.10 amu	6(16.00 amu) = 96.00 amu
Mass/mole of compound	72.06 g	12.10 g	96.00 g





Converting Between Amount, Mass, and Number of Chemical Entities

Mass (g) = no. of moles x
$$\frac{\text{no. of grams}}{1 \text{ mol}} \leftarrow g$$

No. of moles = mass (g) x
$$\frac{1 \text{ mol}}{\text{no. of grams}} \leftarrow \mathcal{M}$$


No. of entities = no. of moles x
$$\frac{6.022 \times 10^{23} \text{ entities}}{1 \text{ mol}}$$

No. of moles = no. of entities x $\frac{1 \text{ mol}}{6.022 \times 10^{23} \text{ entities}}$





Figure 3.2 Mass-mole-number relationships for elements.





Calculating the Mass of a Given Amount of an Element


PROBLEM: Silver (Ag) is used in jewelry and tableware but no

longer in U.S. coins. How many grams of Ag are in

0.0342 mol of Ag?

PLAN: To convert mol of Ag to mass of Ag in g we need the molar

mass of Ag.



SOLUTION:

$$0.0342 \frac{\text{mol Ag}}{\text{1 mol Ag}} \times \frac{107.9 \text{ g Ag}}{1 \frac{\text{mol Ag}}{\text{1 mol Ag}}} = 3.69 \text{ g Ag}$$





Calculating the Number of Entities in a Given Amount of an Element

PROBLEM: Gallium (Ga) is a key element in solar panels, calculators, and other light-sensitive electronic devices. How many Ga atoms are in 2.85 x 10⁻³ mol of gallium?

PLAN: To convert mol of Ga to number of Ga atoms we need to use Avogadro's number.

mol of Ga

multiply by 6.022x10²³ atoms/mol

atoms of Ga





SOLUTION:

2.85 x 10⁻³ mol Ga atoms x 6.022x10²³ Ga atoms 1 mol Ga atoms

= 1.72 x 10²¹ Ga atoms






Calculating the Number of Entities in a Given Mass of an Element

PROBLEM: Iron (Fe) is the main component of steel and is therefore the most important metal in society; it is also essential in the body. How many Fe atoms are in 95.8 g of Fe?

PLAN: The number of atoms cannot be calculated directly from the mass. We must first determine the number of moles of Fe atoms in the sample and then use Avogadro's number.







SOLUTION:

$$95.8 \frac{\text{g Fe}}{\text{55.85 g Fe}} = 1.72 \text{ mol Fe}$$

$$= 1.04 \times 10^{24} \text{ atoms Fe}$$





Figure 3.3 Amount-mass-number relationships for compounds.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display. MASS (g) of compound divide \mathcal{M} (g/mol) chemical formula **AMOUNT** (mol) **AMOUNT** (mol) of elements of compound in compound Avogadro's multiply number (molecules/mol) **MOLECULES** (or formula units) of compound





Calculating the Number of Chemical Entities in a Given Mass of a Compound I

PROBLEM: Nitrogen dioxide is a component of urban smog that forms

from the gases in car exhausts. How many molecules are

in 8.92 g of nitrogen dioxide?

PLAN: Write the formula for the compound and calculate its molar mass. Use the given mass to calculate first the number of moles and then

use the given mass to calculate first the number of moles and the

the number of molecules.

 $mass (g) of NO_2 \\ divide by \mathcal{M} \\ amount (mol) of NO_2 \\ multiply by 6.022 x <math>10^{23}$ formula units/mol $number of NO_2 molecules$





SOLUTION: NO_2 is the formula for nitrogen dioxide.

$$\mathcal{M} = (1 \times \mathcal{M} \text{ of N}) + (2 \times \mathcal{M} \text{ of O})$$

= 14.01 g/mol + 2(16.00 g/mol)
= 46.01 g/mol

$$8.92 \text{ g NO}_2 \text{ x}$$
 $\frac{1 \text{ mol NO}_2}{46.01 \text{ g NO}_2} = 0.194 \text{ mol NO}_2$

=
$$1.17 \times 10^{23}$$
 molecules NO_2





Calculating the Number of Chemical Entities in a Given Mass of a Compound II

PROBLEM:

Ammonium carbonate, a white solid that decomposes on warming, is a component of baking powder.

- a) How many formula units are in 41.6 g of ammonium carbonate?
- b) How many O atoms are in this sample?

PLAN:

Write the formula for the compound and calculate its molar mass. Use the given mass to calculate first the number of moles and then the number of formula units.

The number of O atoms can be determined using the formula and the number of formula units.





mass (g) of $(NH_4)_2CO_3$

divide by \mathcal{M}

amount (mol) of $(NH_4)_2CO_3$

multiply by 6.022 x 10²³ formula units/mol

number of $(NH_4)_2CO_3$ formula units

3 O atoms per formula unit of (NH₄)₂CO₃

number of O atoms

SOLUTION: $(NH_4)_2CO_3$ is the formula for ammonium carbonate.

$$\mathcal{M} = (2 \times M \text{ of N}) + (8 \times \mathcal{M} \text{ of H}) + (1 \times \mathcal{M} \text{ of C}) + (3 \times \mathcal{M} \text{ of O})$$

= $(2 \times 14.01 \text{ g/mol}) + (8 \times 1.008 \text{ g/mol})$
+ $(12.01 \text{ g/mol}) + (3 \times 16.00 \text{ g/mol})$

= 96.09 g/mol



41.6 g (NH₄)₂CO₃ x
$$\frac{1 \text{ mol (NH4)}_2\text{CO}_3}{96.09 \text{ g (NH4)}_2\text{CO}_3} = 0.433 \text{ mol (NH4)}_2\text{CO}_3$$

0.433 mol
$$(NH_4)_2CO_3$$
 x $6.022x10^{23}$ formula units $(NH_4)_2CO_3$ 1 mol $(NH_4)_2CO_3$

= 2.61×10^{23} formula units $(NH_4)_2CO_3$

2.61x10²³ fermula units
$$(NH_4)_2CO_3$$
 x 3 O atoms
1 fermula unit of $(NH_4)_2CO_3$

 $= 7.83 \times 10^{23} \text{ O atoms}$





The Importance of Mass Percent

Mass % of element X =

atoms of X in formula x atomic mass of X (amu)

molecular (or formula) mass of compound (amu)

Mass % of element X =

moles of X in formula x molar mass of X (g/mol) x 100 mass (g) of 1 mol of compound

The individual mass percents added up to 100% (within rounding)





Calculating the Mass Percent of Each Element in a Compound from the Formula

PROBLEM: Glucose $(C_6H_{12}O_6)$ is a key nutrient for generating chemical potential energy in biological systems. What is the mass percent of each element in glucose?


PLAN: Find the molar mass of glucose, which is the mass of 1 mole of glucose. Find the mass of each element in 1 mole of glucose, using the molecular formula.

The mass % for each element is calculated by dividing the mass of that element in 1 mole of glucose by the total mass of 1 mole of glucose, multiplied by 100.





PLAN:







SOLUTION:

In 1 mole of glucose there are 6 moles of C, 12 moles H, and 6 moles O.

6 mol C x
$$\frac{12.01 \text{ g C}}{1 \text{ mol C}}$$
 = 72.06 g C 12 mol H x $\frac{1.008 \text{ g H}}{1 \text{ mol H}}$ = 12.096 g H

6 mol O x
$$\frac{16.00 \text{ g O}}{1 \text{ mol O}} = 96.00 \text{ g O}$$
 $\mathcal{M} = 180.16 \text{ g/mol}$

mass percent of C =
$$\frac{72.06 \text{ g C}}{180.16 \text{ g glucose}} = 0.3999 \text{ x } 100 = 39.99 \text{ mass %C}$$

mass percent of H =
$$\frac{12.096 \text{ g H}}{180.16 \text{ g glucose}}$$
 = 0.06714 x 100 = 6.714 mass %H

mass percent of O =
$$\frac{96.00 \text{ g O}}{180.16 \text{ g glucose}} = 0.5329 \text{ x } 100 = 53.29 \text{ mass } \%\text{O}$$





Determining the Mass of an Element from Its Percent

Mass percent can also be used to calculate the mass of a particular element in any mass of a compound.

Mass of element X present in sample =

mass of compound x

mass of element in 1 mol of compound

mass of 1 mol of compound





Calculating the Mass of an Element in a Compound

PROBLEM: Use the information from Sample Problem 3.6 to determine the mass (g) of carbon in 16.55 g of glucose.

PLAN: The mass percent of carbon in glucose gives us the relative mass of carbon in 1 mole of glucose. We can use this information to find the mass of carbon in any sample of glucose.

mass of glucose sample

multiply by mass percent of C in glucose

mass of C in sample





SOLUTION:

Each mol of glucose contains 6 mol of C, or 72.06 g of C.

Mass (g) of C = mass (g) of glucose x
$$= 6 \text{ mol } x \mathcal{M} \text{ of C (g/mol)}$$

mass (g) of 1 mol of glucose





Empirical and Molecular Formulas

The empirical formula is the simplest formula for a compound that agrees with the elemental analysis. It shows the *lowest* whole number of moles and gives the *relative* number of atoms of each element present.

The empirical formula for hydrogen peroxide is HO.

The molecular formula shows the actual number of atoms of each element in a molecule of the compound.

The molecular formula for hydrogen peroxide is H_2O_2 .





Determining an Empirical Formula from Amounts of Elements

PROBLEM: A sample of an unknown compound contains 0.21

mol of zinc, 0.14 mol of phosphorus, and 0.56 mol of

oxygen. What is its empirical formula?

PLAN: Find the relative number of moles of each element. Divide

by the lowest mol amount to find the relative mol ratios

(empirical formula).

amount (mol) of each element

use # of moles as subscripts

preliminary formula

change to integer subscripts

empirical formula





SOLUTION: Using the numbers of moles of each element given, we write the preliminary formula Zn_{0.21}P_{0.14}O_{0.56}

Next we divide each fraction by the smallest one; in this case 0.14:

$$\frac{0.21}{0.14} = 1.5$$
 $\frac{0.14}{0.14} = 1.0$ $\frac{0.56}{0.14} = 4.0$

$$\frac{0.14}{0.14} = 1.0$$

$$\frac{0.56}{0.14} = 4.0$$

This gives $Zn_{15}P_{10}O_{40}$

We convert to whole numbers by multiplying by the *smallest integer* that gives whole numbers; in this case 2:

$$1.5 \times 2 = 3$$

$$1.0 \times 2 = 2$$

$$1.5 \times 2 = 3$$
 $1.0 \times 2 = 2$ $4.0 \times 2 = 8$

This gives us the empirical formula Zn₃P₂O₈





Determining an Empirical Formula from Masses of Elements

PROBLEM: Analysis of a sample of an ionic compound yields 2.82 g of Na, 4.35 g of Cl, and 7.83 g of O. What is the empirical formula and the name of the compound?

PLAN: Find the relative number of moles of each element. Divide by the lowest mol amount to find the relative mol ratios (empirical formula).

mass (g) of each element

divide by M (g/mol)

amount (mol) of each element

use # of moles as subscripts

preliminary formula

change to integer subscripts

empirical formula





SOLUTION:
$$2.82 \frac{g \text{ Na}}{2} \times \frac{1 \text{ mol Na}}{22.99 \frac{g \text{ Na}}{2}} = 0.123 \text{ mol Na}$$

$$4.35 \, \frac{\text{GCl}}{35.45 \, \frac{\text{Cl}}{\text{GCl}}} = 0.123 \, \text{mol Cl}$$

$$7.83 \frac{\text{g O}}{\text{g O}} \times \frac{1 \text{ mol O}}{16.00 \frac{\text{g O}}{\text{g O}}} = 0.489 \text{ mol O}$$

Na and CI =
$$\frac{0.123}{0.123}$$
 = 1 and O = $\frac{0.489}{0.123}$ = 3.98

The empirical formula is $Na_1CI_1O_{3.98}$ or $NaCIO_4$; this compound is named sodium perchlorate.





Molecular Formulas

The molecular formula gives the *actual* numbers of moles of each element present in 1 mol of compound.

The molecular formula is a *whole-number multiple* of the empirical formula.





Determining a Molecular Formula from Elemental Analysis and Molar Mass

PROBLEM: Elemental analysis of lactic acid (\mathcal{M} = 90.08 g/mol) shows it contains 40.0 mass % C, 6.71 mass % H, and 53.3 mass % O. Determine the empirical formula and the molecular formula for lactic acid.

PLAN:

assume 100 g lactic acid; then mass % = mass in grams

divide each mass by ${\mathcal M}$

amount (mol) of each element

use # mols as subscripts; convert to integers

empirical formula

divide \mathcal{M} by the molar mass for the empirical formula; multiply empirical formula by this number

molecular formula





SOLUTION: Assuming there are 100. g of lactic acid;

$$= 3.33 \text{ mol C}$$

$$= 6.66 \text{ mol H}$$

$$= 3.33 \text{ mol O}$$


$$C_{3.33}$$
 $H_{6.66}$ $O_{3.33}$ 3.33 3.33





Figure 3.4 Combustion apparatus for determining formulas of organic compounds.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



$$C_nH_m + (n + \frac{m}{4}) O_2 = n CO2(g) + \frac{m}{2} H_2O(g)$$





Determining a Molecular Formula from Combustion Analysis

PROBLEM:

When a 1.000 g sample of vitamin C (\mathcal{M} = 176.12 g/mol) is placed in a combustion chamber and burned, the following data are obtained:

mass of CO_2 absorber after combustion = 85.35 g mass of CO_2 absorber before combustion = 83.85 g mass of H_2O absorber after combustion = 37.96 g mass of H_2O absorber before combustion = 37.55 g

What is the molecular formula of vitamin C?

PLAN: The masses of CO₂ and H₂O produced will give us the masses of C and H present in the original sample. From this we can determine the mass of O.





(mass after combustion – mass before) for each absorber = mass of compound in each absorber

mass of each compound x mass % of oxidized element

mass of each oxidized element

mass of vitamin C - (mass of C + H)

mass of O

divide each mass by $\boldsymbol{\mathcal{M}}$

mol of C, H, and O

use # mols as subscripts; convert to integers

empirical formula

molecular formula





SOLUTION: For CO_2 : 85.35 g - 83.85 g = 1.50 g

$$1.50 \text{ g CO}_2 \times \frac{12.01 \text{ g C}}{44.01 \text{ g CO}_2} = 0.409 \text{ g C}$$

For H_2O : 37.96 g - 37.55 g = 0.41 g

$$0.41 \text{ g H}_2\text{O x} = \frac{2.016 \text{ g H}}{18.02 \text{ g H}_2\text{O}} = 0.046 \text{ g H}$$

mass of O = mass of vitamin C – (mass of C + mass of H)
=
$$1.000 \text{ g} - (0.409 + 0.046) \text{ g} = 0.545 \text{ g} \text{ O}$$





Convert mass to moles:

$$\frac{0.409 \text{ g C}}{12.01 \text{ g/mol C}} = 0.0341 \text{ mol C} \qquad \frac{0.046 \text{ g H}}{1.008 \text{ g/mol H}} = 0.0456 \text{ mol H}$$

$$\frac{0.545 \text{ g O}}{16.00 \text{ g/mol O}} = 0.0341 \text{ mol O}$$

Divide by smallest to get the preliminary formula:

C
$$\frac{0.0341}{0.0341} = 1$$
 H $\frac{0.0456}{0.0341} = 1.34$ O $\frac{0.0341}{0.0341} = 1$ C₁H_{1.34}O₁ = C₃H_{4.01}O₃ \longrightarrow C₃H₄O₃

Divide molar mass by mass of empirical formula:

$$\frac{176.12 \text{ g/mol}}{88.06 \text{ g}} = 2.000 \text{ mol} \longrightarrow \boxed{\mathbf{C_6 H_8 O_6}}$$





Table 3.2 Two Constitutional Isomers of C₂H₆O

	C_2H_6O
Property	Ethanol Dimethyl Ether
м (g/mol)	46.07 46.07
Boiling Point	78.5°C -25°C
Density at 20°C	0.789 g/mL
Structural formula	H H H H H H H H H H H H H H H H H H H
Space-filling model	



Writing and Balancing Chemical Equations

A *chemical equation* uses formulas to express the *identities* and *quantities* of substances involved in a physical or chemical change.




Figure 3.5


The formation of HF gas on the macroscopic and molecular levels.





Figure 3.6 A three-level view of the reaction between magnesium and oxygen.


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.







Features of Chemical Equations



The equation must be **balanced**; the same number and type of each atom must appear on both sides.





Balancing a Chemical Equation

translate the statement

magnesium and oxygen gas react to give magnesium oxide:

$$Mg + O_2 \rightarrow MgO$$

balance the atoms using *coefficients*; formulas cannot be changed

$$2Mg + O_2 \rightarrow 2MgO$$

adjust coefficients if necessary

check that all atoms balance

specify states of matter

2Mg
$$(s) + O_2(g) \rightarrow 2$$
MgO (s)





Balancing Chemical Equations

PROBLEM: Within the cylinders of a car's engine, the hydrocarbon octane (C₈H₁₈), one of many components of gasoline, mixes with oxygen from the air and burns to form carbon dioxide and water vapor. Write a balanced equation for this reaction.

PLAN:

translate the statement

balance the atoms

adjust the coefficients

check the atoms balance

specify states of matter

SOLUTION:

$$C_8H_{18} + O_2 \longrightarrow CO_2 + H_2O$$


$$C_8H_{18} + \frac{25}{2}O_2 \longrightarrow 8CO_2 + 9H_2O$$

$$2C_8H_{18} + 25O_2 \longrightarrow 16CO_2 + 18H_2O$$

$$2C_8H_{18} + 25O_2 \longrightarrow 16CO_2 + 18H_2O$$

$$2C_8H_{18}(I) + 25O_2(g) \longrightarrow 16CO_2(g) + 18H_2O(g)$$

Molecular Scene Combustion of Octane








Balancing an Equation from a Molecular Scene

PROBLEM: The following molecular scenes depict an important reaction in nitrogen chemistry. The blue spheres represent nitrogen while the red spheres represent oxygen. Write a balanced equation for this reaction.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



PLAN: Determine the formulas of the reactants and products from their composition. Arrange this information in the correct equation format and balance correctly, including the states of matter.





SOLUTION:

The reactant circle shows only one type of molecule, composed of 2 N and 5 O atoms. The formula is thus N_2O_5 . There are 4 N_2O_5 molecules depicted.

The product circle shows two types of molecule; one has 1 N and 2 O atoms while the other has 2 O atoms. The products are NO_2 and O_2 . There are 8 NO_2 molecules and 2 O_2 molecules shown.

The reaction depicted is $4 N_2 O_5 \rightarrow 8 NO_2 + 2 O_2$.

Writing the equation with the smallest whole-number coefficients and states of matter included;

$$2 N_2 O_5 (g) \rightarrow 4 NO_2 (g) + O_2 (g)$$





Stoichiometric Calculations

- The coefficients in a balanced chemical equation
 - represent the relative number of reactant and product particles
 - and the relative number of moles of each.
- Since moles are related to mass
 - the equation can be used to calculate masses of reactants and/or products for a given reaction.
- The mole ratios from the balanced equation are used as conversion factors.

$$2 N_2 O_5 (g) \rightarrow 4 NO_2 (g) + O_2 (g)$$





UPLOADED BY AHMAD T JUNDI

Table 3.3 Information Contained in a Balanced Equation

Viewed in Terms of


Reactants

 $C_3H_8(g) + 5O_2(g)$

Products

 $3 CO_2(g) + 4 H_2O(g)$

Molecules 1 molecule $C_3H_8 + 5$ molecules $O_2 \rightarrow 3$ molecules $CO_2 + 4$ molecules H_2O



Amount (mol) 1 mol C_3H_8 + 5 mol O_2

 $3 \text{ mol CO}_2 + 4 \text{ mol H}_2\text{O}$

Mass (amu) 44.09 amu C_3H_8 + 160.00 amu $O_2 \rightarrow$ 132.03 amu CO_2 + 72.06 amu H_2O

Mass (g)

 $44.09 \text{ g C}_3H_8 + 160.00 \text{ g O}_2 \longrightarrow 132.03 \text{ g CO}_2 + 72.06 \text{ g H}_2O$

Total Mass (g)

204.09 g

204.09 g





Figure 3.7 Summary of amount-mass-number relationships in a chemical equation.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display. MASS (g) MASS (g) of substance B of substance A \mathcal{M} (g/mol) of \mathcal{M} (g/mol) of substance A substance B molar ratio from balanced equation AMOUNT (mol) AMOUNT (mol) of substance A of substance B Avogadro's Avogadro's number number (entities/mol) (entities/mol) **ENTITIES ENTITIES** of substance A of substance B



Calculating Quantities of Reactants and Products: Amount (mol) to Amount (mol)

PROBLEM:

Copper is obtained from copper(I) sulfide by roasting it in the presence of oxygen gas to form powdered copper(I) oxide and gaseous sulfur dioxide.

How many moles of oxygen are required to roast 10.0 mol of copper(I) sulfide?

PLAN:

write and balance the equation

use the mole ratio as a conversion factor

moles of oxygen

$$2 \text{ Cu}_2\text{S}(s) + 3 \text{ O}_2(g) \rightarrow 2 \text{ Cu}_2\text{O}(s) + 2 \text{ SO}_2(g)$$

10.0 mol
$$Cu_2S \times \frac{3 \text{ mol } O_2}{2 \text{ mol } Cu_2S}$$
 = 15.0 mol O_2





Calculating Quantities of Reactants and Products: Amount (mol) to Mass (g)

PROBLEM: During th

During the process of roasting copper(I) sulfide, how many grams of sulfur dioxide form when 10.0 mol of copper(I) sulfide reacts?

PLAN:

Using the balanced equation from the previous problem, we again use the mole ratio as a conversion factor.

use the mole ratio as a conversion factor
mol of sulfur dioxide
multiply by M of sulfur dioxide
mass of sulfur dioxide





SOLUTION:
$$2 \text{ Cu}_2 \text{S}(s) + 3 \text{ O}_2(g) \rightarrow 2 \text{ Cu}_2 \text{O}(s) + 2 \text{ SO}_2(g)$$

10.0 mol
$$Cu_2S \times 2 \text{ mol } SO_2 \times 64.07 \text{ g } SO_2 \times 2 \text{ mol } Cu_2S \times 1 \text{ mol } SO_2 \times 1 \text{ mol$$






Calculating Quantities of Reactants and Products: Mass to Mass

PROBLEM: During the roasting of copper(I) sulfide, how many

kilograms of oxygen are required to form 2.86 kg of

copper(I) oxide?

PLAN:







SOLUTION:
$$2 \text{ Cu}_2 \text{S}(s) + 3 \text{ O}_2(g) \rightarrow 2 \text{ Cu}_2 \text{O}(s) + 2 \text{ SO}_2(g)$$

2.86 kg
$$\frac{\text{Cu}_2\text{O} \times \text{I}}{1 \text{ kg}}$$
 $\frac{10^3 \text{ g}}{143.10 \text{ g Cu}_2\text{O}}$ = 20.0 mol $\frac{\text{Cu}_2\text{O}}{143.10 \text{ g Cu}_2\text{O}}$

20.0 mol Cu₂O x
$$\frac{3 \text{ mol O}_2}{2 \text{ mol Cu}_2\text{O}}$$
 x $\frac{32.00 \text{ g} \cdot \text{O}_2}{1 \text{ mol O}_2}$ x $\frac{1 \text{ kg}}{10^3 \text{ g}}$ = 0.959 kg O₂





Reactions in Sequence

- Reactions often occur in sequence.
- The product of one reaction becomes a reactant in the next.
- An overall reaction is written by combining the reactions;
 - any substance that forms in one reaction and reacts in the next can be eliminated.

$$\begin{array}{ccc} \mathsf{A} & \to & \mathsf{C} \\ \underline{\mathsf{C}} & \to & \mathsf{B} \\ \mathsf{A} & & & \mathsf{C} \end{array}$$





Limiting Reactants


- So far we have assumed that reactants are present in the correct amounts to react completely.
- In reality, one reactant may *limit* the amount of product that can form.
- The *limiting* reactant will be completely used up in the reaction.
- The reactant that is not limiting is in excess some of this reactant will be left over.





Using Molecular Depictions in a Limiting-Reactant Problem

PROBLEM: Chlorine trifluoride, an extremely reactive substance, is formed as a gas by the reaction of elemental chlorine and fluorine. The molecular scene shows a representative portion of the reaction mixture before the reaction starts. (Chlorine is green, and fluorine is yellow.)




- (a) Find the limiting reactant.
- (b) Write a reaction table for the process.
- (c) Draw a representative portion of the mixture after the reaction is complete. (Hint: The CIF₃ molecule has 1 Cl atom bonded to 3 individual F atoms).





SOLUTION: The balanced equation is $Cl_2(g) + 3F_2(g) \rightarrow 2ClF_3(g)$



There are 3 molecules of Cl₂ and 6 molecules of F₂ depicted:

3 molecules
$$Cl_2 \times \frac{2 \text{ molecules } ClF_3}{1 \text{ molecule } Cl_2} = 6 \text{ molecules } ClF_3$$

6 molecules
$$F_2 \times \frac{2 \text{ molecules CIF}_3}{3 \text{ molecule Cl}_2} = 4 \text{ molecules CIF}_3$$

Since the given amount of F_2 can form less product, it is the limiting reactant.






We use the amount of F_2 to determine the "change" in the reaction table, since F_2 is the limiting reactant:

Molecules	Cl ₂ (g) +	$3F_2(g) \rightarrow$	2CIF ₃ (<i>g</i>)
Initial	3	6	0
Change	-2	-6	+4
Final	1	0	4

The final reaction scene shows that all the F₂ has reacted and that there is Cl₂ left over. 4 molecules of CIF₂ have formed:







Calculating Quantities in a Limiting-Reactant Problem: Amount to Amount

PROBLEM: In another preparation of CIF_3 , 0.750 mol of CI_2 reacts with 3.00 mol of F_2 .

- (a) Find the limiting reactant.
- (b) Write a reaction table.

PLAN: Find the limiting reactant by calculating the amount (mol) of CIF₃ that can be formed from each given amount of reactant. Use this information to construct a reaction table.

SOLUTION: The balanced equation is $Cl_2(g) + 3F_2(g) \rightarrow 2ClF_3(g)$

$$0.750 \frac{\text{mol Cl}_2 \text{ x}}{1 \frac{\text{mol Cl}_2}{\text{mol Cl}_2}} = 1.50 \text{ mol ClF}_3$$

$$3.00 \text{ mol } F_2 \times \frac{2 \text{ mol } ClF_3}{3 \text{ mol } F_2} = 2.00 \text{ mol } ClF_3$$

Cl₂ is limiting, because it yields less CIF₃.





All the Cl_2 reacts since this is the limiting reactant. For every 1 Cl_2 that reacts, 3 F_2 will react, so 3(0.750) or 2.25 moles of F_2 reacts.

Moles	$Cl_2(g)$ +	$3F_2(g) \rightarrow$	2CIF ₃ (<i>g</i>)
Initial	0.750	3.00	0
Change	-0.750	- 2.25	+1.50
Final	0	0.75	1.50





Calculating Quantities in a Limiting-Reactant Problem: Mass to Mass


PROBLEM: A fuel mixture used in the early days of rocketry consisted of two liquids, hydrazine (N_2H_4) and dinitrogen tetraoxide (N_2O_4) , which ignite on contact to form nitrogen gas and water vapor.

- (a) How many grams of nitrogen gas form when 1.00×10^2 g of N_2H_4 and 2.00×10^2 g of N_2O_4 are mixed?
- **(b)** Write a reaction table for this process.

PLAN: Find the limiting reactant by calculating the amount (mol) of CIF₃ that can be formed from each given mass of reactant. Use this information to construct a reaction table.











SOLUTION:
$$2N_2H_4(I) + N_2O_4(I) \rightarrow 3N_2(g) + 4H_2O(g)$$

For
$$N_2H_4$$
: 1.00x 10² g N_2H_4 x $\frac{1 \text{ mol } N_2H_4}{32.05 \text{ g } N_2H_4}$ = 3.12 mol N_2H_4

$$3.12 \text{ mol N}_2H_4 \times \frac{3 \text{ mol N}_2}{2 \text{ mol N}_2H_4} = 4.68 \text{ mol N}_2$$

For
$$N_2O_4$$
: 2.00x 10^2 g N_2O_4 x $\frac{1 \text{ mol } N_2O_4}{92.02 \text{ g } N_2O_4} = 2.17 \text{ mol } N_2$

$$2.17 \text{ mol } N_2 O_4 \times \frac{3 \text{ mol } N_2}{1 \text{ mol } N_2 O_4} = 6.51 \text{ mol } N_2$$

 N_2H_4 is limiting and only 4.68 mol of N_2 can be produced:

$$4.68 \text{ mol N}_2 \times \frac{28.02 \text{ g N}_2}{1 \text{ mol N}_2} = 131 \text{ g N}_2$$





All the N_2H_4 reacts since it is the limiting reactant. For every 2 moles of N_2H_4 that react 1 mol of N_2O_4 reacts and 3 mol of N_2 form:

$$3.12 \text{ mol N}_2H_4 \times \frac{1 \text{ mol N}_2O_4}{2 \text{ mol N}_2H_4} = 1.56 \text{ mol N}_2O_4 \text{ reacts}$$

Moles	2N ₂ H ₄ (I) +	N_2O_4 (1) \rightarrow	$3N_2(g) +$	4H ₂ O (<i>g</i>)
Initial	3.12	2.17	0	0
Change	-3.12	- 1.56	+4.68	+6.24
Final	0	0.61	4.68	6.24





Reaction Yields

The **theoretical yield** is the amount of product calculated using the molar ratios from the balanced equation.

The **actual yield** is the amount of product actually obtained.

The actual yield is usually less than the theoretical yield.

Can expressed in moles or grams





Figure 3.8

The effect of side reactions on the yield of the main product.


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display. A + B(reactants) (side product) (main product)



Calculating Percent Yield

PROBLEM: Silicon carbide (SiC) is made by reacting sand(silicon dioxide, SiO₂) with powdered carbon at high temperature. Carbon monoxide is also formed. What is the percent yield if 51.4 kg of SiC is recovered from processing 100.0 kg of sand?

PLAN:







SOLUTION:
$$SiO_2(s) + 3C(s) \rightarrow SiC(s) + 2CO(g)$$

What is the percent yield if 51.4 kg of SiC is recovered from processing 100.0 kg of sand?

$$100.0 \text{ kg SiO}_2 \times \frac{10^3 \text{ g}}{1 \text{ kg}} \times \frac{1 \text{ mol SiO}_2}{60.09 \text{ g SiO}_2} = 1664 \text{ mol SiO}_2$$

mol SiO₂ = mol SiC = 1664 mol SiC

1664 mol SiC x
$$\frac{40.10 \text{ g SiC}}{1 \text{ mol SiC}}$$
 x $\frac{1 \text{ kg}}{10^3 \text{g}}$ = 66.73 kg $\frac{51.4 \text{ kg}}{66.73 \text{ kg}}$ x 100 $\boxed{= 77.0\%}$





Solution Stoichiometry

- Many reactions occur in solution.
- A solution consists of one or more solutes dissolved in a solvent.
- The concentration of a solution is given by the quantity of solute present in a given quantity of solution.
- *Molarity* (*M*) is often used to express concentration.

$$M = \frac{\text{mol solute}}{\text{L soln}}$$





Calculating the Molarity of a Solution

PROBLEM: What is the molarity of an aqueous solution that contains 0.715 mol of glycine (H₂NCH₂COOH) in 495 mL?

PLAN: SOLUTION:

Molarity is the number of moles of solute per liter of solution.








Figure 3.9

Summary of mass-mole-number-volume relationships in solution.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display. MASS (g) of substance in solution \mathcal{M} (g/mol) Avogadro's number M (mol/L)(entities/mol) AMOUNT (mol) **ENTITIES VOLUME (L)** of substance of substance of solution in solution in solution








Calculating Mass of Solute in a Given Volume of Solution

PROBLEM: How many grams of solute are in 1.75 L of 0.460 *M* sodium monohydrogen phosphate buffer solution?

PLAN: Calculate the moles of solute using the given molarity and volume.

Convert moles to mass using the molar mass of the solute.



SOLUTION:

$$1.75 \pm x \quad 0.460 \text{ moles} = 0.805 \text{ mol Na}_2 \text{HPO}_4$$


$$0.805 \text{ mol Na}_2\text{HPO}_4 \times \frac{141.96 \text{ g Na}_2\text{HPO}_4}{1 \text{ mol Na}_2\text{HPO}_4}$$





Figure 3.10 Converting a concentrated solution to a dilute solution.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



© The McGraw-Hill Companies, Inc./Stephen Frisch Photographer.



Preparing a Dilute Solution from a Concentrated Solution

PROBLEM: "Isotonic saline" is a 0.15 *M* aqueous solution of NaCl. How would you prepare 0.80 L of isotonic saline from a 6.0 *M* stock solution?

PLAN: To dilute a concentrated solution, we add only solvent, so the *moles of solute are the same in both solutions*. The volume and molarity of the dilute solution gives us the moles of solute. Then we calculate the volume of concentrated solution that contains the same number of moles.

volume of dilute soln

multiply by M of dilute soln

moles of NaCl in dilute soln = mol NaCl in concentrated soln

divide by M of concentrated soln

L of concentrated soln





$$M_{\text{dil}} \times V_{\text{dil}} = \# \text{ mol solute} = M_{\text{conc}} \times V_{\text{conc}}$$

SOLUTION:

Using the volume and molarity for the dilute solution:

$$0.80 \text{ L-soln} \times \frac{0.15 \text{ mol NaCl}}{1 \text{ L-soln}} = 0.12 \text{ mol NaCl}$$

Using the moles of solute and molarity for the concentrated solution:

A 0.020 L portion of the concentrated solution must be diluted to a final volume of 0.80 L.






Visualizing Changes in Concentration

PROBLEM: The beaker and circle represent a unit volume of solution. Draw the solution after each of these changes:

- (a) For every 1 mL of solution, 1 mL of solvent is added.
- **(b)** One third of the volume of the solution is boiled off.



PLAN: Only the volume of the solution changes; the total number of moles of solute remains the same. Find the new volume and calculate the number of moles of solute per unit volume.







SOLUTION: $N_{dil} \times V_{dil} = N_{conc} \times V_{conc}$ where N is the number of particles.

(a)
$$N_{dil} = N_{conc} \times \frac{V_{conc}}{V_{dil}} = 8 \text{ particles } \times \frac{1 \text{ mL}}{2 \text{ mL}} = 4 \text{ particles}$$

(b)
$$N_{\text{conc}} = N_{\text{dil}} \times \frac{V_{\text{dil}}}{V_{\text{conc}}} = 8 \text{ particles } \times \frac{1 \text{ mL}}{\frac{2}{3} \text{ mL}} = 12 \text{ particles}$$










Calculating Quantities of Reactants and Products for a Reaction in Solution

PROBLEM: A 0.10 *M* HCl solution is used to simulate the acid concentration of the stomach. How many liters of "stomach acid" react with a tablet containing 0.10 g of magnesium hydroxide?

PLAN: Write a balanced equation and convert the mass of Mg(OH)₂ to moles. Use the mole ratio to determine the moles of HCl, then convert to volume using molarity.







SOLUTION:

$$Mg(OH)_2(s) + 2HCI(aq) \rightarrow MgCI_2(aq) + 2H_2O(l)$$

$$0.10 \text{ g Mg(OH)}_2 \times \frac{1 \text{ mol Mg(OH)}_2}{58.33 \text{ g Mg(OH)}_2} = 1.7 \times 10^{-3} \text{ mol Mg(OH)}_2$$

=
$$1.7x10^{-3} \text{ mol Mg(OH)}_2 \times \frac{2 \text{ mol HCI}}{1 \text{ mol Mg(OH)}_2} = 3.4x10^{-3} \text{ mol HCI}$$






Solving Limiting-Reactant Problems for Reactions in Solution

PROBLEM: In a simulation mercury removal from industrial wastewater, 0.050 L of 0.010 *M* mercury(II) nitrate reacts with 0.020 L of 0.10 *M* sodium sulfide. How many grams of mercury(II) sulfide form? Write a reaction table for this process.

PLAN: Write a balanced chemical reaction. Determine limiting reactant. Calculate the grams of mercury(II) sulfide product.





SOLUTION: $\operatorname{Hg}(NO_3)_2(aq) + \operatorname{Na}_2S(aq) \rightarrow \operatorname{HgS}(s) + 2\operatorname{NaNO}_3(aq)$

$$0.050 \text{ LHg(NO}_3)_2 \times \frac{0.010 \text{ mol Hg(NO}_3)_2}{1 \text{ LHg(NO}_3)_2} \times \frac{1 \text{ mol HgS}}{1 \text{ mol Hg(NO}_3)_2}$$

= $5.0 \times 10^{-4} \text{ mol HgS}$

$$0.020 \text{ L Na}_2\text{S} \times \frac{0.10 \text{ mol Na}_2\text{S}}{1 \text{ L Na}_2\text{S}} \times \frac{1 \text{ mol HgS}}{1 \text{ mol Na}_2\text{S}} = 2.0 \times 10^{-3} \text{ mol HgS}$$

 $Hg(NO_3)_2$ is the limiting reactant because it yields less HgS.






The reaction table is constructed using the amount of $Hg(NO_3)_2$ to determine the changes, since it is the limiting reactant.

Amount	$Hg(NO_3)_2 (aq) +$	$Na_2S(aq) \rightarrow$	HgS (s) +	2NaNO ₃ (<i>aq</i>)
Initial	5.0 x 10 ⁻⁴	2.0 x 10 ⁻³	0	0
Change	-5.0 x 10 ⁻⁴	−5.0 x 10 ⁻⁴	+5.0 x 10 ⁻⁴	+1.0 x 10 ⁻³
Final	0	1.5 x 10 ⁻³	5.0 x 10 ⁻⁴	+1.0 x 10 ⁻³







Chapter 05

Lecture Outline

See separate *Image PowerPoint* slides for all figures and tables pre-inserted into PowerPoint without notes.





Chapter 5

Gases and the Kinetic-Molecular Theory





Gases and the Kinetic Molecular Theory

- 5.1 An Overview of the Physical States of Matter
- **5.2 Gas Pressure and Its Measurement**
- 5.3 The Gas Laws and Their Experimental Foundations
- 5.4 Rearrangements of the Ideal Gas Law
- 5.5 The Kinetic-Molecular Theory: A Model for Gas Behavior
- 5.6 Real Gases: Deviations from Ideal Behavior





An Overview of the Physical States of Matter

Distinguishing gases from liquids and solids.

- Gas volume changes significantly with pressure.
 - Solid and liquid volumes are not greatly affected by pressure.
- Gas volume changes significantly with temperature.
 - Gases expand when heated and shrink when cooled.
 - The volume change is 50 to 100 times greater for gases than for liquids and solids.
- Gases flow very freely.
- Gases have relatively low densities.
- Gases form a solution in any proportions.
 - Gases are freely miscible with each other.






Figure 5.1 The three states of matter.


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.






Gas: Particles are far apart, move freely, and fill the available space.





Liquid: Particles are close together but move around one another.





Solid: Particles are close together in a regular array and do not move around one another.





Gas Pressure and its Measurement

Pressure =
$$\frac{\text{force}}{\text{area}}$$

Atmospheric pressure arises from the force exerted by atmospheric gases on the earth's surface.


Atmospheric pressure decreases with altitude.






Figure 5.2 Effect of atmospheric pressure on a familiar object.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.





© The McGraw-Hill Companies, Inc./Stephen Frisch Photographer





Figure 5.3 A mercury barometer.

$$\frac{h_{\text{H2O}}}{h_{\text{Hg}}} = \frac{d_{\text{Hg}}}{d_{\text{H2O}}}$$

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




Table 5.1 Common Units of Pressure

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Unit	Normal Atmospheric Pressure at Sea Level and 0°C		
pascal (Pa); kilopascal (kPa)	$1.01325 \times 10^5 \text{ Pa}$; 101.325 kPa		
atmosphere (atm)	1 atm*		
millimeters of mercury (mmHg)	760 mmHg*		
torr	760 torr*		
pounds per square inch (lb/in² or psi)	14.7 lb/in ²		
bar	1.01325 bar		

^{*}This is an exact quantity; in calculations, we use as many significant figures as necessary.





Converting Units of Pressure

PROBLEM: A geochemist heats a limestone (CaCO₃) sample and collects the CO₂ released in an evacuated flask attached to a closed-end manometer. After the system comes to room temperature, $\Delta h = 291.4$ mm Hg. Calculate the CO₂ pressure in torrs, atmospheres, and kilopascals.

PLAN: Construct conversion factors to find the other units of pressure.

SOLUTION: 291.4 mmHg x
$$\frac{1 \text{torr}}{1 \text{ mmHg}}$$
 = 291.4 torr

291.4
$$\frac{1 \text{ atm}}{760 \text{ torr}}$$
 = **0.3834 atm**






The Gas Laws

- The gas laws describe the physical behavior of gases in terms of 4 variables:
 - pressure (P)
 - temperature (7)
 - volume (
 - amount (number of moles, n)
- An ideal gas is a gas that exhibits linear relationships among these variables.
- No ideal gas actually exists, but most simple gases behave nearly ideally at ordinary temperatures and pressures.






Figure 5.4 Boyle's law, the relationship between the volume and pressure of a gas.



V (mL)	P (torr)		11	PV	
	Δh ·	+ Patm	= P _{total}	P _{total}	(torr•mL)
20.0	20.0	760	780	0.00128	1.56×10 ⁴
15.0	278	760	1038	0.000963	1.56×10^4
10.0	800	760	1560	0.000641	1.56×10^4
5.0	2352	760	3112	0.000321	1.56×10^4





Boyle's Law

At constant temperature, the volume occupied by a fixed amount of gas is *inversely* proportional to the external pressure.

$$V \propto \frac{1}{P}$$
 or $PV = \text{constant}$

At fixed T and n,
P decreases as V increases
P increases as V decreases





Figure 5.5 AB Charles's law, the relationship between the volume and temperature of a gas.

Copyright @ The McGraw-Hill Companies, Inc. Permission required for reproduction or display.









Figure 5.5 C Absolute zero (0 K) is the temperature at which an ideal gas would have a zero volume.







Charles's Law

At constant pressure, the volume occupied by a fixed amount of gas is *directly* proportional to its absolute (Kelvin) temperature.

$$V \propto T$$
 $\frac{V}{T} = \text{constant}$


At fixed T and n,
P decreases as V increases
P increases as V decreases





Figure 5.6 The relationship between the volume and amount of a gas.

At fixed temperature and pressure, the volume occupied by a gas is directly proportional to the amount of gas.



Avogadro's Law: at fixed temperature and pressure, equal volumes of *any* ideal gas contain equal numbers of particles (or moles).





Gas Behavior at Standard Conditions

STP or standard temperature and pressure specifies a pressure of 1 atm (760 torr) and a temperature of 0°C (273.15 K).


The **standard molar volume** is the volume of 1 mol of an ideal gas at STP.

Standard molar volume = 22.4141 L or **22.4** L





Figure 5.8 Standard molar volume.



<i>n</i> = 1 mol	n = 1 mol	n = 1 mol	
P = 1 atm (760 torr)	P = 1 atm (760 torr)	P = 1 atm (760 torr)	
T = 0°C (273 K)	T = 0°C (273 K)	T = 0°C (273 K)	
V = 22.4 L	V = 22.4 L	V = 22.4 L	
Number of gas particles = 6.022×10 ²³	Number of gas particles = 6.022×10 ²³	Number of gas particles = 6.022×10 ²³	
Mass = 4.003 g	Mass = 28.02 g	Mass = 32.00 g	
d = 0.179 g/L	d = 1.25 g/L	d = 1.43 g/L	



Figure 5.9

The volume of 1 mol (22.4 L) of an ideal gas and of some familiar objects: 1 gal of milk (3.79 L), a basketball (7.50 L) and 2.00 L of a carbonated drink.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.







The Ideal Gas Law

$$pV = nRT$$

$$R = \frac{PV}{nT} = \frac{1 \text{ atm x } 22.414 \text{ L}}{1 \text{ mol x } 273.15 \text{ K}} = \frac{0.0821 \text{ atm \cdot L}}{\text{mol \cdot K}}$$

R is the universal gas constant; the numerical value of *R* depends on the units used.

The ideal gas law can also be expressed by the combined equation:

$$\frac{P_1 V_1}{T_1 n_1} = \frac{P_2 V_2}{T_2 n_2}$$






Figure 5.10 The individual gas laws as special cases of the ideal gas law.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

IDEAL GAS LAW








Applying the Volume-Pressure Relationship

PROBLEM: Boyle's apprentice finds that the air trapped in a J tube occupies 24.8 cm³ at 1.12 atm. By adding mercury to the tube, he increases the pressure on the trapped air to 2.64 atm. Assuming constant temperature, what is the new volume of air (in L)?

PLAN: The temperature and amount of gas are fixed, so this problem involves a change in pressure and volume only.







SOLUTION:

$$P_1 = 1.12 \text{ atm}$$
 $P_2 = 2.64 \text{ atm}$ $V_1 = 24.8 \text{ cm}^3$ $V_2 = \text{unknown}$

n and Tare constant

$$24.8 \text{ cm}^3 \text{ x } \frac{1 \text{ mL}}{1 \text{ cm}^3} \text{ x } \frac{L}{10^3 \text{ mL}} = 0.0248 \text{ L}$$

$$\frac{P_1 V_1}{P_1 T_1} = \frac{P_2 V_2}{P_2 T_2}$$

$$P_1 V_1 = P_2 V_2$$

$$V_2 = V_1 \times \frac{P_1}{P_2} = 0.0248 \text{ L} \times \frac{1.12 \text{ atm}}{2.46 \text{ atm}} = 0.0105 \text{ L}$$






Applying the Pressure-Temperature Relationship

PROBLEM: A steel tank used for fuel delivery is fitted with a safety valve that opens when the internal pressure exceeds 1.00×10^3 torr. It is filled with methane at 23°C and 0.991 atm and placed in boiling water at exactly 100°C. Will the safety valve open?

PLAN:

We must determine if the pressure will exceed 1.00x10³ torr at the new temperature. Since the gas is in a steel tank, the volume remains constant.



UPLOADED BY AHMAD T JUNDI

Sample Problem 5.3

SOLUTION:

$$P_1 = 0.991 \text{ atm}$$
 $P_2 = \text{unknown}$
 $T_1 = 23^{\circ}\text{C}$ $T_2 = 100.^{\circ}\text{C}$

n and *V* are constant

$$0.991 \frac{\text{atm x}}{1 \frac{\text{atm}}{1}} = 753 \text{ torr}$$

$$T_1 = 23 + 273.15 = 296 \text{ K}$$

 $T_2 = 100. + 273.15 = 373 \text{ K}$

$$\frac{P_1 V_1}{P_1 T_1} = \frac{P_2 V_2}{P_2 T_2}$$

$$\frac{P_1}{T_1} = \frac{P_2}{T_2}$$

$$P_2 = P_1 \times \frac{T_2}{T_1} = 753 \text{ torr } \times \frac{373 \text{ K}}{296 \text{ K}} = 949 \text{ torr}$$




The safety valve will not open, since P_2 is less than 1.00 x 10³ torr.



Applying the Volume-Amount Relationship

PROBLEM: A scale model of a blimp rises when it is filled with helium to a volume of 55.0 dm³. When 1.10 mol of He is added to the blimp, the volume is 26.2 dm³. How many more grams of He must be added to make it rise? Assume constant *T* and *P*.

PLAN: The initial amount of helium (n_1) is given, as well as the initial volume (V_1) and the volume needed to make it rise (V_2) . We need to calculate n_2 and hence the mass of He to be added.







SOLUTION:

$$n_1 = 1.10 \text{ mol}$$
 $n_2 = \text{unknown}$
 $V_1 = 26.2 \text{ dm}^3$ $V_2 = 55.0 \text{ dm}^3$

Tand Pare constant

$$\frac{P_1 V_1}{n_1 T_1} = \frac{P_2 V_2}{n_2 T_2}$$

$$\frac{V_1}{n_1} = \frac{V_2}{n_2}$$

$$n_2 = n_1 \times \frac{V_2}{V_1} = 1.10 \text{ mol } \times \frac{55.0 \text{ dm}^3}{26.2 \text{ dm}^3} = 2.31 \text{ mol He}$$

Additional amount of He needed = 2.31 mol – 1.10 mol = 1.21 mol He

1.21 mol He x
$$\frac{4.003 \text{ g He}}{1 \text{ mol He}}$$
 = 4.84 g He





Solving for an Unknown Gas Variable at Fixed Conditions

PROBLEM: A steel tank has a volume of 438 L and is filled with 0.885

kg of O₂. Calculate the pressure of O₂ at 21°C.

PLAN: We are given V, T and mass, which can be converted to

moles (n). Use the ideal gas law to find P.

SOLUTION:
$$V = 438 L$$
 $T = 21^{\circ}C = 294 K$

 $n = 0.885 \text{ kg O}_2$ (convert to mol) P is unknown

$$0.885 \frac{\text{kg } O_2}{4} \times \frac{10^3 \text{ g}}{4} \times \frac{1 \text{ mol } O_2}{32.00 \text{ g } O_2} = 27.7 \text{ mol } O_2$$


$$P = \frac{nRT}{V} = \frac{27.7 \text{ mol } \times 0.0821 \frac{\text{atm} \cdot \text{L}}{\text{mol} \cdot \text{K}} \times 294.15 \text{ K}}{438 \text{ L}} = 1.53 \text{ atm}$$





Using Gas Laws to Determine a Balanced **Equation**

The piston-cylinder is depicted before and after a gaseous PROBLEM: reaction that is carried out at constant pressure. The temperature is 150 K before the reaction and 300 K after



Which of the following balanced equations describes the reaction?

$$(1) A2(g) + B2(g) \rightarrow 2AB(g)$$

(2)
$$2AB(g) + B_2(g) \rightarrow 2AB_2(g)$$

$$(3) A(g) + B_2(g) \rightarrow AB_2(g)$$

(4)
$$2AB_2(g) + A_2(g) + 2B_2(g)$$





PLAN: We are told that *P* is constant for this system, and the depiction shows that *V* does not change either. Since *T* changes, the volume could not remain the same unless the amount of gas in the system also changes.

SOLUTION:
$$n_1 T_1 = n_2 T_2$$
 $\frac{n_2}{n_1} = \frac{T_1}{T_2} = \frac{150 \text{ K}}{300 \text{ K}} = \frac{1}{2}$

Since *T* doubles, the total number of moles of gas must halve – i.e., the moles of product must be half the moles of reactant. This relationship is shown by equation (3).

$$\mathsf{A}(g) + \mathsf{B}_2(g) \to \mathsf{AB}_2(g)$$





The Ideal Gas Law and Gas Density

The density of a gas is

- directly proportional to its molar mass and
- inversely proportional to its temperature.

density =
$$\frac{m}{V}$$
 and moles = $\frac{m}{\mathcal{M}}$

$$PV = \frac{m}{\mathcal{M}} RT$$

$$\frac{m}{V} = d = \frac{\mathcal{M} \times P}{RT}$$





Calculating Gas Density

PROBLEM: Find the density (in g/L) of $CO_2(g)$ and the number of

molecules per liter

(a) at STP and (b) at room conditions (20.°C and 1.00 atm).

PLAN: We can use the molar mass of CO₂ to find its density from

the ideal gas equation.

SOLUTION: (a) At STP, or 273 K and 1.00 atm:

$$d = \frac{\mathcal{M} \times P}{RT} = \frac{44.01 \text{ g/mol} \times 1.00 \text{ atm}}{0.0821 \frac{\text{atm} \cdot \text{L}}{\text{mol} \cdot \text{K}} \times 273 \text{ K}} = 1.96 \text{ g/L}$$







SOLUTION: (b) At 20.°C and 1.00 atm:

$$T = 20.$$
°C + 273.15 = 293 K

$$d = \frac{\mathcal{M} \times P}{RT} = \frac{44.01 \text{ g/mol} \times 1.00 \text{ atm}}{0.0821 \frac{\text{atm} \cdot \text{L}}{\text{mol} \cdot \text{K}} \times 293 \text{ K}} = 1.83 \text{ g/L}$$

= 2.50×10^{22} molecules CO_2/L





Molar Mass from the Ideal Gas Law

$$n = \frac{m}{\mathcal{M}} = \frac{PV}{RT}$$

$$\mathcal{M} = \frac{mRT}{PV}$$





Finding the Molar Mass of a Volatile Liquid

PROBLEM:

An organic chemist isolates a colorless liquid from a petroleum sample. She places the liquid in a preweighed flask and puts the flask in boiling water, causing the liquid to vaporize and fill the flask with gas. She closes the flask and reweighs it. She obtains the following data:

Volume (
$$V$$
) of flask = 213 mL mass of flask + gas = 78.416 g

$$T = 100.0$$
°C $P = 754$ torr mass of flask = 77.834 g

Calculate the molar mass of the liquid.

PLAN: The variables V, T and P are given. We find the mass of the gas by subtracting the mass of the flask from the mass of the flask with the gas in it, and use this information to calculate \mathcal{M} .





SOLUTION:
$$m$$
 of gas = $(78.416 - 77.834) = 0.582$ g

$$V = 213 \text{ mL x} \frac{1 \text{ L}}{10^3 \text{ mL}} = 0.213 \text{ L}$$
 $T = 100.0^{\circ}\text{C} + 273.15 = 373.2 \text{ K}$

$$P = 754 \text{ terr x} \frac{1 \text{ atm}}{760 \text{ terr}} = 0.992 \text{ atm}$$

$$\mathcal{M} = \frac{mRT}{PV} = \frac{0.582 \text{ g x } 0.0821 \frac{\text{atm} \cdot \text{L}}{\text{mol} \cdot \text{K}} \text{ x } 373 \text{ K}}{0.213 \text{ L x } 0.992 \text{ atm}}$$

= 84.4 g/mol





Mixtures of Gases

- Gases mix homogeneously in any proportions.
 - Each gas in a mixture behaves as if it were the only gas present.
- The pressure exerted by each gas in a mixture is called its partial pressure.
- Dalton's Law of partial pressures states that the total pressure in a mixture is the sum of the partial pressures of the component gases.
- The partial pressure of a gas is proportional to its mole fraction:

$$P_{A} = X_{A} \times P_{\text{total}}$$
 $X_{A} = \frac{n_{A}}{n_{\text{total}}}$





Applying Dalton's Law of Partial Pressures

PROBLEM: In a study of O₂ uptake by muscle at high altitude, a physiologist prepares an atmosphere consisting of 79 mole % N₂, 17 mole % ¹⁶O₂, and 4.0 mole % ¹⁸O₂. (The isotope ¹⁸O will be measured to determine the O₂ uptake.) The pressure of the mixture is 0.75 atm to simulate high altitude. Calculate the mole fraction and partial pressure of ¹⁸O₂ in the mixture.

PLAN: Find X_{18O_2} and P_{18O_2} from P_{total} and mol % $^{18}O_2$.

mole % $^{18}O_2$ divide by 100

mole fraction, $X_{^{18}O_2}$ multiply by P_{total} partial pressure $P_{^{18}O_2}$





SOLUTION:

$$X_{18O_2} = \frac{4.0 \text{ mol } \%^{18}O_2}{100} = 0.040$$

$$P_{18O_2} = X_{18O_2} \times P_{total} = 0.040 \times 0.75 \text{ atm}$$
 = **0.030 atm**






Table 5.2 Vapor Pressure of Water (P_{H_2O}) + at Different T

7 (°C)	P _{L2} O(torr)	7 (⁰C)	P _{H₂O} (torr)
0	4.6	40	55.3
5	6.5	45	71.9
10	9.2	50	92.5
12	10.5	55	118.0
14	12.0	60	149.4
16	13.6	65	187.5
18	15.5	70	233.7
20	17.5	75	289.1
22	19.8	80	355.1
24	22.4	85	433.6
26	25.2	90	525.8
28	28.3	95	633.9
30	31.8	100	760.0
35	42.2		





Figure 5.11 Collecting a water-insoluble gaseous product and determining its pressure.



Copyright @ The McGraw-Hill Companies, Inc. Permission required for reproduction or display.





Calculating the Amount of Gas Collected over Water

PROBLEM: Acetylene (C_2H_2) is produced in the laboratory when calcium carbide (CaC_2) reacts with water:

$$CaC_2(s) + 2H_2O(l) \rightarrow C_2H_2(g) + Ca(OH)_2(aq)$$

A collected sample of acetylene has a total gas pressure of 738 torr and a volume of 523 mL. At the temperature of the gas (23°C), the vapor pressure of water is 21 torr. How many grams of acetylene are collected?

PLAN: The difference in pressures will give P for the C_2H_2 . The number of moles (n) is calculated from the ideal gas law and converted to mass using the molar mass.





PLAN:

Ptotal

subtract P for H₂O

Pof C₂H₂

use ideal gas law

n of C_2H_2

multiply by \mathcal{M}

mass of C₂H₂

SOLUTION:

$$P_{C_2H_2} = (738 - 21) \text{ torr} = 717 \text{ torr}$$

$$P = 717 \frac{\text{torr}}{\text{760 torr}} = 0.943 \text{ atm}$$

$$V = 523 \text{ mL } \times \frac{1 \text{ L}}{10^3 \text{ mL}} = 0.523 \text{ L}$$

$$T = 23$$
°C + 273.15 K = 296 K





SOLUTION:

$$n_{C_2H_2} = \frac{PV}{RT} = \frac{0.943 \text{ atm.} \times 0.523 L}{0.0821 \text{ mol·K}} \times 296 K}$$
 = **0.0203 mol**

$$0.0203 \frac{\text{mol}}{\text{mol}} \times \frac{26.04 \text{ g C}_2 \text{H}_2}{1 \frac{\text{mol}}{\text{C}_2 \text{H}_2}} = 0.529 \text{ g C}_2 \text{H}_2$$





The Ideal Gas Law and Stoichiometry




Figure 15.12

The relationships among the amount (mol, n) of gaseous reactant (or product) and the gas pressure (P), volume (V), and temperature (T).






Using Gas Variables to Find Amounts of Reactants and Products I

PROBLEM: What volume of H₂ gas at 765 torr and 225°C is needed to reduce 35.5 g of copper(II) oxide to form pure copper and water?

PLAN: Write a balanced equation. Convert the mass of copper (II) oxide to moles and find the moles of H₂, using the mole ratio from the balanced equation. Calculate the corresponding volume of H₂ using the ideal gas law.







SOLUTION:
$$CuO(s) + H_2(g) \rightarrow Cu(s) + H_2O(g)$$

$$35.5 \, \text{g CuO} \times \frac{1 \, \text{mol CuO}}{79.55 \, \text{g CuO}} \times \frac{1 \, \text{mol H}_2}{1 \, \text{mol CuO}} = 0.446 \, \text{mol H}_2$$

$$P = 765 \frac{\text{torr}}{760 \frac{\text{torr}}{\text{torr}}} = 1.01 \text{ atm}$$
 $T = 225 ^{\circ}\text{C} + 273.15 \text{ K} = 498 \text{ K}$

$$V = \frac{nRT}{P} = \frac{0.446 \text{ mol H}_2 \times 0.0821 \frac{\text{atm} \cdot \text{L}}{\text{mol K}} \times 498 \text{ K}}{1.01 \text{ atm}}$$





Using Gas Variables to Find Amounts of Reactants and Products II

PROBLEM: What mass of potassium chloride forms when 5.25 L of

chlorine gas at 0.950 atm and 293 K reacts with 17.0 g of

potassium metal?

PLAN: First we must write a balanced equation. Since the quantities of both reactants are given, we must next determine which reactant is limiting. We will use the ideal gas law to calculate the moles of Cl₂ present.

SOLUTION: The balanced equation is:

$$Cl_2(g) + 2K(s) \rightarrow 2KCl(s)$$

$$P = 0.950 \text{ atm}$$
 $V = 5.25 \text{ L}$

$$T$$
= 293 K n = unknown





$$n_{\text{Cl}_2} = \frac{PV}{RT} = \frac{0.950 \text{ atm} \times 5.25 \text{ L}}{0.0821 \times 10^{-10} \text{ mol} \cdot \text{K}} = 0.207 \text{ mol Cl}_2$$

For
$$Cl_2$$
: 0.207 $\frac{\text{mol } Cl_2}{\text{1-mol } Cl_2} = 0.414 \text{ mol } KCl_2$

For K:
$$17.0 \frac{g}{g} \frac{K}{K} \times \frac{1 \text{ mol } K}{39.10 \frac{g}{g} \frac{K}{K}} \times \frac{2 \text{ mol } KCl}{2 \text{ mol } K} = 0.435 \text{ KCl}$$

Cl₂ is the limiting reactant.





The Kinetic Molecular Theory: A Model for Gas Behavior

Postulate 1: Particle volume

Gas particles are tiny with large spaces between them. The volume of each particle is so small compared to the total volume of the gas that it is assumed to be zero.

Postulate 2: Particle motion

Gas particles are in constant, random, straight-line motion except when they collide with each other or with the container walls.

Postulate 3: Particles collision


Collisions are elastic, meaning that colliding particles exchange energy but do not lose any energy due to friction. Their *total kinetic energy is constant*. Between collisions the particles do not influence each other by attractive or repulsive forces.

* Gas conforms to theses postulates is called ideal.





Figure 5.13 Distribution of molecular speeds for N₂ at three temperatures.




Each particle in container it has molecular speed (u), most particles are moving near the most probable speed but some are much faster and other much slower.





A Movecontain eview of the gas law

- 1- Origin of pressure
- Pressure arises from countless collisions between gas particles and walls.
- Grater number of particles the more frequently colloid with the container, and so the greater the pressure.










2-A molecular view of Boyle's law.

Figure 5.1





 P_{ext} increases, T and n fixed



At any T, $P_{gas} = P_{ext}$ as particles hit the walls from an average distance, d_1 .

Higher $P_{\rm ext}$ causes lower V, which results in more collisions, because particles hit the walls from a shorter average distance ($d_2 < d_1$) collision frequency increase. As a result, $P_{\rm gas}$ increase untie it again = $P_{\rm ext.}$






3- A molecular view of Dalton's law

$P_{total} = P_A + P_B$

Figure 5.16



Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Increase number of particles lead to increase number of collision with the wall per second.






4- AUPhoteculatiview of Charles's law



Figure 5.17



At T_1 , $P_{gas} = P_{atm}$.

Higher T increases collision frequency, so $P_{gas} > P_{atm.}$


Thus, V increases until $P_{gas} = P_{atm}$ at T_2 .





5- A molecular view of Avogadro's law $V \propto n$

Figure 5.18



For a given amount, n_1 , of gas, $P_{gas} = P_{atm}$.

When gas is added to reach n_2 the collision frequency of the particles increases, so $P_{\text{gas}} > P_{\text{atm}}$.

As a result, V increases until $P_{gas} = P_{atm}$ again.





Kinetic Energy and Gas Behavior

At a given T, all gases in a sample have the same **average** kinetic energy.

$$E_{\rm k} = \frac{1}{2}$$
 mass x speed²

Kinetic energy depends on both the mass and the speed of a particle.


One mole of H_2 (2 g/mol) and 1 mole of O_2 (32 g/mol) occupied the same volume.

At the same T, a heavier gas particle moves more slowly than a lighter one. increase mass reduce the speed.





Figure 5.19 The relationship between molar mass and molecular speed.







Graham's Law of Effusion

Effusion is the process by which a gas escapes through a small hole in its container into an evacuated space.

Graham's law of effusion states that the rate of effusion of a gas is inversely proportional to the square root of its molar mass.


A lighter gas moves more quickly and therefore has a higher rate of effusion than a heavier gas at the same T.

Rate of effusion
$$\infty \frac{1}{\sqrt{\mathcal{M}}}$$






Figure 5.20 Effusion. Lighter (*black*) particles effuse faster than heavier (*red*) particles.



Rate
$$_{\mathcal{A}} = \sqrt{\frac{\mathcal{M}_{\mathcal{B}}}{\mathcal{M}_{\mathcal{A}}}}$$





Applying Graham's Law of Effusion

PROBLEM: A mixture of helium (He) and methane (CH₄) is placed in an effusion apparatus. Calculate the ratio of their effusion rates.

PLAN: The effusion rate is inversely proportional $\sqrt{\mathcal{M}}$ for each gas, so we find the molar mass for each substance using its formula and take the square root. The ratio of the effusion rates is the inverse of the ratio of these square roots.

SOLUTION: \mathcal{M} of CH₄ = 16.04 g/mol \mathcal{M} of He = 4.003 g/mol

$$\frac{\text{rate}}{\text{rate}} = \sqrt{\frac{16.04}{4.003}} = 2.002$$





Real Gases: Deviations from Ideal Behavior

- The kinetic-molecular model describes the behavior of ideal gases. Real gases deviate from this behavior.
- Real gases have real volume.
 - Gas particles are **not** points of mass, but have volumes determined by the sizes of their atoms and the bonds between them.
- Real gases do experience attractive and repulsive forces between their particles.
- Real gases deviate most from ideal behavior at *low temperature* and *high pressure*.





Table 5.3 Molar Volume of Some Common Gases at STP (0°C and 1 atm)

Gas	Molar Volume (L/mol)	Boiling Point (°C)
He	22.435	-268.9
H_2	22.432	-252.8
Ne	22.422	-246.1
Ideal gas	22.414	_
Ar	22.397	-185.9
N_2	22.396	-195.8
O_2	22.390	-183.0
CO	22.388	-191.5
Cl_2	22.184	-34.0
$\overline{NH_3}$	22.079	-33.4

Deviation increase as a boiling point increase





Figure 5.21 Deviations from ideal behavior with increasing external pressure







Figure 5.22

The effect of interparticle attractions on measured gas pressure.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display,



At ordinary P_{ext}, particles are too far apart to interact.

At moderately high P_{ext}, particles are close enough to interact.

Interparticle attractions (red arrows) lower the force of collisions with the container wall.






Figure 5.23 The effect of particle volume on measured gas volume.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

At ordinary P_{ext}, free volume ≈ container volume because particle volume is an insignificant portion of container volume.



At very high P_{ext}, free volume < container volume because particle volume becomes a significant portion of container volume.





The van der Waals equation

- The van der Waals equation adjusts the ideal gas law to take into account
 - the real volume of the gas particles and
 - the effect of interparticle attractions.

Van der Waals equation for *n* moles of a real gas

$$(P + \frac{n^2 a}{V^2})(V - nb) = nRT$$

The constant (*a)* relates to factors that influence the attraction between particles.

The constant *b* relates to particle volume.






Table 5.4 Van der Waals Constants for Some Common Gases

Gas	$a \left(\frac{\text{atm} \cdot L^2}{\text{mol}^2} \right)$	$b\left(\frac{L}{mol}\right)$
He	0.034	0.0237
Ne	0.211	0.0171
Ar	1.35	0.0322
Kr	2.32	0.0398
Xe	4.19	0.0511
H_2	0.244	0.0266
$\overline{N_2}$	1.39	0.0391
O_2	1.36	0.0318
\overline{Cl}_2	6.49	0.0562
CH_4	2.25	0.0428
CO	1.45	0.0395
CO_2	3.59	0.0427
NH_3	4.17	0.0371
H_2O	5.46	0.0305





Chapter 06

Lecture **Outline**

See separate *Image PowerPoint* slides for all figures and tables pre-inserted into PowerPoint without notes.





Chapter 6

Thermochemistry: Energy Flow and Chemical Change





Thermochemistry: Energy Flow and Chemical Change

- 6.1 Forms of Energy and Their Interconversion
- 6.2 Enthalpy: Chemical Change at Constant Pressure
- 6.3 Calorimetry: Measuring the Heat of a Chemical or Physical Change
- 6.4 Stoichiometry of Thermochemical Equations
- 6.5 Hess's Law: Finding ΔH of Any Reaction
- 6.6 Standard Enthalpies of Reaction (ΔH_{rxn}^{o})



Transfer and Interconversion of Energy

Thermodynamics is the study of energy and its transformations.

Thermochemistry is a branch of thermodynamics that deals with the heat involved in chemical and physical changes.

When energy is transferred from one object to another, it appears as *work* and *heat*.

The System and Its Surroundings

A meaningful study of any transfer of energy requires that we first clearly define both the *system* and its *surroundings*.

System + Surroundings = Universe

The *internal energy*, *E*, of a system is the sum of the potential and kinetic energies of all the particles present.

The total energy of the universe remains *constant*. A change in the energy of the system must be accompanied by an *equal* and *opposite* change in the energy of the surroundings.






UPLOADED BY AHMAD T JUNDI



Figure 6.1 Energy diagrams for the transfer of internal energy (*E*) between a system and its surroundings.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



$$\Delta E = E_{\text{final}} - E_{\text{initial}} = E_{\text{products}} - E_{\text{reactants}}$$








Figure 6.2 The two cases where energy is transferred as heat only.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



The system releases heat




B Energy absorbed as heat.

The system absorbs heat





Figure 6.3A The two cases where energy is transferred as work only.




A Energy released as work.

The system does work on the surroundings.





Figure 6.3B The two cases where energy is transferred as work only.



B Energy absorbed as work.

The system has work done on it by the surroundings.





Table 6.1 The Sign Conventions* for q, w, and ΔE

q	+	W	=	ΔΕ
+ (heat absorbed) + (heat absorbed) - (heat released) - (heat released)	-	(work done on) (work done by) (work done on) (work done by)		+ (energy <i>absorbed</i>) Depends on the <i>sizes</i> of <i>q</i> and <i>w</i> Depends on the <i>sizes</i> of <i>q</i> and <i>w</i> - (energy <i>released</i>)

^{*}From the perspective of the system.

Which of these are **not state functions?**


- A) volume B) temperature C) pressure D) heat E) energy
- A) internal energy B) volume C) work D) pressure E) enthalpy

A system which undergoes an adiabatic change (i.e., q = 0)

A system which undergoes an isothermal change (i.e., constant temperature)



Figure 6.4 Two different paths for the energy change of a system.



Even though q and w for the two paths are different, the total ΔE is the same for both.





The Law of Energy Conservation

The first law of Thermodynamics states that the total energy of the universe is constant.

Energy is conserved, and is neither created nor destroyed. Energy is transferred in the form of heat and/or work.

$$\Delta E_{\text{universe}} = \Delta E_{\text{system}} + \Delta E_{\text{surroundings}} = 0$$



Units of Energy

The SI unit of energy is the joule (J).

 $1 J = 1 kg \cdot m^2/s^2$

The **calorie** was once defined as the quantity of energy needed to raise the temperature of 1 g of water by 1°C.

1 cal = 4.184 J

The **British Thermal Unit (Btu)** is often used to rate appliances.

1 Btu is equivalent to 1055 J.



Determining the Change in Internal Energy of a System

PROBLEM: When gasoline burns in a car engine, the heat released causes the products CO_2 and H_2O to expand, which pushes the pistons outward. Excess heat is removed by the car's radiator. If the expanding gases do 451 J of work on the pistons and the system releases 325 J to the surroundings as heat, calculate the change in energy (ΔE) in J, kJ, and kcal.

PLAN: Define the system and surroundings and assign signs to q and w correctly. Then $\Delta E = q + w$. The answer can then be converted from J to kJ and to kcal.





SOLUTION:

Heat is given out by a chemical reaction, so it makes sense to define the system as the reactants and products involved. The pistons, the radiator and the rest of the car then comprise the surroundings.

Heat is given out by the system, so q = -325 J

The gases expand to push the pistons, so the system does work on the surroundings and w = -451 J

$$\Delta E = q + w = -325 \text{ J} + (-451 \text{ J}) = \boxed{-776 \text{ J}}$$

$$-776 - J \times \frac{1 \text{ kJ}}{10^3 \text{ J}} = -0.776 \text{ kJ} \times \frac{1 \text{ kcal}}{4.184 \text{ kJ}} = -0.185 \text{ kcal}$$





Enthalpy: Chemical Change at Constant Pressure


$$W = - P \Delta V$$

$$H = E + PV$$

where H is enthalpy

$$\Delta H = \Delta E + P \Delta V$$

$$q_p = \Delta E + P \Delta V = \Delta H$$



 $W = -P\Delta V$

An expanding gas pushing back the atmosphere does PV work ($w = -P \triangle V$).





When∆*H*≈ ∆*E*?

$\Delta H = \Delta E + P \Delta V$


- 1. Reactions that do not involve gases.
- 2. Reactions in which the number of moles of gas does not change.
- 3. Reactions in which the number of moles of gas does change but q is >>> $P\Delta V$.

Figure 6.6


Enthalpy diagrams for exothermic and endothermic processes.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

$$CH_4(g) + 2O_2(g) \rightarrow CO_2(g) + H_2O(g)$$



 $H_2O(s) \rightarrow H_2O(h)$



A Exothermic process

A Exothermic process Heat is given out.

B Endothermic process Heat is taken in.





Drawing Enthalpy Diagrams and Determining the Sign of ΔH

PROBLEM: In each of the following cases, determine the sign of ΔH , state whether the reaction is exothermic or endothermic, and draw an enthalpy diagram.

(a)
$$H_2(g) + \frac{1}{2}O_2(g) \rightarrow H_2O(h) + 285.8 \text{ kJ}$$

(b) 40.7 kJ +
$$H_2O(I) \rightarrow H_2O(g)$$

PLAN: From each equation, note whether heat is a "reactant" or a "product". If heat is taken in as a "reactant", the process is endothermic. If heat is released as a "product", the process is exothermic.


For the enthalpy diagram, the arrow always points from reactants to products. For endothermic reactions, the products are at a higher energy than the reactants, since the reactants take in heat to form the products.





SOLUTION: (a) $H_2(g) + \frac{1}{2}O_2(g) \rightarrow H_2O(1) + 285.8 \text{ kJ}$

Heat is a "product" for this reaction and is therefore given out, so the reaction is exothermic. The reactants are at a higher energy than the products.




EXOTHERMIC



SOLUTION: (b)
$$40.7 \text{ kJ} + \text{H}_2\text{O}(1) \longrightarrow \text{H}_2\text{O}(g)$$

Heat is a "reactant" in this reaction and is therefore absorbed, so the reaction is endothermic. The reactants are at a lower energy than the products.







UPLOADED BY AHMAD T JUND Calorimetry

heat capacity of a substance is the quantity of heat required to change its temperature by **1** K.

$$q \dot{\alpha} \Delta T$$

$$q = c$$
onstant x ΔT

$$q = heat \ capacity \times \Delta T$$

$$q$$
 = heat lost or gained c = specific heat capacity m = mass in g $\Delta T = T_{\text{final}} - T_{\text{initial}}$

The *specific heat capacity* (*c*) of a substance is the quantity of heat required to change the temperature of *1 gram* of the substance by *1 K*. Unit of c is (J/g·K)

$$q = c \times m \times \Delta T$$





Table 6.2 5 per chind Hear Capacities (c) of Some Elements, Compounds,

and Materials

Substance	Specific Heat Capacity (J/g·K)	Substance	Specific Heat Capacity (J/g·K)*		
Elements	Solid materials				
aluminum, Al	0.900	wood	1.76		
graphite,C	0.711	cement	0.88		
iron, Fe	0.450	glass	0.84		
copper, Cu	0.387	granite	0.79		
gold, Au	0.129	steel	0.45		
Compounds					
V	water, H ₂ O(<i>I</i>)				
ϵ	ethyl alcohol, C ₂ H ₅ OH(<i>I</i>)	2.46			
ϵ	ethylene glycol, (CH ₂ OH)	₂ (<i>I</i>) 2.42			
C	carbon tetrachloride, CCl	(<i>I</i>) 0.862			





Finding the Quantity of Heat from a Temperature Change

PROBLEM: A layer of copper welded to the bottom of a skillet weighs 125 g. How much heat is needed to raise the temperature of the copper layer from 25°C to 300.°C? The specific heat capacity (*c*) of Cu is 0.387 J/g·K.

PLAN: We know the mass (125 g) and c (0.387 J/g·K) of Cu and can find ΔT in °C, which equals ΔT in K. We can use the equation $q = c \times m \times \Delta T$ to calculate the heat.

SOLUTION: $\Delta T = T_{\text{final}} - T_{\text{initial}} = 300. - 25 = 275^{\circ}\text{C} = 275 \text{ K}$

$$q = c \times m \times \Delta T = \frac{0.387 \text{ J}}{-g \cdot \text{K}} \times 125 \text{ g} \times 275 \text{ K} = 1.33 \times 10^4 \text{ J}$$





Common types of calorimetry


1- Constant pressure calorimetry

Coffee-cup calorimeter.

Device used to find the specific heat capacity for solid does not dissolve or react with water.

$$-q_{solid} = q_{H_2O}$$

$$C_{solid} = \frac{C_{\text{H}_2\text{O}} \times \text{mass}_{\text{H}_2\text{O}} \times \Delta T_{\text{H}_2\text{O}}}{\text{mass}_{\text{solid}} \times \Delta T_{\text{solid}}}$$



Copyright @ The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

This device measures the heat transferred at constant pressure (q_P) .

UPLOADED BY:Ahmad Al-Jundi





Determining the Specific Heat Capacity of a Solid

PROBLEM: A 22.05 g solid is heated in a test-tube to 100.00°C and added to 50.00 g of water in a coffee-cup calorimeter. The water temperature changes from 25.10°C to 28.49°C. Find the specific heat capacity of the solid. $C_{H_2O} = 4.184 \text{ J/g·K}$

PLAN: Since the water and the solid are in contact, heat is transferred from the solid to the water until they reach the same T_{final} . In addition, the heat given out by the solid $(-q_{\text{solid}})$ is equal to the heat absorbed by the water (q_{water}) .

SOLUTION:

$$\Delta T_{\text{water}} = T_{\text{final}} - T_{\text{initial}} = (28.49^{\circ}\text{C} - 25.10^{\circ}\text{C}) = 3.39^{\circ}\text{C} = 3.39 \text{ K}$$

$$\Delta T_{\text{solid}} = T_{\text{final}} - T_{\text{initial}} = (28.49^{\circ}\text{C} - 100.00^{\circ}\text{C}) = -71.51^{\circ}\text{C} = -71.51 \text{ K}$$





UPLOADED BY AHMAD T JUNDI

Sample Problem 6.4

$$c_{solid} = \frac{c_{\text{H}_2\text{O}} \times \text{mass} \times \Delta T_{\text{H}_2\text{O}}}{\text{mass}_{\text{solid}} \times \Delta T_{\text{solid}}}$$

$$= \frac{4.184 \text{ J/g·K} \times 50.00 \text{ g x } 3.39 \text{ K}}{22.05 \text{ g x } (-71.51 \text{ K})} = \mathbf{0.450 \text{ J/g·K}}$$





Determining the Enthalpy Change of an Aqueous Reaction

PROBLEM: 50.0 mL of 0.500 M NaOH is placed in a coffee-cup calorimeter at 25.00°C and 25.0 mL of 0.500 M HCl is carefully added, also at 25.00°C. After stirring, the final temperature is 27.21°C. Calculate $q_{\rm soln}$ (in J) and the change in enthalpy, ΔH ,

(in kJ/mol of H₂O formed).

Assume that the total volume is the sum of the individual volumes, that d = 1.00 g/mL and c = 4.184 J/g·K

PLAN: Heat flows from the reaction (the system) to its surroundings (the solution). Since $-q_{rxn} = q_{soln}$, we can find the heat of the reaction by calculating the heat absorbed by the solution.





SOLUTION:

(a) To find q_{soln} :

Total mass (g) of the solution = $(25.0 \text{ mL} + 50.0 \text{ mL}) \times 1.00 \text{ g/mL} = 75.0 \text{ g}$

$$\Delta T_{\text{soln}} = 27.21^{\circ}\text{C} - 25.00^{\circ}\text{C} = 2.21^{\circ}\text{C} = 2.21 \text{ K}$$

$$q_{\text{soln}} = c_{\text{soln}} \times \text{mass}_{\text{soln}} \times \Delta T_{\text{soln}} = (4.184 \text{ J/g·K})(75.0 \text{ g})(2.21 \text{ K})$$

(b) To find ΔH_{rxn} we first need a balanced equation:

$$HCI(aq) + NaOH(aq) \rightarrow NaCI(aq) + H_2O(l)$$



For HCI:

25.0 mL HCl x
$$\frac{1 \text{ L}}{10^3 \text{ mL}}$$
 x $\frac{0.500 \text{ mol}}{1 \text{ L}}$ x $\frac{1 \text{ mol H}_2\text{O}}{1 \text{ mol HCl}}$ = 0.0125 mol H₂O

For NaOH:

50.0 mL NaOH x
$$\frac{1 \text{ L}}{10^3 \text{ mL}}$$
 x $\frac{0.500 \text{ mol}}{1 \text{ L}}$ x $\frac{1 \text{ mol H}_2\text{O}}{1 \text{ mol NaOH}}$ = 0.0250 mol H₂O

HCl is limiting, and the amount of H₂O formed is 0.0125 mol.

$$\Delta H_{\text{rxn}} = \frac{q_{\text{rxn}}}{\text{mol H}_2\text{O}} = \frac{-693 \text{ J}}{0.0125 \text{ mol}} \times \frac{1 \text{ kJ}}{10^3 \text{J}} = -55.4 \text{ kJ/mol H}_2\text{O}$$






1- Constant volume calorimetry

A bomb calorimeter.

Device used to measure the heat of combustion reaction

$$q_{v} = \Delta E + P \Delta V$$

$$q_v = \Delta E + 0$$



This device measures the heat released at constant volume (q_{V}) .





Calculating the Heat of a Combustion Reaction

PROBLEM: A manufacturer claims that its new dietetic dessert has "fewer than 10 Calories per serving." To test the claim, a chemist at the Department of Consumer Affairs places one serving in a bomb calorimeter and burns it in O₂. The initial temperature is 21.862°C and the temperature rises to 26.799°C. If the heat capacity of the calorimeter is 8.151 kJ/K, is the manufacturer's claim correct?

PLAN: When the dessert (system) burns, the heat released is absorbed by the calorimeter:

$$-q_{\text{system}} = q_{\text{calorimeter}}$$

To verify the energy provided by the dessert, we calculate $q_{
m calorimeter}$



SOLUTION:

$$\Delta T_{\text{calorimeter}} = T_{\text{final}} - T_{\text{initial}}$$

= 26.799°C - 21.862°C = 4.937°C = 4.937 K

 $q_{\text{calorimeter}}$ = heat capacity x ΔT = 8.151 kJ/K x 4.937 K = 40.24 kJ

The manufacturer's claim is true, since the heat produced is less than 10 Calories.



Stoichiometry of Thermochemical Equations

- A *thermochemical equation* is a balanced equation that includes ΔH_{rxn} .
- The sign of \(\Delta H \) indicates whether the reaction is exothermic or endothermic.
- The magnitude of ∆H is proportional to the amount of substance.
- The value of ∆H can be used in a calculation in the same way as a mole ratio.



Figure 6.11

The relationship between amount (mol) of substance and the energy (kJ) transferred as heat during a reaction.





Using the Enthalpy Change of a Reaction (ΔH) to Find Amounts of Substance

PROBLEM: The major source of aluminum in the world is bauxite (mostly aluminum oxide). Its thermal decomposition can be represented by the equation

$$Al_2O_3(s) \rightarrow 2Al(s) + \frac{3}{2}O_2(g) \Delta H_{rxn} = 1676 \text{ kJ}$$

If aluminum is produced this way, how many grams of aluminum can form when 1.000x10³ kJ of heat is transferred?

PLAN: From the balanced equation and ΔH , we see that 2 mol of Al is formed when 1676 kJ of heat is absorbed.







SOLUTION:

$$1.000 \times 10^{3} \text{ kJ x} \quad \frac{2 \text{ mol Al}}{1676 \text{ kJ}} \times \quad \frac{26.98 \text{ g Al}}{1 \text{ mol Al}} = 32.20 \text{ g Al}$$





Hess' Law

Hess's law states that the enthalpy change of an overall process is the sum of the enthalpy changes of its individual steps.

$$\Delta H_{\text{overall}} = \Delta H_1 + \Delta H_2 + \dots + \Delta H_n$$

 ΔH for an overall reaction can be calculated if the ΔH values for the individual steps are known.



Calculating ΔH for an overall process

- Identify the target equation, the step whose ΔH is unknown.
 - Note the amount of each reactant and product.
- Manipulate each equation with known ∆H values so that the target amount of each substance is on the correct side of the equation.
 - Change the sign of ΔH when you reverse an equation.
 - Multiply amount (mol) and ΔH by the same factor.
- Add the manipulated equations and their resulting ΔH
 values to get the target equation and its ΔH.
 - All substances except those in the target equation must cancel.



Using Hess's Law to Calculate an Unknown Δ*H*

PROBLEM: Two gaseous pollutants that form in auto exhausts are CO and NO. An environmental chemist is studying ways to convert them to less harmful gases through the following reaction:

$$CO(g) + NO(g) \rightarrow CO_2(g) + \frac{1}{2}N_2(g) \Delta H = ?$$

Given the following information, calculate the unknown ΔH :

Equation A:
$$CO(g) + \frac{1}{2}O_2(g) \rightarrow CO_2(g) \Delta H_A = -283.0 \text{ kJ}$$

Equation B:
$$N_2(g) + O_2(g) \rightarrow 2NO(g) \Delta H_B = 180.6 \text{ kJ}$$

PLAN: Manipulate Equations A and/or B and their ΔH values to get to the target equation and its ΔH . All substances except those in the target equation must cancel.





SOLUTION:

Multiply Equation B by ½ and reverse it:

$$NO(g) \rightarrow \frac{1}{2} N_2(g) + \frac{1}{2} O_2(g); \Delta H = -90.3 \text{ kJ}$$

Add the manipulated equations together:

Equation A:
$$CO(g) + \frac{1}{2}O_2(g) \rightarrow CO_2(g)$$

$$\Delta H = -283.0 \text{ kJ}$$

$$\frac{1}{2}$$
 Equation B: NO(g) $\rightarrow \frac{1}{2} N_2(g) + \frac{1}{2} O_2(g) \Delta H = -90.3 \text{ kJ}$

$$\Delta H = -90.3 \text{ kJ}$$

(reversed) _____

$$CO(g) + NO(g) \rightarrow CO_2(g) + \frac{1}{2}N_2(g)$$

$$\Delta H_{\rm rxn} = -373.3 \text{ kJ}$$



Table 6.3 Selected Standard Enthalpies of Formation at 25°C (298K)

Formula $\Delta H_{f}^{\circ}(kJ/m)$	ol) Formula	$\Delta H_{\rm f}^{\prime}$ (kJ/mol)	Formula $\Delta H_f^{\prime}(kJ/mol)$
Calcium $Ca(s) 0$ $CaO(s) -635.1$ $CaCO_3(s) -1206.9$	$Cl_2(g)$ $HCl(g)$ Hydrogen	0 -92.3	Silver $Ag(s) \qquad 0$ $AgCl(s) \qquad -127.0$
Carbon C(graphite) 0	$H(g)$ $H_2(g)$	218 0	Sodium Na(<i>s</i>) 0
C(diamond) 1.9 $CO(g)$ -110.5	Nitrogen N ₂ (<i>g</i>)	0	Na(<i>g</i>) 107.8 NaCl(<i>s</i>) -411.1
$CO_2(g)$ -393.5 $CH_4(g)$ -74.9	NH ₃ (<i>g</i>) NO(<i>g</i>)	-45.9 90.3	Sulfur S ₈ (rhombic) 0
$CH_3OH(I)$ -238.6 HCN(g) 135 $CS_s(I)$ 87.9	Oxygen $O_2(g)$	0	S_8 (monoclinic) 0.3 $SO_2(g)$ -296.8
Chlorine $Cl(g)$ 121.0	$O_3(g)$ $H_2O(g)$ $H_2O(h)$	143 -241.8 -285.8 Y:Ahmad Al-Jundi	SO ₃ (<i>g</i>) -396.0

Writing Formation Equations

PROBLEM: Write balanced equations for the formation of 1 mol of the following compounds from their elements in their standard states and include ΔH_f .

- (a) Silver chloride, AgCl, a solid at standard conditions.
- **(b)** Calcium carbonate, CaCO₃, a solid at standard conditions.
- (c) Hydrogen cyanide, HCN, a gas at standard conditions.

PLAN: Write the elements as reactants and 1 mol of the compound as the product formed. Make sure all substances are in their standard states. Balance the equations and find the value of ΔH_f in Table 6.3 or Appendix B.



SOLUTION:

(a) Silver chloride, AgCl, a solid at standard conditions.

$$Ag(s) + \frac{1}{2}Cl_2(g) \longrightarrow AgCl(s)$$
 $\Delta H_f^s = -127.0 \text{ kJ}$

(b) Calcium carbonate, CaCO₃, a solid at standard conditions.

$$Ca(s) + C(graphite) + \frac{3}{2}O_2(g) \rightarrow CaCO_3(s)$$
 $\Delta H_f^s = -1206.9 \text{ kJ}$

(c) Hydrogen cyanide, HCN, a gas at standard conditions.

$$\frac{1}{2}H_2(g) + C(graphite) + \frac{1}{2}N_2(g) \rightarrow HCN(g)$$
 $\Delta H_f^e = 135 \text{ kJ}$







Figure 6.10

The two-step process for determining ΔH_{rxn}^{o} from ΔH_{f}^{o} values.





Calculating ΔH_{rxn}^{\prime} from ΔH_{f}^{\prime} Values

PROBLEM: Nitric acid, whose worldwide annual production is about 8 billion kilograms, is used to make many products, including fertilizer, dyes, and explosives. The first step in the industrial production process is the oxidation of ammonia:

$$4NH_3(g) + 5O_2(g) \rightarrow 4NO(g) + 6H_2O(g)$$

Calculate ΔH_{rxn}^{\prime} from ΔH_f^{\prime} values.

PLAN: Use the ΔH_f values from Table 6.3 or Appendix B and apply the equation

$$\Delta H_{\text{rxn}} = \sum m \Delta H_{\text{f}}^{\prime} \text{ (products)} - \sum n \Delta H_{\text{f}}^{\prime} \text{ (reactants)}$$



SOLUTION:

$$\Delta H_{\rm rxn} = \sum m \Delta H_{\rm f}^{\prime} \, ({\rm products}) - \sum n \Delta H_{\rm f}^{\prime} \, ({\rm reactants})$$

$$\Delta H_{\rm rxn} = [4(\Delta H_{\rm f}^{\prime} \, {\rm of} \, {\rm NO}(g) + 6(\Delta H_{\rm f}^{\prime} \, {\rm of} \, {\rm H_2O}(g)]$$

$$- [4(\Delta H \, {\rm of} \, {\rm NH_3}(g) + 5(\Delta H \, {\rm of} \, {\rm O_2}(g)]$$

=
$$(4 \text{ mol})(90.3 \text{ kJ/mol}) + (6 \text{ mol})(-241.8 \text{ kJ/mol}) -$$

$$[(4 \text{ mol})(-45.9 \text{ kJ/mol}) + (5 \text{ mol})(0 \text{ kJ/mol})]$$

$$= -906 \text{ kJ}$$


$$\Delta H_{\rm rxn} = -906 \text{ kJ}$$





Figure 6.11 The trapping of heat by the atmosphere.


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.







Connect UPLOADED BY AHMAD T JUNDI



Chapter 07

Lecture Outline

See separate *Image PowerPoint* slides for all figures and tables pre-inserted into PowerPoint without notes.





Chapter 7

Quantum Theory and Atomic Structure





Quantum Theory and Atomic Structure

7.1 The Nature of Light

7.2 Atomic Spectra

7.3 The Wave-Particle Duality of Matter and Energy

7.4 The Quantum-Mechanical Model of the Atom





The Wave Nature of Light

Visible light is a type of *electromagnetic radiation*.

The wave properties of electromagnetic radiation are described by three variables:

- frequency (v), cycles per second
- wavelength (λ), the distance a wave travels in one cycle
- amplitude, the height of a wave crest or depth of a trough.

The *speed of light* is a constant:

$$c = v \times \lambda$$

 $= 3.00 \times 10^{8} \text{ m/s in a vacuum}$





Figure 7.1 The reciprocal relationship of frequency and wavelength.

Copyright @ The McGraw-Hill Companies, Inc. Permission required for reproduction or display. Wavelength = distance per cycle $\lambda_A = 2\lambda_B = 4\lambda_C$ Wavelength A В C Frequency = cycles per second $v_{\rm A} = \frac{1}{2}v_{\rm B} = \frac{1}{4}v_{\rm C}$

UPLODED BY Ahmad Jundi





Figure 7.2 Differing amplitude (brightness, or intensity) of a wave.









Figure 7.3 Regions of the electromagnetic spectrum.



Light of single wavelength is called **monochromatic** (one color) Whereas light of many wavelength is **polychromatic**.



Interconverting Wavelength and Frequency

PROBLEM: A dental hygienist uses x-rays (λ = 1.00Å) to take a series of dental radiographs while the patient listens to a radio station (λ = 325 cm) and looks out the window at the blue sky (λ = 473 nm). What is the frequency (in s⁻¹) of the electromagnetic radiation from each source? (Assume that the radiation travels at the speed of light, 3.00x10⁸ m/s.)

PLAN: Use the equation $c = v\lambda$ to convert wavelength to frequency. Wavelengths need to be in meters because c has units of m/s.

wavelength in units given

use conversion factors $1 \text{ Å} = 10^{-10} \text{ m}$

wavelength in m

$$v = \frac{c}{\lambda}$$

frequency (s-1 or Hz)





SOLUTION:

For the x-rays:
$$\lambda = 1.00 \text{ Å x} \frac{10^{-10} \text{ m}}{1 \text{ Å}} = 1.00 \text{ x} 10^{-10} \text{ m}$$

$$v = \frac{C}{\lambda} = \frac{3.00 \times 10^8 \text{ m/s}}{1.00 \times 10^{-10} \text{ m}} = 3.00 \times 10^{18} \text{ s}^{-1}$$

$$= 3.00 \times 10^{18} \text{ s}^{-1}$$

For the radio signal:
$$v = \frac{c}{\lambda} = \frac{3.00 \times 10^8 \text{ m/s}}{325 \text{ cm x } \frac{10^{-2} \text{ m}}{1 \text{ cm}}} = 9.23 \times 10^7 \text{ s}^{-1}$$

$$= 9.23 \times 10^7 \text{ s}^{-1}$$

For the blue sky:
$$v = \frac{c}{\lambda} = \frac{3.00 \times 10^8 \text{ m/s}}{473 \text{ nm x} \frac{10^{-9} \text{ m}}{1 \text{ cm}}} = 6.34 \times 10^{14} \text{ s}^{-1}$$

$$= 6.34 \times 10^{14} \text{ s}^{-1}$$





Classical distinction between energy and matter


Different behaviors of waves of energy and particles of matter

1- Refraction and Dispersion

When the light wave pass from one media into another the speed of the wave change.

White light separates (disperses) into its components colors when it pass

through the prism






2- Diffraction and interference

When wave strike strike the edge of an object it bend around it (diffraction). When of light pass through two adjacent slits the nearby emerging circular waves interact through the process of interference.

- If the crest of waves coincide in phase (the interference is constructive).
- If the crest coincide with troughs out of phase (the interference is destructive).

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.





Energy and frequency

A solid object emits visible light when it is heated to about 1000 K. This is called *blackbody radiation*.

The *color* (and the intensity) of the light changes as the temperature changes. Color is related to *wavelength* and *frequency*, while temperature is related to *energy*.

Energy is therefore related to frequency and wavelength:

In frequency lundi

$$E = nhv$$

E = energyn is a positive integer (quantum number)h is Planck's constant





The Quantum Theory of Energy

Any object (including atoms) can emit or absorb only *certain quantities* of energy.

Energy is *quantized*, it occurs in fixed quantities, rather than being continuous. Each fixed quantity of energy is called a *quantum*.

An atom changes its energy state by emitting or absorbing one or more *quanta* of energy.

 $\Delta E = nhv$ where *n* can only be a whole number.



UPigure 7.6 MAD T JUThe photoelectric effect.


Einstine proposed that the light itself is particulate quantized in to tiny "bundles" of energy (photons).

Each atom change its energy one its absorb or emit one photon(one particles of light) whose energy related to frequency not to its amplitude.

$$\mathsf{E}_{\mathsf{photon}} = h_{\mathcal{V}} = \mathsf{A} \mathsf{E}$$

Two features of the photoelectric effect:

- 1- **Threshold frequency**: Intensity is related to number of photon but not to energy of each, a photon of certain minimum energy must be absorb to free an electron from the surface, since energy depends on frequency the theory predicts a threshold frequency.
- 2- **Absence of the time lag**: the current is week in dim light because fewer photons of enough energy can free fewer electrons per unit time.





Calculating the Energy of Radiation from Its Wavelength

PROBLEM: A cook uses a microwave oven to heat a meal. The wavelength of the radiation is 1.20 cm. What is the energy of one photon of this microwave radiation?

PLAN: We know λ in cm, so we convert to m and find the frequency using the speed of light. We then find the energy of one photon using E = hv.

SOLUTION:

$$E = h_{V} = \frac{hc}{\lambda} = \frac{(6.626 \times 10^{-34}) \text{ J} \cdot \text{s})(3.00 \times 10^{8} \text{ m/s})}{(1.20 \text{ cm})(\frac{10^{-2} \text{ m}}{1 \text{ cm}})} = \frac{1.66 \times 10^{-23} \text{ J}}{}$$





Figure 7.7A The line spectrum of hydrogen.






Figure 7.7B The line spectra of Hg and Sr.









Figure 7.8 Three series of spectral lines of atomic hydrogen.



Rydberg equation
$$\frac{1}{\lambda} = R \left[\frac{1}{n_1^2} - \frac{1}{n_2^2} \right]$$

R is the Rydberg constant = 1.096776x10⁷ m⁻¹

for the visible series, $n_1 = 2$ and $n_2 = 3, 4, 5, ...$





The Bohr Model of the Hydrogen Atom

Bohr's atomic model postulated the following:

- The H atom has only certain energy levels, which Bohr called stationary states.
 - Each state is associated with a fixed circular orbit of the electron around the nucleus.
 - The higher the energy level, the farther the orbit is from the nucleus.
 - When the H electron is in the first orbit, the atom is in its lowest energy state, called the *ground state*.



- The atom does not radiate energy while in one of its stationary states.
- The atom changes to another stationary state only by absorbing or emitting a photon.
 - The energy of the photon ($h\nu$) equals the difference between the energies of the two energy states.
 - When the E electron is in any orbit higher than n = 1, the atom is in an *excited state*.

Figure 7.9 A quantum "staircase" as an analogy for atomic energy levels.








Figure 7.10 The Bohr explanation of three series of spectral lines emitted by the H atom.








UPLOADED BY AHMAD T JUNDI A tabletop analogy for the H atom's energy.





$$\Delta E = E_{\text{final}} - E_{\text{initial}} = -2.18 \times 10^{-18} \text{ J} \left(\frac{1}{n^2_{\text{final}}} - \frac{1}{n^2_{\text{initial}}} \right)$$





Determining ΔE and λ of an Electron Transition

PROBLEM: A hydrogen atom absorbs a photon of UV light (see Figure 7.10) and its electron enters the *n* = 4 energy level. Calculate (a) the change in energy of the atom and (b) the wavelength (in nm) of the photon.

PLAN: (a) The H atom absorbs energy, so $E_{\text{final}} > E_{\text{initial}}$. We are given $n_{\text{final}} = 4$, and Figure 7.10 shows that $n_{\text{initial}} = 1$ because a UV photon is absorbed. We apply Equation 7.4 to find ΔE .

(b) Once we know ΔE , we find frequency and wavelength.





SOLUTION:

(a)
$$\Delta E = -2.18 \times 10^{-18} \text{ J} \left(\frac{1}{n^2_{\text{final}}} - \frac{1}{n^2_{\text{initial}}} \right) = -2.18 \times 10^{-18} \text{ J} \left(\frac{1}{4^2} - \frac{1}{1^2} \right)$$

= -2.18x10⁻¹⁸ J
$$\left(\frac{1}{16} - \frac{1}{1}\right)$$
 = 2.04x10⁻¹⁸ J

(b)
$$\lambda = \frac{hc}{\Delta E} = \frac{(6.626 \times 10^{-34} \text{ J} \cdot \text{s})(3.00 \times 10^8 \text{ m/s})}{2.04 \times 10^{-18} \text{ J}} = 9.74 \times 10^{-8} \text{ m}$$


$$9.74 \times 10^{-8} \text{ m x} \quad \frac{1 \text{ nm}}{10^{-9} \text{ m}} = 97.4 \text{ nm}$$






Figure 7.11 Measuring chlorophyll a concentration in leaf extract.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.





Two types of spectra

- 1- Excitation spectrum
- 2- Emission spectrum
 UPLODED BY Ahmad Jundi





The Wave-Particle Duality of Matter and Energy

Matter and Energy are alternate forms of the same entity.

$$E = mc^2$$

All matter exhibits properties of both particles and waves. Electrons have wave-like motion and therefore have only certain allowable frequencies and energies.

Matter behaves as though it moves in a wave, and the *de Broglie wavelength* for any particle is given by:


$$\lambda = \frac{h}{mu} \qquad m = \text{mass}$$

$$u = \text{speed in m/s}$$





Figure 7.12 Wave motion in restricted systems.



ST28ENTS-HUB.com

Table 7.1 The de Broglie Wavelengths of Several Objects

Substance	Mass (g)	Speed (m/s)	λ (m)
slow electron	9x10 ⁻²⁸	1.0	7x10 ⁻⁴
fast electron	9x10 ⁻²⁸	5.9x10 ⁶	1x10 ⁻¹
alpha particle	6.6x10 ⁻²⁴	1.5x10 ⁷	$7x^{0}10^{-1}$
one-gram mass	1.0	0.01	7x ⁵ 10 ⁻²⁹
baseball	142	25.0	2x10 ⁻³⁴
Earth	6.0x10 ²⁷	3.0x10 ⁴	4x10 ⁻⁶³





Calculating the de Broglie Wavelength of an Electron

PROBLEM: Find the de Broglie wavelength of an electron with a

speed of $1.00x10^6$ m/s (electron mass = $9.11x10^{-31}$ kg;

 $h = 6.626 \times 10^{-34} \text{ kg} \cdot \text{m}^2/\text{s}$).

PLAN: We know the speed and mass of the electron, so we substitute these into Equation 7.5 to find λ .

SOLUTION:

$$\lambda = \frac{h}{mu}$$

$$\lambda = \frac{6.626 \times 10^{-34} \text{ kg} \cdot \text{m}^2/\text{s}}{9.11 \times 10^{-31} \text{ kg} \times 1.00 \times 10^6 \text{ m/s}}$$


$$= 7.27 \times 10^{-10} \,\mathrm{m}$$






Figure 7.13 Diffraction patterns of aluminum with x-rays and electrons.

Copyright @ The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



PSSC Physics @ 1965, Education Development Center, Inc.

Copyright @ The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



D.C. Heath & Company/Education Development Center, Inc.

x-ray diffraction of aluminum foil

electron diffraction of aluminum foil



CLASSICAL THEORY

Matter particulate, massive

Energy continuous, wavelike

Figure 7.14

Major observations and theories leading from classical theory to quantum theory

Since *matter* is discontinuous and particulate, perhaps *energy* is discontinuous and particulate.

Observation

Theory

Blackbody radiation — Planck: Energy is quantized; only certain values

allowed

Atomic line spectra —— Bohr: Energy of atoms is quantized; photon

emitted when electron changes orbit.





Figure 7.14 continued

Since <i>energy</i> is wavelike, perhaps <i>matter</i> is wavelike.				
Observation	Theory			
Davisson/Germer: Electron beam is diffracted by metal crystal		tter travels in waves; energy of s quantized due to wave motion of ons		
Since <i>matte</i> r has mass, perhaps <i>energy</i> has mass				
Observation	Theory			
Compton: Photon's wavelength increases (momentum decreases) after colliding with electron	Einstein/deBroglie	e: Mass and energy are equivalent; particles have wavelength and photons have momentum.		

QUANTUM THEORY

UPLODED BY Alfrida Jungi particulate, massive, wavelike ploaded By: mariam Qadah





Heisenberg's Uncertainty Principle

Heisenberg's Uncertainty Principle states that it is not possible to know both the position *and* momentum of a moving particle at the same time.

$$\Delta x m \Delta u \ge \frac{h}{4\pi}$$
 $x = \text{position}$ $u = \text{speed}$

The more accurately we know the speed, the less accurately we know the position, and vice versa.






The Quantum-Mechanical Model of the Atom

The matter-wave of the electron occupies the space near the nucleus and is continuously influenced by it.

The **Schrödinger wave equation** allows us to solve for the energy states associated with a particular atomic orbital.

The square of the wave function gives the *probability* density, a measure of the *probability* of finding an electron of a particular energy in a particular region of the atom.









Figure 7.15

Electron probability density in the ground-state H atom.







Quantum Numbers and Atomic Orbitals

An atomic orbital is specified by three quantum numbers.

The *principal* quantum number (*n*) is a positive integer.

The value of *n* indicates the relative *size* of the orbital and therefore its relative *distance* from the nucleus.

The *angular momentum* quantum number *(I)* is an integer from 0 to (n-1).

The value of / indicates the **shape** of the orbital.

The *magnetic* quantum number (m_j) is an integer with values from -1 to +1

The value of m_i indicates the spatial *orientation* of the orbital.



Table 7.2 The Hierarchy of Quantum Numbers for Atomic Orbitals

Name, Symbol (Property)

Allowed Values


Quantum Numbers


Principal, *n* (size, energy)

Positive integer (1, 2, 3, ...)

Angular momentum, *I* 0 to *n* – 1 (shape)

Magnetic, m_I (orientation) -I,...,0,...,+I







Determining Quantum Numbers for an Energy Level

PROBLEM: What values of the angular momentum (I) and magnetic (m_I) quantum numbers are allowed for a principal quantum number (I) of 3? How many orbitals are allowed for I = 3?

PLAN: Values of l are determined from the value for n, since l can take values from 0 to (n-1). The values of m_l then follow from the values of l.

SOLUTION: For n = 3, allowed values of l are l and 2

For
$$l = 0$$
 $m_l = 0$

For
$$I = 1$$
 $m_I = -1$, 0, or +1

For
$$l = 2$$
 $m_l = -2, -1, 0, +1, or +2$

There are 9 m_I values and therefore 9 orbitals with n = 3.





Determining Sublevel Names and Orbital Quantum Numbers

PROBLEM: Give the name, magnetic quantum numbers, and number of orbitals for each sublevel with the following quantum numbers:

(a)
$$n = 3$$
, $l = 2$ (b) $n = 2$, $l = 0$ (c) $n = 5$, $l = 1$ (d) $n = 4$, $l = 3$

PLAN: Combine the n value and l designation to name the sublevel. Knowing l, we can find m_l and the number of orbitals.

SOLUTION:

	n	1	sublevel name	possible m_I values #	f of orbitals
(a)	3	2	3 <i>d</i>	-2, -1, 0, 1, 2	5
(b)	2	0	2 <i>s</i>	0	1
(c)	5	1	5 <i>p</i>	-1, 0, 1	3
(d)	4	3	4 <i>f</i>	-3, -2, -1, 0, 1, 2, 3	7





Identifying Incorrect Quantum Numbers

PROBLEM: What is wrong with each of the following quantum numbers designations and/or sublevel names?

	n	1	m_{l}	Name
(a)	1	1	0	1 <i>p</i>
(b)	4	3	+1	4 <i>d</i>
(c)	3	1	-2	3 <i>p</i>

SOLUTION:

- (a) A sublevel with n = 1 can only have l = 0, not l = 1. The only possible sublevel name is 1s.
- **(b)** A sublevel with I = 3 is an f sublevel, to a d sublevel. The name should be 4f.
- (c) A sublevel with I = 1 can only have m_I values of -1, 0, or +1, not -2.





Figure 7.16

The 1s, 2s, and 3s orbitals.








Figure 7.17 The 2p orbitals.






Figure 7.18 The 3*d* orbitals.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.






Figure 7.18





Figure 7.19 The $4f_{xyz}$ orbital, one of the seven 4f orbitals.




Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Figure 7.20 Energy levels of the H atom.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




Energy

1*s* –







Chapter 08

Lecture Outline

See separate *Image PowerPoint* slides for all figures and tables pre-inserted into PowerPoint without notes.





UPLODED BY Ahmad Jundi

Chapter 8

Electron Configuration and Chemical Periodicity





Electron Configuration and Chemical Periodicity

8.1 Characteristics of Many-Electron Atoms

8.2 The Quantum-Mechanical Model and the Periodic Table

8.3 Trends in Three Atomic Properties

8.4 Atomic Properties and Chemical Reactivity



Figure 8.1 The effect of electron spin.






Table 8.1 Summary of Quantum Numbers of Electrons in Atoms

Name	Symbol	Permitted Values	Property
principal	n	positive integers (1, 2, 3,)	orbital energy (size)
angular momentum	1	integers from 0 to <i>n</i> -1	orbital shape (The <i>I</i> values 0, 1, 2, and 3 correspond to <i>s</i> , <i>p</i> , <i>d</i> , and <i>f</i> orbitals, respectively.)
magnetic	m_{l}	integers from -/to 0 to +/	orbital orientation
spin	$m_{_S}$	$+\frac{1}{2}$ or $-\frac{1}{2}$	direction of e ⁻ spin





Quantum Numbers and The Exclusion Principle

Each electron in any atom is described completely by a set of *four* quantum numbers.

The first three quantum numbers describe the orbital, while the fourth quantum number describes electron spin.

Pauli's **exclusion principle** states that *no two electrons in the same atom can have the same four quantum numbers*.

An atomic orbital can hold a *maximum of two electrons* and they must have *opposing spins*.



Factors Affecting Atomic Orbital Energies

- The energies of atomic orbitals are affected by
 - nuclear charge (Z) and
 - shielding by other electrons.
- A higher nuclear charge increases nucleus-electron interactions and lowers sublevel energy.
- **Shielding** by other electrons reduces the full nuclear charge to an *effective nuclear charge* (Z_{eff}).
 - Z_{eff} is the nuclear charge an electron actually experiences.
- Orbital shape also affects sublevel energy.




Shielding and Orbital Energy

- Electrons in the same energy level shield each other to some extent.
- Electrons in *inner* energy levels shield the outer electrons very effectively.
 - The farther from the nucleus an electron is, the lower the $Z_{\rm eff}$ for that particular electron.

Figure 8.2 Penetration and sublevel energy.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Orbital shape causes electrons in some orbitals to "penetrate" close to the nucleus.

Penetration increases nuclear attraction and decreases shielding.

Splitting of Levels into Sublevels

Each energy level is split into *sublevels* of differing energy. Splitting is caused by penetration and its effect on shielding.

For a given *n* value, a lower *l* value indicates a lower energy sublevel.

Order of sublevel energies: s








Figure 8.3

Order for filling energy sublevels with electrons.

In general, energies of sublevels increase as n increases (1 < 2 < 3, etc.) and as l increases (s).

As *n* increases, some sublevels overlap.



Electron Configurations and Orbital Diagrams

Electron configuration is indicated by a shorthand notation:

Orbital diagrams make use of a box, circle, or line for each orbital in the energy level. An arrow is used to represent an electron *and* its spin.










A vertical orbital diagram for the Li ground state.





Building Orbital Diagrams

The **aufbau principle** is applied – electrons are always placed in the lowest energy sublevel available.

The **exclusion principle** states that each orbital may contain a maximum of 2 electrons, which must have opposite spins.

He
$$(Z=2)$$
 1s² $\uparrow \downarrow$






Building Orbital Diagrams

Hund's rule specifies that when orbitals of equal energy are available, the lowest energy electron configuration has the maximum number of unpaired electrons with parallel spins.

N (
$$Z = 7$$
) 1 $s^2 2s^2 2p^3$





Determining Quantum Numbers from Orbital Diagrams

PROBLEM: Write a set of quantum numbers for the third electron and a set for the eighth electron of the F atom.

PLAN: Identify the electron of interest and note its level (n), sublevel, (l), orbital (m_l) and spin (m_s). Count the electrons in the order in which they are placed in the diagram.

SOLUTION:

F (Z=9)
$$1s^2 2s^2 2p^5$$

$$1s \quad 2s \quad 2p$$

For the 3rd electron: n = 2, I = 0, $m_I = 0$, $m_S = +\frac{1}{2}$

For the 8th electron: n = 2, I = 1, $m_I = -1$, $m_S = -\frac{1}{2}$





Partial Orbital Diagrams and Condensed Configurations

A *partial orbital diagram* shows only the highest energy sublevels being filled.

A *condensed electron configuration* has the element symbol of the *previous* noble gas in square brackets.

All has the condensed configuration [Ne]3 s^2 3 p^1



Table 8.2 Partial Orbital Diagrams and Electron Configurations* for the Elements in Period 3.

Atomic Number	Element		al Orbital Diagram nd 3 <i>p</i> Sublevels Only)	Full Electron Configuration [†]	Condensed Electron Configuration
		3s	3 <i>p</i>		
11	Na	\uparrow		$[1s^22s^22p^6]$ 3s ¹	[Ne] $3s^1$
12	Mg	$\uparrow\downarrow$		$[1s^22s^22p^6]$ $3s^2$	[Ne] $3s^2$
13	Al	$\uparrow\downarrow$	\uparrow	$[1s^22s^22p^6] 3s^23p^1$	[Ne] $3s^2 3p^1$
14	Si	$\uparrow\downarrow$	\uparrow \uparrow	$[1s^22s^22p^6] 3s^23p^2$	[Ne] $3s^2 3p^2$
15	P	$\uparrow\downarrow$	\uparrow \uparrow \uparrow	$[1s^22s^22p^6] 3s^23p^3$	[Ne] $3s^2 3p^3$
16	S	$\uparrow\downarrow$	$\uparrow\downarrow$ \uparrow \uparrow	$[1s^22s^22p^6] 3s^23p^4$	[Ne] $3s^2 3p^4$
17	C1	$\uparrow\downarrow$	$\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow$	$[1s^22s^22p^6] 3s^23p^5$	[Ne] $3s^2 3p^5$
18	Ar	$\uparrow\downarrow$	$\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow\downarrow$	$[1s^22s^22p^6]\ 3s^23p^6$	[Ne] $3s^2 3p^6$

^{*}Colored type indicates the sublevel to which the last electron is added.

[†]The full configuration is not usually written with square brackets; they are included here to show how the [Ne] designation arises.



Electron Configuration and Group

Elements in the same group of the periodic table have the same outer electron configuration.

Elements in the same group of the periodic table exhibit similar chemical behavior.

Similar outer electron configurations correlate with similar chemical behavior.



Figure 8.4 Condensed electron configurations in the first three periods.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

		1A (1)						
	1	1 H 1 <i>s</i> ¹	2A (2)	3A (13)	4A (14)	5A (15)	6A (16)	7A (17)
Period	2	3 Li [He] 2 <i>s</i> ¹	4 Be [He] 2 <i>s</i> ²	5 B [He] 2s ² 2p ¹	6 C [He] 2 <i>s</i> ² 2 <i>p</i> ²	7 N [He] 2 <i>s</i> ² 2 <i>p</i> ³	8 O [He] 2 <i>s</i> ² 2 <i>p</i> ⁴	9 F [He] 2 <i>s</i> ² 2 <i>p</i> ⁵
	3	11 Na [Ne] 3 <i>s</i> ¹	12 Mg [Ne] 3 <i>s</i> ²	13 AI [Ne] 3 <i>s</i> ² 3 <i>p</i> ¹	14 Si [Ne] 3 <i>s</i> ² 3 <i>p</i> ²	15 P [Ne] 3 <i>s</i> ² 3 <i>p</i> ³	16 S [Ne] 3 <i>s</i> ² 3 <i>p</i> ⁴	17 CI [Ne] 3 <i>s</i> ² 3 <i>p</i> ⁵





8A (18)

He

 $1s^{2}$

10

Ne

[He] $2s^22p^6$

18 **Ar**

[Ne] $3s^23p^6$

Table 8.3 Partial Orbital Diagrams and Electron Configurations* for the Elements in Period 4.

Atomic Number	Element	Partial Orbital Diagram (4 <i>s</i> , 3 <i>d</i> , and 4 <i>p</i> Sublevels Only)	Full Electron Configuration	Condensed Electron Configuration
19	K	4s 3d 4p	$1s^22s^22p^63s^23p^64s^1$	[Ar] $4s^1$
20	Ca	↑	$1s^2 2s^2 2p^6 3s^2 3p^6 4s^2$	$[Ar] 4s^2$
21	Sc	$\uparrow\downarrow$ \uparrow	$1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^1$	$[Ar] 4s^2 3d^1$
22	Ti	$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	$1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^2$	$[Ar] 4s^2 3d^2$
23	V	$\uparrow \downarrow \qquad \qquad \uparrow \qquad \uparrow \qquad \uparrow \qquad \qquad \boxed{}$	$1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^3$	$[Ar] 4s^2 3d^3$
24	Cr	$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	$1s^2 2s^2 2p^6 3s^2 3p^6 4s^1 3d^5$	$[Ar] 4s^1 3d^5$
25	Mn	$\uparrow \downarrow \qquad \qquad \uparrow \qquad \uparrow \qquad \uparrow \qquad \uparrow \qquad \uparrow \qquad \qquad $	$1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^5$	[Ar] $4s^2 3d^5$
26	Fe	$\uparrow \downarrow \qquad \uparrow \downarrow \uparrow $	$1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^6$	$[Ar] 4s^2 3d^6$
27	Co	$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	$1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^7$	$[Ar] 4s^2 3d^7$

^{*}Colored type indicates the sublevel to which the last electron is added.





Table 8.3 Partial Orbital Diagrams and Electron Configurations* for the Elements in Period 4.

Atomic Number	Element		artial Orbital Diagran 4 <i>s,</i> 3 <i>d,</i> and 4 <i>p</i> Sublev		Full Electron Configuration	Condensed Electron Configuration
28	Ni	$\uparrow\downarrow$	$\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow\uparrow\uparrow$		$1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^8$	$[Ar] 4s^2 3d^8$
29	Cu	\uparrow			$1s^2 2s^2 2p^6 3s^2 3p^6 4s^1 3d^{10}$	[Ar] $4s^13d^{10}$
30	Zn	$\uparrow\downarrow$			$1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^{10}$	[Ar] $4s^23d^{10}$
31	Ga	$\uparrow\downarrow$	$\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow$	\uparrow	$1s^22s^22p^63s^23p^64s^23d^{10}4p^1$	[Ar] $4s^2 3d^{10}4p^1$
32	Ge	$\uparrow\downarrow$	$\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow\uparrow\downarrow$	\uparrow \uparrow	$1s^22s^22p^63s^23p^64s^23d^{10}4p^2$	[Ar] $4s^2 3d^{10} 4p^2$
33	As	$\uparrow\downarrow$	$\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow\uparrow\downarrow$	\uparrow \uparrow \uparrow	$1s^22s^22p^63s^23p^64s^23d^{10}4p^3$	[Ar] $4s^2 3d^{10}4p^3$
34	Se	$\uparrow\downarrow$	$\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow$	$\uparrow\downarrow$ \uparrow \uparrow	$1s^22s^22p^63s^23p^64s^23d^{10}4p^4$	[Ar] $4s^2 3d^{10} 4p^4$
35	Br	$\uparrow\downarrow$	$\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow$	$\uparrow\downarrow\uparrow\downarrow\uparrow$	$1s^22s^22p^63s^23p^64s^23d^{10}4p^5$	[Ar] $4s^2 3d^{10}4p^5$
36	Kr	$\uparrow\downarrow$	$\boxed{\uparrow\downarrow \uparrow\downarrow \uparrow\downarrow \uparrow\downarrow \uparrow\downarrow \uparrow\downarrow}$	$\uparrow\downarrow\uparrow\uparrow\downarrow\uparrow\downarrow$	$1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^{10} 4p^6$	[Ar] $4s^2 3d^{10} 4p^6$

^{*}Colored type indicates the sublevel to which the last electron is added.





Figure 8.5 A periodic table of partial ground-state electron configurations.


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

						0							2.5						
		Main- Elem (s bl	nents												Ma	in-Grou (<i>p</i> bl	p Eleme lock)	nts	
		1A (1) ns ¹																	8A (18) ns ² np ⁶
	1	1 H	2A (2)											3A (13)	4A (14)	5A (15)	6A (16)	7A (17)	2 He
		1s ¹	ns ²											ns ² np ¹	ns ² np ²	ns ² np ³	ns ² np ⁴	ns ² np ⁵	152
Ne	2	3 Li	4 Be											5 B	6 C	7 N	8 O	9 F	10 Ne
energy level	-	2s ¹	2s ²				Tr	ansition		nts				2s ² 2p ¹	2s ² 2p ²	2s ² 2p ³	2s ² 2p ⁴	2s ² 2p ⁵	2s ² 2p ⁶
ener		11	12					(0 0	lock)					13	14	15	16	17	18
ө рө	3	Na	Mg	3B	4B	5B	6B	7B		— 8B —		1B	2B	Al	Si	P	S	CI	Ar
cupi		3s1	3s ²	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	3s ² 3p ¹	3s ² 3p ²	3s ² 3p ³	3s ² 3p ⁴	3s ² 3p ⁵	3s ² 3p ⁶
t oc	4	19 K	20 Ca	21 Sc	22 Ti	23 V	24 Cr	25 Mn	26 Fe	27 Co	28 Ni	29 Cu	30 Zn	31 Ga	32 Ge	33 As	34 Se	35 Br	36 K r
Period number: highest occupied	-	4s ¹	4s ²	4s ² 3d ¹	4s ² 3d ²	4s ² 3d ³	4s ¹ 3d ⁵	4s ² 3d ⁵	4s ² 3d ⁶	4s ² 3d ⁷	4s ² 3d ⁸	4s ¹ 3d ¹⁰			4s ² 4p ²	4s ² 4p ³	4s ² 4p ⁴	4s ² 4p ⁵	4s ² 4p ⁶
r: hić		37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
nbei	5	Rb	Sr	Υ	Zr	Nb	Мо	Тс	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Те	1	Xe
nu		5s ¹	5s ²	5s ² 4d ¹	5s ² 4d ²	5s ¹ 4d ⁴	5s ¹ 4d ⁵	5s ² 4d ⁵	5s ¹ 4d ⁷	5s ¹ 4d ⁸	4d ¹⁰	5s ¹ 4d ¹⁰		-	5s ² 5p ²	5s ² 5p ³	5s ² 5p ⁴	5s ² 5p ⁵	5s ² 5p ⁶
riod	6	55 Cs	56 Ba	57 La*	72 Hf	73 Ta	74 W	75 Re	76 Os	77 Ir	78 Pt	79 Au	80 Hg	81 TI	82 Pb	83 Bi	84 Po	85 At	86 Rn
Pe	0	6s ¹	$6s^2$	$6s^25d^1$	$6s^25d^2$	$6s^25d^3$	6s ² 5d ⁴	6s ² 5d ⁵	$6s^25d^6$	6s ² 5d ⁷	6s ¹ 5d ⁹	6s ¹ 5d ¹⁰	The second second		$6s^26p^2$	$6s^{2}6p^{3}$	6s ² 6p ⁴	6s ² 6p ⁵	6s ² 6p ⁶
		87	88	89	104	105	106	107	108	109	110	111	112	113	114	115	116		118
	7	Fr	Ra	Ac**	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg	Cn			200 000			
		7s ¹	7s ²	7s ² 6d ¹	$7s^26d^2$	7s ² 6d ³	7 <i>s</i> ² 6 <i>d</i> ⁴	7s ² 6d ⁵	7 <i>s</i> ² 6 <i>d</i> ⁶	7s ² 6d ⁷	7s ² 6d ⁸	7s ² 6d ⁹	7s ² 6d ¹⁰	$7s^27p^1$	7s ² 7p ²	$7s^27p^3$	7s ² 7p ⁴		7s ² 7p ⁶
	Inner Transition Elements (f block)																		
			2020	58	59	60	61	62	63	64	65	66	67	68	69	70	71		
	6	*Lanth	anides	Се	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er 212	Tm	Yb	Lu		
				6s ² 4f ¹ 5d ¹	6s ² 4f ³	6s ² 4f ⁴	6s ² 4f ⁵	6s ² 4f ⁶	6s ² 4f ⁷	6s ² 4f ⁷ 5d ¹	6s ² 4f ⁹	6s ² 4f ¹⁰		6s ² 4f ¹²	6s ² 4f ¹³		6s ² 4f ¹⁴ 5d ¹		
	7	**Ac	tinides	90 Th	91 Pa	92 U	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr		
	,	7.0					7s ² 5f ⁴ 6d ¹												
												11171			1000	0.000		4	





Figure 8.6 Orbital filling and the periodic table.



The order in which the orbitals are filled can be obtained directly from the periodic table.






UPLOADED BY AHMAD T JUNDI

Aid to memorizing sublevel filling order.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



The *n* value is constant horizontally. The *I* value is constant vertically. *n* + *I* is constant diagonally.





Categories of Electrons

Inner (core) electrons are those an atom has in common with the pervious noble gas and any *completed* transition series.

Outer electrons are those in the *highest* energy level (highest *n* value).

Valence electrons are those involved in forming compounds.

For **main group** elements, the valence electrons **are** the outer electrons.

For **transition elements**, the valence electrons include the outer electrons and any (*n*-1)*d* electrons.



elements:

Sample Problem 8.2

Determining Electron Configurations

PROBLEM: Using the periodic table on the inside cover of the text (not Figure 8.5 or Table 8.3), give the full and condensed electron configurations, partial orbital diagrams showing valence electrons only, and number of inner electrons for the following

(a) potassium (b) technetium (c) lead (K;
$$Z=19$$
) (Tc; $Z=43$) (Pb; $Z=82$)

PLAN: The atomic number gives the number of electrons, and the periodic table shows the order for filling orbitals. The partial orbital diagram includes all electrons added after the previous noble gas except those in filled inner sublevels.



SOLUTION:

(a) For K (Z = 19)


full configuration

 $1s^22s^22p^63s^23p^64s^1$


condensed configuration


[Ar] $4s^1$

partial orbital diagram



45





There are 18 inner electrons.

SOLUTION:

(b) For Tc (Z = 43)


full configuration

 $1s^22s^22p^63s^23p^64s^23d^{10}4p^65s^24d^6$

condensed configuration

[Kr]5*s*²4*d*⁵

partial orbital diagram






There are 36 inner electrons.

SOLUTION:

(a) For Pb (Z = 82)

full configuration $1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^{10} 4p^6 5s^2 4d^{10} 5p^6 6s^2 4f^{14} 5d^{10} 6p^2$ condensed configuration [Xe] $6s^2 4f^{14} 5d^{10} 6p^2$

partial orbital diagram




There are 78 inner electrons.




Trends in Atomic Size


Defining atomic size.



A. The metallic radius of aluminum.



B. The covalent radius of chlorine.



C. Known covalent radii and distances between nuclei can be used to find unknown radii.





UPLOADED BY AHMAD TTKENDS IN Atomic Size

Atomic size varies within both groups and periods as a result of two opposing influences:

1- Atomic size *increases* as the principal quantum number *n increases*.

As *n* increases, the probability that the outer electrons will be farther from the nucleus increases.

2- Atomic size *decreases* as the effective nuclear charge Z_{eff} *increases*.

As Z_{eff} increases, the outer electrons are pulled closer to the nucleus.

For *main group elements*:

atomic size *increases* down a group in the periodic table and *decreases* across a period.






Transition elements

- 1- **Down a transition group**, n increase, but shielding by an additional level of inner electrons results in only a small size increase from period 4 to 5 and non from 5 to 6.
- 2- Across a transition series, atomic size shrinks through the first two or three elements because of increasing the nuclear charge, but from then on, size remain relatively constant because the shielding by the inner d electrons counteract the increase in Zeff.
- 3- Transition series affects atomic size in neighboring main groups, d electrons cause a size decrease from group 2A to group 3A.



Figure 8.8

Atomic radii of the main-group and transition elements.











Figure 8.9 Periodicity of atomic radius.







Ranking Elements by Atomic Size

PROBLEM: Using only the periodic table (not Figure 8.8), rank each set of main-group elements in order of *decreasing* atomic size:

- (a) Ca, Mg, Sr (b) K, Ga, Ca
- **(c)** Br, Rb, Kr **(d)** Sr, Ca, Rb

PLAN: Locate each element on the periodic table. Main-group elements increase in size down a group and decrease in size across the period.

SOLUTION:

(a) Sr > Ca > Mg

Ca, Mg, and Sr are in Group 2A. Size increases down the group.

(b) K > Ca > Ga

K, Ga, and Ca are all in Period 4. Size decreases across the period.

(c) Rb > Br > Kr

Rb is the largest because it has one more energy level than the other elements. Kr is smaller than Br because Kr is further to the right in the same period.

(d) Rb > Sr > Ca

Ca is the smallest because it has one fewer energy level. Sr is smaller than Rb because it is smaller to the right in the same period.



Trends in Ionization Energy

lonization energy (IE) is the energy required for the *complete removal* of 1 mol of electrons from 1 mol of gaseous atoms or ions.

Atoms with a *low IE* tend to form *cations*.

Atoms with a *high IE* tend to form *anions* (except the

noble gases).

Ionization energy tends to *decrease* down a group and *increase* across a period.

Atom_(g)
$$\longrightarrow$$
 Ion⁺ + e⁻ $\Delta E = IE_1 > 0$
Ion⁺ \longrightarrow Ion²⁺ + e⁻ $\Delta E = IE_2$ (always> IE_1)





Figure 8.10 Periodicity of first ionization energy (IE₁).





Figure 8.11 First ionization energies of the main-group elements.





Ranking Elements by First Ionization Energy

PROBLEM: Using the periodic table only, rank the elements in each of the following sets in order of *decreasing* IE₁:

- (a) Kr, He, Ar
- **(b)** Sb, Te, Sn

(c) K, Ca, Rb

(d) I, Xe, Cs

PLAN: Find each element on the periodic table. IE₁ generally decreases down a group and increases across a period.

SOLUTION:

(a) He > Ar > Kr

Kr, He, and Ar are in Group 8A. IE₁ decreases down the group.

SOLUTION:

(b) Te > Sb > Sn

Sb, Te, and Sn are in Period 5. IE₁ increases across a period.

(c) Ca > K > Rb

K has a higher IE₁ than Rb because K is higher up in Group 1A. Ca has a higher IE₁ than K because Ca is further to the right in Period 4.

(d) Xe > I > Cs

Xe has a higher IE₁ than I because Xe is further to the right in the same period. Cs has a lower IE₁ than I because it is further to the left in a higher period.



Figure 8.12 The first three ionization energies of beryllium.




Table 8.4 Successive Ionization Energies of the Elements Lithium Through Sodium

		Number		Ionization Energy (MJ/mol)*								
Z	Element	of Valence Electrons	IE ₁	IE ₂	IE ₃	IE ₄	IE ₅	IE ₆	IE ₇	IE ₈	IE ₉	IE ₁₀
3	Li	1	0.52	7.30	11.81							
4	Be	2	0.90	1.76	14.85	21.01			CORE	ELECTR	ONS	
5	В	3	0.80	2.43	3.66	25.02	32.82					
6	C	4	1.09	2.35	4.62	6.22	37.83	47.28				
7	N	5	1.40	2.86	4.58	7.48	9.44	53.27	64.36			
8	O	6	1.31	3.39	5.30	7.47	10.98	13.33	71.33	84.08		
9	F	7	1.68	3.37	6.05	8.41	11.02	15.16	17.87	92.04	106.43	
10	Ne	8	2.08	3.95	6.12	9.37	12.18	15.24	20.00	23.07	115.38	131.43
11	Na	1	0.50	4.56	6.91	9.54	13.35	16.61	20.11	25.49	28.93	141.37

^{*}MJ/mol, or megajoules per mole = 10^3 kJ/mol.





Identifying an Element from Its Ionization Energies

PROBLEM: Name the Period 3 element with the following ionization energies (in kJ/mol) and write its electron configuration:

IE ₁	IE ₂	IE ₃	IE_4	IE ₅	IE ₆
1012	1903	2910	4956	6278	22,230


PLAN: Look for a large increase in IE, which occurs after all valence electrons have been removed.

SOLUTION:

The largest increase occurs after IE_5 , that is, after the 5th valence electron has been removed. The Period 3 element with 5 valence electrons is **phosphorus** (**P**; Z=15).

The complete electron configuration is $1s^22s^22p^63s^23p^3$.





Trends in Electron Affinity

Electron Affinity (EA) is the energy change that occurs when 1 mol of electrons is added to 1 mol of gaseous atoms or ions.

Atoms with a *low EA* tend to form *cations*. Atoms with a *high EA* tend to form *anions*.

The trends in electron affinity are not as regular as those for atomic size or IE.

Figure 8.13 Electron affinities of the main-group elements (in kJ/mol).

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Сорупе		COIGWI	iii Companii	CO, 1110. 1 CI1	mosion requ	anca for rep	roddollori or	uispiay.
1A (1)							V	8A (18)
H -72.8	2A (2)	-	3A (13)	4A (14)	5A (15)	6A (16)	7A (17)	He (0.0)
Li -59.6	Be ≤0		B -26.7	C – 122	N +7	O -141	F - 328	Ne (+29)
Na -52.9	Mg ≤0		AI -42.5	Si - 134	P -72.0	S -200	CI -349	Ar (+35)
K -48.4	Ca -2.37		Ga -28.9	Ge - 119	As -78.2	Se -195	Br -325	Kr (+39)
Rb -46.9	Sr -5.03		In –28.9	Sn – 107	Sb -103	Te – 190	l -295	Xe (+41)
Cs -45.5	Ba -13.95		TI –19.3	Pb -35.1	Bi -91.3	Po – 183	At -270	Rn (+41)





Behavior Patterns for IE and EA

Reactive nonmetals have high IEs and highly negative EAs.

These elements attract electrons strongly and tend to form negative ions in ionic compounds.


Reactive metals have low IEs and slightly negative EAs. These elements lose electrons easily and tend to form positive ions in ionic compounds.

Noble gases have very high IEs and slightly positive EAs. These elements tend to neither lose nor gain electrons.



Figure 8.14 Trends in three atomic properties.

`Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.








Metallic Behavior

- Metals are typically shiny solids with moderate to high melting points.
- Metals are good conductors of heat and electricity, and can easily be shaped.
- Metals tend to lose electrons and form cations, i.e., they are easily oxidized.
- Metals are generally strong reducing agents.
- Most metals form ionic oxides, which are basic in aqueous solution.



Figure 8.15 Trends in metallic behavior.





Acid-Base Behavior of Oxides

Main-group metals form *ionic oxides*, which are *basic* in aqueous solution.

Main-group nonmetals form *covalent oxides*, which are *acidic* in aqueous solution.


Some metals and metalloids form *amphoteric oxides*, which can act as acids or bases in water:

$$Al_2O_3(s) + 6HCl(aq) \rightarrow 2AlCl_3(aq) + 3H_2O(1)$$

$$Al_2O_3(s) + 2NaOH(aq) \rightarrow 2NaAl(OH)_4(aq)$$



Figure 8.16 Acid-base behavior of some element oxides.



Oxides become more basic down a group and more acidic across a period.





Electron configurations of Monatomic Ions

Elements at either end of a period gain or lose electrons to attain a filled outer level. The resulting ion will have a *noble gas electron configuration* and is said to be *isoelectronic* with that noble gas.

Na(1
$$s^22s^22p^63s^1$$
) \rightarrow e⁻ + Na⁺([He]2 s^22p^6)
[isoelectronic with Ne]

Br([Ar]
$$4s^23d^{10}4p^5$$
) + $e^- \rightarrow Br^-$ ([Ar] $4s^23d^{10}4p^6$)
[isoelectronic with Kr]



Figure 8.47 BY AHMain group elements whose ions have noble gas electron configurations.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display. **8A** (18)2A 1A (2)(1) 6A **7A** He Li 2 (16)Ne Na F 3 Mg Ar Period 8 S CI K Ca Kr 5 Se Br Sr Rb Xe 5 Cs Ba 6 Gain Lose





electrons

electrons

Electron Configurations of Monatomic Ions

A *pseudo-noble gas configuration* is attained when a metal atom empties its highest energy level.

The ion attains the stability of empty ns and np sublevels and a filled (n-1)d sublevel.

$$Sn([Kr]5s^24d^{10}5p^2) \rightarrow 4e^- + Sn^{4+} ([Kr]4d^{10})$$

A metal may lose only the *np* electrons to attain an *inert pair configuration*.

The ion attains the stability of a filled ns and (n-1)d sublevels.

$$Sn([Kr]5s^24d^{10}5p^2) \rightarrow 2e^- + Sn^{2+} ([Kr]5s^24d^{10})$$



Sample Problem 8.6

Writing Electron Configurations of Main-Group lons

PROBLEM: Using condensed electron configurations, write reactions for the formation of the common ions of the following elements:

(a) Iodine (
$$Z = 53$$
) (b) Potassium ($Z = 19$) (c) Indium ($Z = 49$)

PLAN: Identify the position of each element on the periodic table and recall that:

- Ions of elements in Groups 1A(1), 2A(2), 6A(16), and 7A(17) are usually isoelectronic with the nearest noble gas.
- Metals in Groups 3A(13) to 5A(15) can lose the *ns* and *np* electrons or just the *np* electrons.






Sample Problem 8.6

SOLUTION:

- (a) Iodine (Z = 53) is in Group 7A(17) and will gain one electron to be isoelectronic with Xe: I ([Kr] $5s^24d^{10}5p^5$) + e⁻ \rightarrow I⁻ ([Kr] $5s^24d^{10}5p^6$)
- (b) Potassium (Z = 19) is in Group 1A(1) and will lose one electron to be isoelectronic with Ar: $K([Ar] 4s^1) \rightarrow K^+([Ar]) + e^-$
- (c) Indium (Z = 49) is in Group 3A(13) and can lose either one electron or three electrons: In ([Kr] $5s^24d^{10}5p^1$) \rightarrow In⁺ ([Kr] $5s^24d^{10}$) + e⁻ In ([Kr] $5s^24d^{10}5p^1$) \rightarrow In³⁺([Kr] $4d^{10}$) + 3e⁻



Figure 8.18 The crossover of sublevel energies in Period 4.







Magnetic Properties of Transition Metal Ions

A species with one or more unpaired electrons exhibits *paramagnetism* – it is attracted by a magnetic field.


Ag
$$(Z=47)$$

$$\uparrow \qquad \uparrow \downarrow \qquad 5p$$

A species with all its electrons paired exhibits diamagnetism – it is not attracted (and is slightly repelled) by a magnetic field.



Figure 8.19 Measuring the magnetic behavior of a sample.



The apparent mass of a diamagnetic substance is unaffected by the magnetic field.


The apparent mass of a paramagnetic substance increases as it is attracted by the magnetic field.





Magnetic Properties of Transition Metal ions

Magnetic behavior can provide evidence for the electron configuration of a given ion.



Ti²⁺ has 2 unpaired electrons and is paramagnetic, providing evidence that the 4*s* electrons are lost before the 3*d* electrons.



Sample Problem 8.7

Writing Electron Configurations and Predicting Magnetic Behavior of Transition Metal Ions

PROBLEM:

Use condensed electron configurations to write the reaction for the formation of each transition metal ion, and predict whether the ion is paramagnetic or diamagnetic.

(a)
$$Mn^{2+}(Z = 25)$$

(b)
$$Cr^{3+}(Z = 24)$$

(c)
$$Hg^{2+}(Z = 80)$$

PLAN:

Write the condensed electron configuration for each atom, recalling the irregularity for Cr. Remove electrons, beginning with the *ns* electrons, and determine if there are any unpaired electrons.



Sample Problem 8.7

SOLUTION:

(a) $Mn^{2+}(Z = 25)$ $Mn ([Ar] 4s^23d^5) \rightarrow Mn^{2+}([Ar] 3d^5) + 2e^-$ Since there are 5 d electrons they are all unpaired. Mn^{2+} is **paramagnetic**.

(b)
$$Cr^{3+}(Z = 24)$$
 $Cr([Ar] 4s^{1}3d^{5}) \rightarrow Cr^{3+}([Ar] 3d^{8}) + 3e^{-}$

Since there are 3 d electrons they are all unpaired. Cr^{3+} is paramagnetic.

(c)
$$Hg^{2+}(Z = 80)$$
 $Hg([Xe] 6s^24f^{4}5d^{10}) \rightarrow Hg^{2+}([Xe] 4f^{4}5d^{10}) + 2e^{-}$

The 4f and the 5d sublevels are filled, so there are no unpaired electrons. Hg²⁺ is **diamagnetic**.



Ionic Size vs. Atomic Size

Cations are *smaller* than their parent atoms while anions are *larger*.

Ionic radius *increases* down a group as *n* increases.

Cation size *decreases* as charge *increases*.

An *isoelectronic series* is a series of ions that have the same electron configuration. Within the series, ion size *decreases* with increasing nuclear charge.



Figure 8.20

Ionic radius.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.









Figure 8.21

Ionic vs. atomic radii.







Sample Problem 8.8

Ranking lons by Size

PROBLEM: Rank each set of ions in order of *decreasing* size, and

explain your ranking:

(a) Ca^{2+} , Sr^{2+} , Mg^{2+} (b) K^+ , S^{2-} , Cl^- (c) Au^+ , Au^{3+}

PLAN: Find the position of each element on the periodic table and

apply the trends for ionic size.

SOLUTION:

(a) $Sr^{2+} > Ca^{2+} > Mg^{2+}$

All these ions are from Group 2A, so size increases down the group.



Sample Problem 8.8

SOLUTION:

(b)
$$S^{2-} > CI^- > K^+$$


These ions are isoelectronic, so size decreases as nuclear charge increases.

(c) $Au^+ > Au^{3+}$

Cation size decreases as charge increases.







Chapter 09

Lecture Outline

See separate *Image PowerPoint* slides for all figures and tables pre-inserted into PowerPoint without notes.





UPLOADED BY:Ahmad Jundi

Chapter 9

Models of Chemical Bonding






Models of Chemical Bonding

- 9.1 Atomic Properties and Chemical Bonds
- 9.2 The Ionic Bonding Model
- 9.3 The Covalent Bonding Model
- 9.4 Bond Energy and Chemical Change
- 9.5 Between the Extremes: Electronegativity and Bond Polarity





Figure 9.1 A comparison of metals and nonmetals.



METAL NONMETAL ATOM		
Larger	Smaller	
Lower	Higher	
Lower	Higher	
Less negative	More negative	
	Larger Lower Lower	





Types of Chemical Bonding

lonic bonding involves the *transfer* of electrons and is usually observed when a *metal* bonds to a *nonmetal*.

Covalent bonding involves the **sharing** of electrons and is usually observed when a **nonmetal** bonds to a **nonmetal**.


Metallic bonding involves *electron pooling* and occurs when a *metal* bonds to another *metal*.






Figure 9.2 Three models of chemical bonding.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.




A lonic bonding



B Covalent bonding






C Metallic bonding





Figure 9.3 AHMAD T Jordations in bond type among Period 3 (black type) and Group 4A (red type) elements.








Lewis Electron-Dot Symbols

To draw the Lewis symbol for any main-group element:

- Note the A-group number, which gives the number of valence electrons.
- Place one dot at a time on each of the four sides of the element symbol.
- Keep adding dots, pairing them, until all are used up.

Example:

Nitrogen, N, is in Group 5A and therefore has 5 valence electrons.







Lewis Symbols and Bonding

For a *metal*, the *total* number of dots in the Lewis symbol is the number of electrons the atom loses to form a cation.

For a *nonmetal*, the number of unpaired dots equals

- the number of electrons the atom *gains* to form an anion
- or the number it **shares** to form covalent bonds.

The *octet rule* states that when atoms bond, they lose, gain, or share electrons to attain a *filled outer level of 8 electrons* (or 2, for H and Li).





Figure 9.4

Lewis electron-dot symbols for elements in Periods 2 and 3.

	1A(1)	2A(2)	
	ns ¹	ns ²	
2	• Li	•Be•	
3	• Na	•Mg•	

3A(13)	4A(14)	5A(15)	6A(16)	7A(17)	8A(18)
ns ² np ¹	ns ² np ²	ns ² np ³	ns ² np ⁴	ns ² np ⁵	ns ² np ⁶
• B •		: Z ·	: 0 •	: F.	Ne:
• AI •	· Si ·	. P	: s ·	: CI :	: Ar

The Ionic Bonding Model

An ionic bond is formed when a metal *transfers* electrons to a nonmetal to form *ions*, which attract each other to give a solid compound.

The total number of electrons lost by the metal atom(s) equals the total number of electrons gained by the nonmetal atoms.





Figure 9.5 Three ways to depict electron transfer in the formation of Li⁺ and F⁻.

Electron configurations Li $1s^22s^1 + F 1s^22p^5 \rightarrow Li^+ 1s^2 + F^- 1s^22s^22p^6$

Orbital diagrams

Lewis electron-dot symbols





Sample Problem 9.1

Depicting Ion Formation

PROBLEM: Use partial orbital diagrams and Lewis symbols to depict the formation of Na⁺ and O²⁻ ions from the atoms, and determine the formula of the compound formed.

PLAN: Draw orbital diagrams and Lewis symbols for Na and O atoms. To attain filled outer levels, Na loses one electron and O gains two. Two Na atoms are needed for each O atom so that the number of electrons lost equals the number of electrons gained.

SOLUTION:


Na•
$$\longrightarrow$$
 2Na+ $\stackrel{\cdot \circ}{\longrightarrow}$ 2Na• $\stackrel{\cdot \circ}{\longrightarrow}$ 1





UPLOADED BY AHMAD T JUNDI

Sample Problem 9.1





The formula is Na₂O





Figure 9.6 The exothermic formation of sodium bromide.







Periodic Trends in Lattice Energy

Lattice energy is the energy required to separate 1 mol of an ionic solid into gaseous ions.

Lattice energy is a measure of the strength of the ionic bond.

Coloumb's Law

Electrostatic energy ∞ char

charge A x charge B distance

Electrostatic energy ∞

cation charge x anion charge cation radius + anion radius

 $\propto \Delta \mathcal{H}_{ ext{lattice}}$



Periodic Trends in Lattice Energy

Lattice energy is affected by *ionic size* and *ionic charge*.


As ionic size *increases*, lattice energy *decreases*. Lattice energy therefore decreases down a group on the periodic table.

As ionic charge *increases*, lattice energy *increases*.





Figure 9.7 Trends in lattice energy.





Properties of Ionic Compounds

- Ionic compounds tend to be hard, rigid, and brittle, with high melting points.
- Ionic compounds do not conduct electricity in the solid state.
 - In the solid state, the ions are fixed in place in the lattice and do not move.
- Ionic compounds conduct electricity when melted or dissolved.
 - In the liquid state or in solution, the ions are free to move and carry a current.





Figure 9.8 Why ionic compounds crack.













Figure 9.9 Electrical conductance and ion mobility.



Solid ionic compound



Molten ionic
UPL®RBEPBY!Ahmad Jundi



lonic compound dissolved in water



Table 9.1 Melting and Boiling Points of Some Ionic Compounds

Compound	mp (°C)	bp (°C)
CsBr	636	1300
Nal	661	1304
MgCl ₂	714	1412
KBr	734	1435
CaCl ₂	782	>1600
NaCl	801	1413
LiF	845	1676
KF	858	1505
MgO	2852	3600





UPLOADED BY AHMAD T JUNDI Figure 9.10 Covalent bond formation in H₂.









Figure 9.11 Distribution of electron density in H_2 .



At some distance (bond length), attractions balance repulsions.

Electron density is high around and between the nuclei.





Bonding Pairs and Lone Pairs

Atoms share electrons to achieve a full outer level of electrons. The shared electrons are called a *shared pair* or *bonding pair*.

The shared pair is represented as a pair of dots or a line:

An outer-level electron pair that is not involved in bonding is called a *lone pair*, or *unshared pair*.





Properties of a Covalent Bond

The **bond order** is the number of electron pairs being shared by a given pair of atoms.

A single bond consists of one bonding pair and has a bond order of 1.

The **bond energy** (BE) is the energy needed to overcome the attraction between the nuclei and the shared electrons. The **stronger** the bond the **higher** the bond energy.

The **bond length** is the distance between the nuclei of the bonded atoms.





Trends in bond order, energy, and length

For a given pair of atoms, a *higher bond order* results in a *shorter bond length* and *higher bond energy*.

For a given pair of atoms, a shorter bond is a stronger bond.

Bond length *increases* down a group in the periodic table and *decreases* across the period.

Bond energy shows the opposite trend.





UPLOADED BY AHMAD T JUND!

Table 9.2 Average Bond Energies (kJ/mol) and Bond Lengths (pm)

Bond	Energy	Length	Bond	Energy	Length	Bond	Energy	Length	Bond	Energy	Length
Single Bonds											
H— H	432	74	N-H	391	101	Si—H	323	148	S-H	347	134
H— F	565	92	N-N	160	146	Si—Si	226	234	S-S	266	204
H—Cl	427	127	N-P	209	177	Si—O	368	161	S—F	327	158
H—Br	363	141	N-O	201	144	Si—S	226	210	S—Cl	271	201
H— I	295	161	N-F	272	139	Si—F	565	156	S—Br	218	225
			N—Cl	200	191	Si—Cl	381	204	S-I	~170	234
C-H	413	109	N—Br	243	214	Si—Br	310	216			
C-C	347	154	N-I	159	222	Si—I	234	240	F-F	159	143
C—Si	301	186							F—Cl	193	166
C-N	305	147	O-H	467	96	Р—Н	320	142	F—Br	212	178
C-O	358	143	O-P	351	160	P—Si	213	227	F—I	263	187
C-P	264	187	0 - 0	204	148	P-P	200	221	Cl—Cl	243	199
C-S	259	181	o-s	265	151	P—F	490	156	Cl—Br	215	214
C—F	453	133	O-F	190	142	P—Cl	331	204	Cl—I	208	243
C-C1	339	177	O—Cl	203	164	P—Br	272	222	Br—Br	193	228
C—Br	276	194	O—Br	234	172	P—I	184	246	Br—I	175	248
C—I	216	213	O—I	234	194				I—I	151	266
Multiple	Bonds										
C=C	614	134	N=N	418	122	$C \equiv C$	839	121	$N \equiv N$	945	110
C=N	615	127	N=0	607	120	$C \equiv N$	891	115	$N \equiv 0$	631	106
C=0	745	123	O_2	498	121	$C \equiv O$	1070	113			
(799 in CO	(2)									





Table 9.3 The Relation of Bond Order, Bond Length, and Bond Energy

Bond	Bond Order	Average Bond Length (pm)	Average Bond Energy (kJ/mol)
С—О	1	143	358
C=O	2	123	745
C≡O	3	113	1070
C-C	1	154	347
C = C	2	134	614
$C \equiv C$	3	121	839
N-N	1	146	160
N=N	2	122	418
$N \equiv N$	3	110	945





Figure 9.12

Bond length and covalent radius.


Internuclear distance (bond length)

Covalent radius


Internuclear distance (bond length)

Covalent radius


114 pm



72 pm




Internuclear distance (bond length)



Covalent radius

100 pm

Internuclear distance (bond length)



Covalent radius

133 pm





Sample Problem 9.2

Comparing Bond Length and Bond Strength

PROBLEM: Using the periodic table, but not Tables 9.2 or 9.3, rank the bonds in each set in order of *decreasing* bond length and decreasing bond strength:

PLAN: (a) S is singly bonded to three different halogen atoms, so the bond order is the same. Bond length increases and bond strength decreases as the atomic radius of the halogen increases.

(b) The same two atoms are bonded in each case, but the bond orders differ. Bond strength increases and bond length decreases as bond order increases.





Sample Problem 9.2

SOLUTION:

(a) Atomic size increases going down a group, so F < Cl < Br.

Bond length: S-Br > S-Cl > S-F Bond strength: S-F > S-Cl > S-Br

(b) By ranking the bond orders, we get

Bond length: C=O > C=O > C=OBond strength: C=O > C=O > C=O





Figure 9.13

Strong forces within molecules and weak forces between them.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.










Figure 9.14 Covalent bonds of network covalent solids: quartz and diamond.









Figure 9.15 The infrared (IR) spectra of 2-butanol (*green*) and diethyl ether (*red*).







Bond Energies and ΔH^{0}_{rxn}

The heat released or absorbed during a chemical change is due to differences between the bond energies of reactants and products.

$$\Delta H^{o}_{rxn} = \Sigma \Delta H^{o}_{reactant bonds broken} + \Sigma \Delta H^{o}_{product bonds formed}$$






Figure 9.16 Using bond energies to calculate ΔH^{o}_{rxn} for HF formation.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display. **ATOMS** $\Sigma \Delta H_{\text{reactant bonds broken}}^{\circ} =$ + 591 kJ $\Sigma \Delta H_{\text{product bonds formed}}^{\circ} =$ BE = 432 kJ/mol BE = 159 kJ/mol-1130 kJ $+\Sigma BE$ Enthalpy, H BE = 565 kJ/mol $-\Sigma BE$ **REACTANTS** ∆**H**° rxn **PRODUCTS**





Figure 9.17 Using bond energies to calculate ΔH^o_{rxn} for the combustion of methane.







Sample Problem 9.3

Using Bond Energies to Calculate ΔH^{o}_{rxn}

PROBLEM: Calculate ΔH_{rxn}^{\prime} for the chlorination of methane to form chloroform.

PLAN:

All the reactant bonds break, and all the product bonds form. Find the bond energies in Table 9.2 and substitute the two sums, with correct signs, into Equation 9.2.





UPLOADED BY AHMAD T JUNDI

Sample Problem 9.3

SOLUTION:

For bonds broken:

$$4 \times C-H = (4 \text{ mol})(413 \text{ kJ/mol}) = 1652 \text{ kJ}$$

 $3 \times CI-CI = (3 \text{ mol})(243 \text{ kJ/mol}) = 729 \text{ kJ}$

$$\Sigma \Delta H_{\text{bonds broken}}^{\prime} = 2381 \text{ kJ}$$

For bonds formed:

$$3 \times \text{C-Cl} = (3 \text{ mol})(-339 \text{ kJ/mol}) = -1017 \text{ kJ}$$

 $1 \times \text{C-H} = (1 \text{ mol})(-413 \text{ kJ/mol}) = -413 \text{ kJ}$
 $3 \times \text{H-Cl} = (3 \text{ mol})(-427 \text{ kJ/mol}) = -1281 \text{ kJ}$
 $\Sigma \Delta \mathcal{H}^{\circ}_{\text{bonds formed}} = -2711 \text{ kJ}$

$$\Delta \mathcal{H}_{\text{reaction}}^{\circ} = \Sigma \Delta \mathcal{H}_{\text{bonds broken}}^{\circ} + \Sigma \Delta \mathcal{H}_{\text{bonds formed}}^{\circ}$$

= 2381 kJ + (-2711 kJ) = -330 kJ





Electronegativity and Bond Polarity

A covalent bond in which the shared electron pair is not shared equally, but remains closer to one atom than the other, is a *polar covalent bond*.


The ability of an atom in a covalent bond to attract the shared electron pair is called its *electronegativity*.

Unequal sharing of electrons causes the more electronegative atom of the bond to be *partially* negative and the less electronegative atom to be *partially* positive.




Figure 9,418 D BY AHMAD BORDING between the models.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



Pure ionic




Polar covalent bonds are much more common than either pure ionic or pure covalent bonds.







Figure 9.19 The Pauling electronegativity (EN) scale.







Trends in Electronegativity

The most electronegative element is *fluorine*.

In general electronegativity *decreases* down a group as atomic size *increases*.

In general electronegativity *increases* across a period as atomic size *decreases*.

Nonmetals are *more* electronegative than metals.





Electronegativity and Oxidation Number

Electronegativities can be used to assign oxidation numbers:

- The more electronegative atom is assigned all the shared electrons.
- The less electronegative atom is assigned *none* of the shared electrons.
- Each atom in a bond is assigned all of its unshared electrons.
- O.N. = # of valence e⁻ (# of shared e⁻ + # of unshared e⁻)





Example: H—CI

Cl is more electronegative than H, so for Cl:

valence $e^{-} = 7$

shared $e^{-} = 2$

unshared $e^{-} = 6$

O.N.
$$= 7 - (2 + 6) = -1$$

H is less electronegative than Cl, so for H:

valence $e^{-}=1$

shared e⁻ = 0 (all shared e⁻ assigned to CI)

unshared $e^{-} = 0$


O.N.
$$= 1 - (0 + 0) = +1$$





Depicting Polar Bonds

The unequal sharing of electrons can be depicted by a polar arrow. The head of the arrow points to the *more electronegative element*.




A polar bond can also be marked using δ + and δ - symbols.

$$\delta + \delta - F$$





Figure 9.20 Electron density distributions in H_2 , F_2 , and HF.



In HF, the electron density shifts from H to F.

The H–F bond has partial ionic character.




Figure 9.21 ΔEN ranges for classifying the partial ionic character of bonds.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display. 3.3 Mostly ionic δ + δ -Polar covalent Mostly Nonpolar UP**coxadeot**BY:Ahmad Jundi





Figure 9.22 Percent ionic character as a function of ΔEN.




Determining Bond Polarity from EN Values

- **PROBLEM:** (a) Use a polar arrow to indicate the polarity of each bond: N–H, F–N, I–CI.
 - **(b)** Rank the following bonds in order of increasing polarity: H–N, H–O, H–C.
- **PLAN:** (a) We use Figure 9.21 to find the EN values for each element. The polar arrow points toward the more electronegative element.
 - (b) The greater the Δ EN between the atoms, the more polar the bond.

SOLUTION: (a) The EN values are:

$$N = 3.0$$
, $H = 2.1$; $F = 4.0$; $I = 2.5$, $CI = 3.0$







(b) The EN values are:

$$N = 3.0, H = 2.1; O = 3.5; C = 2.5$$

$$\Delta$$
EN for H-N = 3.0 - 2.1 = 0.9

$$\Delta$$
EN for H $-$ O = 3.5 $-$ 2.1 = 1.4

$$\Delta$$
EN for H–C = 2.5 – 2.1 = 0.4


H-C < H-N < H-O






Figure 9.23 Properties of the Period 3 chlorides.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.






As Δ EN decreases, melting point and electrical conductivity decrease because the bond type changes from ionic to polar covalent to nonpolar covalent.







Chapter 12

Lecture Outline

See separate *Image PowerPoint* slides for all figures and tables pre-inserted into PowerPoint without notes.



Chapter 12

Intermolecular Forces:

Liquids, Solids, and Phase Changes





Intermolecular Forces: Liquids, Solids, and Phase Changes

- 12.1 An Overview of Physical States and Phase Changes
- 12.2 Quantitative Aspects of Phase Changes
- 12.3 Types of Intermolecular Forces
- 12.4 Properties of the Liquid State
- 12.5 The Uniqueness of Water
- 12.6 The Solid State: Structure, Properties, and Bonding



Phases of Matter

Each physical state of matter is a *phase*, a physically distinct, homogeneous part of a system.

The properties of each phase are determined by the balance between the *potential* and *kinetic* energy of the particles.

The *potential* energy, in the form of *attractive forces*, tends to draw particles together.

The *kinetic* energy associated with movement tends to disperse particles.



Attractive Forces


Intramolecular or *bonding* forces are found *within* a molecule. The *chemical* behavior of each phase of matter is the same because the same basic particle is present in each case.

H₂O molecules are present whether the substance is in the solid, liquid, or gas phase.

Intermolecular or *nonbonding* forces are found *between* molecules. The *physical* behavior of each phase of matter is different because the *strength* of these forces differs from state to state.



UPLOADED BY AHMAD T JUNDI



Intramolecular Forces -

Force which keeps molecule together, i.e., bonds.

Intermolecular Forces -

Attractive force between molecules.

Responsible for keeping matter in solid or liquid phase.

UPLOADED BY:Ahmad Jundi



Table 12.1 A Macroscopic Comparison of Gases, Liquids, and Solids

State	Shape and Volume	Compressibility	Ability to Flow
Gas	Conforms to shape and volume of container	High	High
Liquid	Conforms to shape of container; volume limited by surface	Very low	Moderate
Solid	Maintains its own shape and volume	Almost none	Almost none



Kinetic Molecular View of the Three States

Attractive Forces	VS.
Kinetic Energy	

Properties

Gas

Attractive forces are weak relative to kinetic energy.

Particles are far apart. A gas has no fixed shape or volume.

Liquid

Attractive forces are stronger because particles have less kinetic energy.

A liquid can flow and change shape, but has a fixed volume.

Solid


Attractions dominate motion. Particles are fixed in place relative to each other.

A solid has a fixed shape and volume.



Phase Changes

sublimation



<u>exothermic</u>

Deposition (Desublimation)



UPLOADED BY: Ahmad Jundi



Figure 12.1 Heats of vaporization and fusion for several common substances.







Figure 12.2 Phase changes and their enthalpy changes.





Quantitative Aspects of Phase Changes

Within a phase, heat flow is accompanied by a *change* in *temperature*, since the average E_k of the particles changes.


 $q = \text{(amount)} \times \text{(heat capacity)} \times \Delta T$

During a phase change, heat flow occurs at **constant temperature**, as the average distance between particles changes.

 $q = (amount)(\Delta H of phase change)$



Figure 12.3 A cooling curve for the conversion of gaseous water to ice.







WDENTS-HUB.com

Finding the Heat of a Phase Change Depicted by Molecular Scenes

PROBLEM: The scenes below represent a phase change of water. Select data from the previous text discussion to find the heat (in kJ) released or absorbed when 24.3 g of H₂O undergoes this change.






PLAN: The scenes show a disorderly, condensed phase at 85.0°C changing to separate particles at 117°C. A liquid is becoming a gas, so the scene shows vaporization. We must identify the number of stages in the process and calculate the heat absorbed in each.

SOLUTION:

There are 3 stages involved in this process:

- 1) heating of the liquid to its boiling point
- 2) the phase change from liquid to gas
- 3) heating the gas to the final temperature



Copyright © The McGraw-Hill Companies, Inc. Permission required for

mol H₂O = 24.3 g H₂O
$$\times \frac{1 \text{ mol H}_2\text{O}}{18.02 \text{ g H}_2\text{O}} = 1.35 \text{ mol H}_2\text{O}$$

For Stage 1:

$$q = n \times C_{\text{water}(1)} \times \Delta T$$

Molar heat capacity
$$(C) = q/moles \times \Delta T$$
 $(J/mol \cdot ^{\circ}C)$

$$C$$
 of H₂O(I) = 4.184 J/g.K x 18.02 g/mol = 75.40 J/mol.K

$$q = n \times C_{\text{water}(\Lambda)} \times \Delta T$$

$$= (1.35 \text{ mol})(75.4 \text{ J/mol} \cdot ^{\circ}\text{C})(100. - 85.0 ^{\circ}\text{C})$$

$$= 1527 J = 1.53 kJ$$



For Stage 2:

$$q = n(\Delta H^{\prime}_{vap}) = (1.35 \text{ mol})(40.7 \text{ kJ/mol}) = 54.9 \text{ kJ}$$

For Stage 3:

$$q = n \times C_{\text{water}(g)} \times \Delta T$$

= (1.35 mol)(33.1 J/mol·°C)(117 – 100.°C)
= 759.6 J = 0.760 kJ

$$C$$
 of H₂O(g) = 1.843 J/g.K

$$q_{\text{total}} = 1.53 + 54.9 + 0.760 \text{ kJ} = 57.2 \text{ kJ}$$



Figure 12.4 Liquid-gas equilibrium.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Vacuum

Time

Liquid

Liquid


A Molecules in the liquid vaporize.

B Molecules vaporize and condense at the same rate.

In a closed flask, the system reaches a state of *dynamic equilibrium*, where molecules are leaving and entering the liquid at the *same rate*.




Figure 12.4 continued



The *vapor pressure* is the pressure exerted by the vapor on the liquid. The pressure increases until equilibrium is reached; **at equilibrium the pressure is constant**.



Figure 12.5 The effect of temperature on the distribution of molecular speeds.



Factors affecting Vapor Pressure

As temperature *increases*, the fraction of molecules with enough energy to enter the vapor phase *increases*, and the vapor pressure *increases*.


higher $T \longrightarrow$ higher P

The *weaker* the intermolecular forces, the *more* easily particles enter the vapor phase, and the *higher* the vapor pressure.

weaker forces \longrightarrow higher P



Figure 12.6 Vapor pressure as a function of temperature and intermolecular forces.



Vapor pressure *increases* as temperature *increases*.

Vapor pressure *decreases* as the strength of the intermolecular forces *increases*.



The Clausius-Clapeyron Equation

This equation relates vapor pressure to temperature.


$$\ln P = \frac{-\Delta H_{\text{vap}}}{R} \left(\frac{1}{T}\right) + C$$

The two-point form is used when the vapor pressures at two different temperatures are known.

$$\ln \frac{P_2}{P_1} = \frac{-\Delta H_{\text{vap}}}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$$



Figure 12.7 Linear plots of the relationship between vapor pressure and temperature.



$$\ln P = \frac{-\Delta H_{\text{vap}}}{R} \left(\frac{1}{T}\right) + C$$

slope =
$$\frac{-\Delta H_{\text{vap}}}{R}$$

Applying the Clausius-Clapeyron Equation

PROBLEM: The vapor pressure of ethanol is 115 torr at 34.9°C. If ΔH_{vap} of ethanol is 40.5 kJ/mol, calculate the temperature (in °C) when the vapor pressure is 760 torr.

PLAN: We are given 4 of the 5 variables in the Clausius-Clapeyron equation, so we substitute these into the equation and solve for T_2 . Tvalues must be converted to K.

SOLUTION:

In
$$\frac{P_2}{P_1} = -\frac{\Delta H_{\text{vap}}}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$$
 $T_1 = 34.9^{\circ}\text{C} + 273.15 = 308.0 \text{ K}$

$$\ln \frac{760 \text{ torr}}{115 \text{ torr}} = -\frac{40.5 \times 10^3 \text{ J/mol}}{8.314 \text{ J/mol} \cdot \text{K}} \left(\frac{1}{T_2} - \frac{1}{308.0 \text{ K}} \right)$$



$$T_2 = 350$$
. K $- 273.15 = 77$ °C

Vapor Pressure and Boiling Point

The *boiling point* of a liquid is the temperature at which the vapor pressure equals the external pressure.

The *normal boiling point* of a substance is observed at standard atmospheric pressure or 760 torr.

As the external pressure on a liquid *increases*, the boiling point *increases*.



Figure 12.8 Phase diagram for CO₂.







Figure 12.8 Phase diagram for H_2O .



The solid-liquid line slants to the *left* for H₂O, because the solid is *less* dense than the liquid. Water expands on freezing.



The Nature of Intermolecular Forces


Intermolecular forces arise from the attraction between molecules with partial charges, or between ions and molecules.

Intermolecular forces are relatively *weak* compared to bonding forces because they involve smaller charges that are farther apart.



Figure 12.9A Covalent and van der Waals radii.

The van der Waals *distance* is the distance between two *nonbonded* atoms in *adjacent* molecules.



The van der Waals *radius* is one-half the closest distance between the nuclei of two *nonbonded* atoms. The VDW radius is *always larger* than the covalent radius.

UPLOADED BY: Ahmad Jundi



Figure 12.9B Periodic trends in covalent and van der Waals radii.






Table 12.2 Comparison of Bonding and Nonbonding (Intermolecular) Forces

Force	Model	Basis of Attraction	Energy (kJ/mol)	Example
Bonding Ionic	+ - + - +	Cation–anion	400–4000	NaCl
Covalent	•••	Nuclei–shared e ⁻ pair	150–1100	Н—Н
Metallic	+ + + + + + + + + + + + + + + + + + + +	Cations–delocalized electrons	75–1000	Fe



Table 12.2 Comparison of Bonding and Nonbonding (Intermolecular) Forces (continued)

Force	Model	Basis of Attraction	Energy (kJ/mol)	Example
Nonbonding (Int	termolecular)			
Ion-dipole	+	Ion charge— dipole charge	40-600	Na+···O
H bond	δ- δ+ δ- -A-H·····B-	Polar bond to H- dipole charge (high EN of N, O,	10–40 F)	:Ö— н···:Ö— н Н Н
Dipole-dipole	·····	Dipole charges	5-25	I—CI…I—CI
Ion-induced dipole	+	Ion charge— polarizable e [—] cloud	3–15	Fe ²⁺ ···O ₂
Dipole-induced dipole		Dipole charge— polarizable e cloud	2–10	H—CI····CI—CI
Dispersion (London)		Polarizable e clouds	0.05-40	F—F···F—F



dipole-dipole forces.





Figure 12.11 Dipole moment and boiling point.



The Hydrogen Bond


Hydrogen bonding is possible for molecules that have a *hydrogen atom* covalently bonded to a small, highly electronegative atom with *lone electron pairs*, specifically *N, O, or F*.

An intermolecular *hydrogen bond* is the attraction between the *H atom* of one molecule and a *lone pair of the N, O, or F atom* of another molecule.



ST2DB6S-HUB.com

Figure 12.12 Hydrogen bonding and boiling point.



Drawing Hydrogen Bonds Between Molecules of a Substance

PROBLEM: Which of the following substances exhibits H bonding? For any that do, draw the H bonds between two of its molecules.

(a)
$$C_2H_6$$

PLAN: If the molecule does not contain N, O, or F it cannot form H bonds. If it contains any of these atoms covalently bonded to H, we draw two molecules in the pattern –B:----H–A.


SOLUTION:

(a) C_2H_6 has no N, O, or F, so no H-bonds can form.



(b) CH₃OH contains a covalent bond between O and H. It can form H bonds between its molecules:

(c) CH₃C — NH₂ can form H bonds at two sites:





Polarizability and Induced Dipoles

A nearby electric field can *induce* a distortion in the electron cloud of an atom, ion, or molecule.

- For a *nonpolar* molecule, this induces a *temporary dipole moment*.
- For a *polar* molecule, the field *enhances* the existing dipole moment.

The *polarizability* of a particle is the ease with which its electron cloud is distorted.



Trends in Polarizability

Smaller particles are **less polarizable** than larger ones because their electrons are held more tightly.

Polarizability *increases down a group* because atomic size increases and larger electron clouds distort more easily.

Polarizability *decreases across a period* because of increasing Z_{eff} .

Cations are smaller than their parent atoms and less polarizable; anions show the opposite trend.



Figure 12.13 Dispersion forces among nonpolar particles.






A. When atoms are far apart they do not influence one other.



B. When atoms are close together, the instantaneous dipole in one atom induces a dipole in the other.



C. The process occurs throughout the sample.



Dispersion (London) Forces

Dispersion forces or **London forces** arises when an **instantaneous dipole** in one particle **induces** a dipole in another, resulting in an attraction between them.

Dispersion forces exist between *all particles*, increasing the energy of attraction in all matter.

Dispersion forces are *stronger* for *more polarizable* particles.

In general, larger particles experience stronger dispersion forces than smaller ones.



Figure 12.14 Molar mass and trends in boiling point.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Dispersion forces are stronger for larger, more polarizable particles.

Polarizability correlates closely with molar mass for similar particles.








Figure 12.15 Molecular shape, intermolecular contact, and boiling point.



 CH_3 CH_2 CH_2 CH_2 CH_3 n-Pentane, bp = 36.1°C

There are more points at which dispersion forces act.

There are fewer points at which dispersion forces act.




Neopentane, bp = 9.5° C



UPLOADED BY:Ahmad Jundi

Figure 12.16 Determining the intermolecular forces in a sample.



DISPERSION FORCES ALSO PRESENT



Predicting the Types of Intermolecular Forces

PROBLEM: For each pair of substances, identify the key bonding and/or intermolecular force(s), and predict which one of the pair has the higher boiling point:

(a) $MgCl_2$ or PCl_3 (b) CH_3NH_2 or CH_3F (c) CH_3OH or CH_3CH_2OH

PLAN: We examine the formulas and structures for key differences between the members of each pair: Are ions present? Are molecules polar or nonpolar? Is N, O, or F bonded to H? Do molecular compounds have different masses or shapes?



Remember that:

- Bonding forces are stronger than nonbonding (intermolecular) forces.
- Hydrogen bonding is a strong type of dipole-dipole force.
- Dispersion forces are decisive when the difference is molar mass or molecular shape.

SOLUTION:

(a) MgCl₂ consists of Mg²⁺ and Cl⁻ ions held together by **ionic bonding forces**; PCl₃ consists of polar molecules, so intermolecular **dipole-dipole** forces are present. The ionic bonding forces in MgCl₂ are stronger than the dipole-dipole forces in PCl₃.

MgCl₂ has a higher boiling point than PCl₃.



(b) CH₃NH₂ and CH₃F both consist of **polar molecules** of about the same molar mass. CH₃NH₂ has covalent N-H bonds, so it can form **H bonds** between its molecules. CH₃F contains a C-F bond but no H-F bond, so **dipole-dipole** forces occur but not H bonds.

CH₃NH₂ has a higher boiling point than CH₃F.




(c) CH₃OH and CH₃CH₂OH are both **polar molecules** and both contain a covalent O-H bond. Both can therefore form **H bonds**.

CH₃CH₂OH has a larger molar mass than CH₃OH and its **dispersion forces** are therefore stronger.

CH₃CH₂OH has a higher boiling point than CH₃OH.




(d) Hexane and 2,2-dimethylbutane are both **nonpolar molecules** and therefore experience **dispersion forces** as their only intermolecular force. They have equal molar masses but different molecular shapes.



Cylindrical hexane molecules make more intermolecular contact than the more compact 2,2-dimethylbutane molecules.

Hexane has a higher boiling point than 2,2-dimethylbutane.



CHECK:

The actual boiling points show our predictions are correct:

- (a) MgCl₂ (1412°C) and PCl₃ (76°C)
- (b) CH_3NH_2 (-6.3°C) and CH_3F (-78.4°C)
- (c) CH₃OH (64,7°C) and CH₃CH₂OH (78.5°C)
- (d) Hexane (69°C) and 2,2-dimethylbutane (49.7°C)

Remember that *dispersion forces are always present*, but in (a) and (b) they are much less significant than the other forces that occur.




Figure 12.17 The molecular basis of surface tension.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

A surface molecule experiences a *net attraction downward*. This causes a liquid surface to have the smallest area possible.

An interior molecule is attracted by others on all sides.




Surface tension is the energy required to increase the surface area of a liquid. The *stronger* the forces between the particles the *higher* the surface tension.

Table	12.3	Surface	Tension ar	nd Forces	Between	Particles

Substance	Formula	Surface Tension (J/m²) at 20°C	Major Force(s)
Diethyl ether	CH ₃ CH ₂ OCH ₂ CH ₃	1.7x10 ⁻²	Dipole-dipole; dispersion
Ethanol	CH ₃ CH ₂ OH	2.3x10 ⁻²	H bonding
Butanol	CH ₃ CH ₂ CH ₂ CH ₂ OH	H 2.5x10 ⁻²	H bonding; dispersion
Water	H ₂ O	7.3x10 ⁻²	H bonding
Mercury	Hg	48x10 ⁻²	Metallic bonding



Figure 12.18 Capillary action and the shape of the water or mercury meniscus in glass.



A. Water displays a concave B. Mercury d meniscus.

UPLOADED BY: Ahmad Meniscus.

ST2D54TS-HUB.com

B. Mercury displays a convex _{mad}നുക്വiscus. 54

Table 12.4 Viscosity of Water at Several Temperatures

Viscosity is resistance of a fluid to flow.


Temperature (°C)	Viscosity (N·s/m²)*
20	1.00x10 ⁻³
40	0.65x10 ⁻³
60	0.47x10 ⁻³
80	0.35x10 ⁻³

^{*}The units of viscosity are Newton-seconds per square meter.



The uniqueness not water

H-bonding ability of water.




Each H₂O molecule can form *four* H bonds to other molecules, resulting in a tetrahedral arrangement.




- 1- Solvent properties: its results of its <u>polarity</u> and <u>hydrogen bond</u> ability. dissolve ionic compounds through ion-dipole forces
- 2- Thermal properties: water has high specific heat capacity (water has so many strong hydrogen bonding.
- 3- surface properties of water: water has high surface tension and high capillary
- 4- Density of solid and liquid water: ice has hexagonal structure, this organization explain the negative slope of solid liquid line in phase diagram. The large space within ice give the Solid state lower density than the liquid.



Figure 12.20 The hexagonal structure of ice.





© Scott Camazine/Photo Researchers, Inc.

Ice has an open structure due to H bonding. Ice is therefore *less* dense than liquid water.

