



Patient package insert
Prescription Only medicine
Carefully read the entire leaflet before starting use
SITAGLU 50 & 100 FILM COATED TABLETS

COMPOSITION:

Sitaglu 50 Film Coated Tablets: Each tablet contains 50 mg Sitagliptin (as monohydrate phosphate).
Sitaglu 100 Film Coated Tablets: Each tablet contains 100 mg Sitagliptin (as monohydrate phosphate).

THERAPEUTIC ACTIVITY: Is a DDP-4 enzyme inhibitors (Dipeptidyl peptidase-4 enzyme inhibitors).

INDICATIONS: It is indicated as an adjunct to diet and exercise to improve glycemic control (lowers blood sugar levels) in adults' patients with type 2 diabetes mellitus.

WHEN SHOULD THE PREPARATION NOT BE USED?

- If you have type 1 diabetes mellitus.
- If you have diabetic ketoacidosis (increased ketones in the blood or urine: diabetes complication which includes high blood sugar, rapid weight loss, nausea or vomiting).
- If you are allergic (sensitive) to the active ingredient. Symptoms of a serious allergic reaction to Sitaglu may include: rash, raised red patches on your skin (hives), or swelling of the face, lips, tongue, and throat that may cause difficulty in breathing or swallowing.

DO NOT USE THIS MEDICINE WITHOUT CONSULTING A PHYSICIAN: If you have or have had in the past pancreatitis (inflammation of the pancreas), gallstones, alcoholism or very high blood triglyceride levels. These medical conditions can increase your chance of getting pancreatitis (see "Side Effects" section). If you have kidney problems. If you have any other medical conditions. If you are pregnant or plan to become pregnant talk with your doctor about the best way to control your blood sugar while you are pregnant. It is not recommended to take JANUVIA during pregnancy or if you are breast-feeding or plan to breast-feed.

WARNINGS & PRECAUTIONS: Do not use this medicine frequently or for long periods of time without consulting your doctor. If you are sensitive to any type of food or medicine, consult your doctor prior to commencing treatment.

DRIVING AND USING MACHINE: This medicine has no known influence on the ability to drive and use machines. However, dizziness and drowsiness could occur, which may affect your ability to drive or use machines. Taking Sitaglu in combination with medicines called sulfonylurea or with insulin, can cause hypoglycemia, which may affect your ability to drive and use machines.

DRUG INTERACTIONS: Tell your doctor or pharmacist if you are taking, have recently taken, or might take any other medicines including non-prescription medicines and nutritional supplements, you should inform the attending doctor or pharmacist.

SIDE EFFECTS: In addition to the desired effect of the medicine, adverse reactions may occur during the course of taking this medicine, for example: