

سوال نمبر 3


Principles of Physics (10th edition)

Phy 132

CH 21: Coulomb's Law

Problems: 3, 6, 13, 31, 35, 37

P₃: In Fig 21.11, the particles have charges $q_1 = -q_2 = 300 \text{ nC}$ and $q_3 = -q_4 = 200 \text{ nC}$ and distance $a = 5.0 \text{ cm}$. What are the (a) magnitude and (b) angle (relative to +x direction) of the net force on particle 3?




Sol: $a = 5 \text{ cm} = 5 \times 10^{-2} \text{ m}$, $q_1 = -q_2 = 300 \text{ nC} = 300 \times 10^{-9} \text{ C}$, $q_3 = -q_4 = 200 \text{ nC} = 200 \times 10^{-9} \text{ C}$

$$\vec{F}_{13} = \frac{k q_1 q_3}{a^2}$$

$$= \frac{9 \times 10^9 \times 300 \times 10^{-9} \times 200 \times 10^{-9}}{(5 \times 10^{-2})^2}$$

$$= \frac{5.4 \times 10^{-4}}{25 \times 10^{-4}}$$

$$\vec{F}_{13} = 0.216 \text{ N} \quad (-y)$$




(2)

kclw

$$\vec{F}_{23} = \frac{k q_2 q_3}{r^2} = \frac{9 \times 10^9 \times 300 \times 10^{-9} \times 200 \times 10^{-9}}{(\sqrt{50} \times 10^{-2})^2}$$

$$= 0.108 \text{ N}$$




$$\vec{F}_{43} = \frac{k q_4 q_3}{a^2} = \frac{9 \times 10^9 \times 200 \times 10^{-9} \times 200 \times 10^{-9}}{(5 \times 10^{-2})^2}$$

$$= 0.144 \text{ N (+x)}$$

$$\Sigma F_x = F_4 + F_2 \cos 45$$

$$= 0.144 + 0.108 \times \frac{1}{\sqrt{2}}$$

$$= 0.22036 \text{ N (+x)}$$



$$\Sigma F_y = F_2 \sin 45 - F_1$$

$$= -0.1396$$

$$= 0.1396 \text{ N (-y)}$$

$$|F_{\text{net}}| = \sqrt{F_x^2 + F_y^2}$$

$$= \sqrt{(0.2203)^2 + (0.1396)^2}$$


$$= 0.2608 \text{ N}$$

$$\tan \theta = \frac{\Sigma F_y}{\Sigma F_x} \Rightarrow \theta = \tan^{-1} \left(\frac{\Sigma F_y}{\Sigma F_x} \right)$$

$$\theta = \tan^{-1} \left(\frac{0.1396}{0.22036} \right) = \tan^{-1}(0.633)$$

$$= -32 \text{ clock wise}$$

$$\text{or } 360 - 32 = 328 \text{ counter-clockwise}$$




(3)


سوال ٣

P₆: Three particles are fixed on an x axis. Particle 1 of charge q_1 is at $x = -a$, and particle 2 of charge q_2 is at $x = +a$. If their net electrostatic force on particle 3 of charge $+Q$ is to be zero, what must be the ratio q_1/q_2 when particle 3 is at (a) $x = +0.750a$ and (b) $x = +1.50a$?

Sol:



a)



$$\Sigma F = 0 \quad (\text{net electrostatic force equal zero})$$

$$F_1 = F_2$$

$$\frac{k q_1 Q}{(a + 0.75a)^2} = \frac{k q_2 Q}{(a - 0.75a)^2}$$


$$\frac{q_1}{(1.75a)^2} = \frac{q_2}{(0.25a)^2}$$

$$\frac{q_1}{q_2} = \frac{(1.75a)^2}{(0.25a)^2} = 49$$

4

سوال 4

6) دو نقطه بار q_1 و q_2 در فاصله $2.5a$ و $0.5a$ از یکدیگر قرار دارند. اگر نیروهای الکتریکی بین آنها متعادل باشد، نسبت بارها را بیابید.



$$F_1 = F_2$$

$$\frac{k q_1}{(2.5a)^2} = \frac{k q_2}{(0.5a)^2}$$

$$\frac{q_1}{(2.5a)^2} = \frac{q_2}{(0.5a)^2}$$

$$\frac{q_1}{q_2} = \frac{(2.5)^2}{(0.5)^2} \frac{a^2}{a^2}$$


$$\frac{q_1}{q_2} = 25$$

Since $q_1 \oplus$ $q_2 \ominus$ then $\frac{q_1}{q_2} = -25$

P13: A nonconducting spherical shell with an inner radius of 4.0 cm and outer radius of 5.0 cm, has charge spread nonuniformly through its volume between its inner and outer surfaces.

The volume charge density ρ is the charge per unit volume with the unit coulomb per cubic meter. For this shell $\rho = br$, where r is the distance in meters from the center of the shell and $b = 3.0 \text{ nC/m}^3$. What is the net charge in the shell?


sol: $b = 3 \times 10^{-6} \text{ C/m}^3$



$$\rho = \frac{Q}{V}$$

(5)

سارہ جاتا



$$Q = \rho V$$

$$dQ = \rho dV$$

but $V = \text{Area} \times \text{height or thickness}$

$$V = A \cdot L$$

but in spherical shell

$$L = dr$$

$$A = 4\pi r^2$$

Surface area

$$dQ = \rho A dr$$

$$dQ = \rho 4\pi r^2 dr$$

$$\int dQ = \int_{r_1}^{r_2} \rho 4\pi r^2 dr, \quad \rho = b/r$$

$$Q = \int_{r_1}^{r_2} \frac{b}{r} 4\pi r^2 dr$$

$$Q = 4\pi b \int_{r_1}^{r_2} r dr$$

$$Q = 4\pi b \frac{r^2}{2} \Big|_{r_1}^{r_2}$$

$$Q = 2\pi b r^2 \Big|_{r_1}^{r_2}$$

$$Q = 2\pi b (r_2^2 - r_1^2)$$

$$Q = 2\pi \times 3 \times 10^{-6} ((5 \times 10^{-2})^2 - (4 \times 10^{-2})^2)$$


$$Q = 2\pi \times 3 \times 10^{-6} \times 9 \times 10^{-4}$$

$$Q = 1.7 \times 10^{-8} \text{ C}$$

(6)

سوال 31

P31: In Fig 21-21, particles 1 and 2 of charge $q_1 = q_2 = +4e$ are on y-axis at distance $d = 17.0 \text{ cm}$ from the origin. Particle 3 of charge $q_3 = +8e$ is moved gradually along the x-axis from $x=0$ to $x = +5.0 \text{ m}$. At what values of x will the magnitude of electrostatic force on the third particles be (a) minimum (b) maximum? what are the (c) minimum and (d) maximum magnitudes?



حل المسألة

$$q_1 = q_2 = +4e \Rightarrow d = 17 \text{ cm}$$

$$q_3 = +8e \Rightarrow x = 0 \rightarrow 5 \text{ m}$$


sol:

* The magnitude of the electrostatic force between any two particles

$$F = k \frac{|q_1| |q_2|}{r^2} \quad (1)$$

θ : angle between F and x -axis

$$\cos \theta = \frac{x}{\sqrt{x^2 + d^2}} \quad (2)$$



* due to symmetry, there is no y component to the net force on the third particle.

$$\Sigma F_x = F_1 \cos \theta + F_2 \cos \theta = 2F_1 \cos \theta \quad , \quad \Sigma F_y = F_1 \sin \theta - F_2 \sin \theta = 0$$

$$F_{\text{net}} = F_1 + F_2 = 2F \cos \theta = \frac{2K(4e)(8e)}{(\sqrt{x^2 + d^2})^2} \frac{x}{\sqrt{x^2 + d^2}} = \frac{64Ke^2 x}{(x^2 + d^2)^{3/2}}$$

(7)

سارہ خان

لا یبار

$$F_{\text{net}} = \frac{64ke^2 x}{(x^2 + d^2)^{3/2}} \quad \text{--- (3)}$$

$F_{\text{net}} = 0$ at $x = 0$ according to eq (3)

(a) So the minimum value for x is zero

(b) taking the derivative of eq (3) and equating it to zero to find the max value

$$\Rightarrow x = \frac{d}{\sqrt{2}} = \frac{17 \text{ cm}}{\sqrt{2}} = 12 \text{ cm}$$

$$(c) \quad F_{\text{net}}' = \frac{64ke^2 x}{(x^2 + d^2)^{3/2}} = \frac{64ke^2 (0)}{(0 + d^2)^{3/2}} = 0$$

$$(d) \quad F_{\text{net}} = \frac{64ke^2 x}{(x^2 + d^2)^{3/2}}$$

$$= \frac{64 \times 9 \times 10^9 \times (1.6 \times 10^{-19})^2 \times (0.12)}{(0.12)^2 + (0.17)^2)^{3/2}}$$

$$= \frac{1.769 \times 10^{-27}}{9.010 \times 10^{-3}}$$

$$= 1.96 \times 10^{-25} \text{ N}$$

$$\approx 2.0 \times 10^{-25} \text{ N}$$

derivative for part (b)

$$F = \frac{64ekx}{(d^2+x^2)^{3/2}}$$

$$F = 64e^2kx (d^2+x^2)^{-3/2}$$

$$\frac{dF}{dx} = 64e^2k \left[(d^2+x^2)^{-3/2} + x \left(-\frac{3}{2} \right) (d^2+x^2)^{-5/2} 2x \right]$$

$$= 64e^2k \left[\frac{1}{(d^2+x^2)^{3/2}} - \frac{3x^2}{(d^2+x^2)^{5/2}} \right]$$

$$= 64e^2k \left[\frac{d^2+x^2-3x^2}{(d^2+x^2)^{5/2}} \right]$$

$$\frac{dF}{dx} = 64e^2k \left[\frac{d^2-2x^2}{(d^2+x^2)^{5/2}} \right]$$

$$0 = 64e^2k \left[\frac{d^2-2x^2}{(d^2+x^2)^{5/2}} \right]$$

$$d^2-2x^2=0 \Rightarrow d^2=2x^2$$

$$x = \pm \frac{d}{\sqrt{2}}$$

سأردنا


P35: In Fig 21-24, three charged particles lie on an x axis. Particles 1 and 2 are fixed in place. Particle 3 is free to move, but the net electrostatic force on it from particles 1 and 2 happens to be zero. If $2.0 L_{23} = L_{12}$, what is the ratio q_1/q_2 ?

لما أن عمله القوى = صفر لنقطة
تقع خارج الشحنتين

مع الشحنتين متساوية في المقدار
وتكونا لهما أحزاب للاحتراف
أي $q_1 > q_2$ إذن نتوقع


$$\left| \frac{q_1}{q_2} \right| > 1$$

والا $L_{12} < L_{23}$



دول:

$$F_{\text{net}} = F_{13} - F_{23}$$



$$\text{but } F_{\text{net}} = 0$$

$$\ominus q_2 < \oplus q_1 \quad \text{لأن } q_1 > q_2$$

$$\Rightarrow F_{13} = F_{23}$$

$$\frac{k q_1 q_3}{(L_{12} + L_{23})^2} = \frac{k q_2 q_3}{(L_{23})^2}$$

$$\frac{q_1}{(L_{12} + L_{23})^2} = \frac{q_2}{L_{23}^2} \quad \text{but } L_{12} = 2 L_{23}$$

$$\frac{q_1}{(2 L_{23} + L_{23})^2} = \frac{q_2}{(L_{23})^2}$$

$$\frac{q_1}{9 L_{23}^2} = \frac{q_2}{L_{23}^2}$$

9


ساده

$$\frac{q_1}{q_2} = \frac{q_1 L_{23}^2}{L_{13}^2}$$


$$\Rightarrow \left| \frac{q_1}{q_2} \right| = 9 \quad \Rightarrow \frac{q_1}{q_2} = -9 \quad \text{because } q_1 (+) \quad q_2 (-)$$

P37: If the charge Q initially on a tiny sphere, a portion q is to be transferred to a second, nearby sphere. Both spheres can be treated as particles and are fixed with a certain separation. (a) For what value of q/Q will the electrostatic force between the two spheres be maximized? What are (b) smaller and (c) larger values of q/Q that give a force magnitude that is 75% of that maximum?

sol: Before After



Q



zero



$Q-q$



q

$$|F| = \frac{k q_1 q_2}{r^2} = \frac{k (Q-q) q}{r^2}$$

$$\Rightarrow |F| = \frac{k (Qq - q^2)}{r^2}$$

$$a) \frac{dF}{dq} = \frac{k}{r^2} \frac{d}{dq} (Qq - q^2) = \frac{k}{r^2} (Q - 2q)$$

سواء

$$\text{maximized } \frac{dF}{dq} = 0$$

$$\Rightarrow \frac{k}{r^2} (q - 2q) = 0, \quad \frac{k}{r^2} \text{ const}$$

$$q - 2q = 0$$

$$q = 2q$$

$$\boxed{\frac{q}{q} = \frac{1}{2}}$$

$$b) F_{\max} \text{ at } \frac{q}{q} = \frac{1}{2} \Rightarrow \text{at } q = \frac{q}{2}$$

$$\text{sub } q = q/2 \text{ in } F = \frac{k(qq - q^2)}{r^2}$$

$$F_{\max} = \frac{k \left(\frac{q \times q}{2} - \left(\frac{q}{2} \right)^2 \right)}{r^2}$$

$$= \frac{k \left(\frac{q^2}{2} - \frac{q^2}{4} \right)}{r^2}$$

$$F_{\max} = \frac{kq^2}{4r^2}$$

$$\Rightarrow 75\% \text{ of } F_{\max} \text{ is } \frac{3}{4} \text{ of } F_{\max}$$

$$\Rightarrow 75\% \text{ of } F_{\max} = \frac{3}{4} \times \frac{kq^2}{4r^2} = \frac{3}{16} \frac{kq^2}{r^2}$$

* to find the smaller value and larger value of q/q of 75 %

$$|F| = 75\% F_{\max}$$

$$\frac{k(qq - q^2)}{r^2} = \frac{3}{16} \frac{kq^2}{r^2}$$

ساده

$$Q^2 - Q^2 = \frac{3}{16} Q^2$$

$$Q^2 - Q^2 + \frac{3}{16} Q^2 = 0$$

$$a=1, b=-Q, c=\frac{3}{16} Q^2 \quad \text{معادله درجه دوم}$$

$$Q = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{+Q \pm \sqrt{Q^2 - 4 \times \frac{3}{16} Q^2}}{2}$$

$$= \frac{Q \pm \sqrt{Q^2 - \frac{12}{16} Q^2}}{2}$$

$$= \frac{Q \pm \sqrt{\frac{1}{4} Q^2}}{2}$$

$$Q = \frac{Q \pm \frac{Q}{2}}{2}$$

$$Q = \frac{Q + \frac{Q}{2}}{2} = \frac{3}{4} Q$$

$$Q = \frac{Q - \frac{Q}{2}}{2} = \frac{Q}{2}$$

so b) smaller value $\Rightarrow Q = Q/2$

c) larger value $\Rightarrow Q = \frac{3}{4} Q$