

# Trigonometric Integrals

33

Exp

$$\int \sec^2 x \tan x \, dx$$

$$u = \tan x$$

$$du = \sec^2 x \, dx$$

$$\int u \, du$$

$$= \frac{u^2}{2} + C$$

$$= \frac{\tan^2 x}{2} + C$$

45

$$\int 4 \tan^3 x \, dx$$

$n=3$  odd

$$\int 4 \tan^2 x \tan x \, dx$$

$$1 + \tan^2 x = \sec^2 x$$

$$\tan^2 x = \sec^2 x - 1$$

$$\int 4 (\sec^2 x - 1) \tan x \, dx$$

$$u = \tan x$$

$$du = \sec^2 x \, dx$$

$$4 \int \sec^2 x \tan x \, dx - 4 \int \tan x \, dx$$

33

$$\frac{4}{2} \frac{\tan^2 x}{2} + 4 \int \frac{-\sin x}{\cos x} dx$$

$$= 2 \tan^2 x + 4 \int \frac{-\sin x}{\cos x} dx$$

$$= 2 \tan^2 x + 4 \ln |\cos x| + C$$

(67)  $\int x \sin^2 x dx$

$$\int x \left( \frac{1 - \cos 2x}{2} \right) dx$$

$$\int x \left( \frac{1}{2} - \frac{\cos 2x}{2} \right) dx$$

$$\int \frac{x}{2} dx - \int \frac{x}{2} \cos 2x dx$$

$$\frac{1}{2} \frac{x^2}{2} - \left[ \frac{x}{4} \sin 2x + \frac{1}{8} \cos 2x \right]$$

$$\frac{x^2}{4} - \frac{x}{4} \sin 2x - \frac{1}{8} \cos 2x + C$$

$$\sin^2 x = \frac{1 - \cos 2x}{2}$$

$$\cos^2 x = \frac{1 + \cos 2x}{2}$$

$$\cos 2x = 1 - 2 \sin^2 x = 2 \cos^2 x - 1$$

$\frac{x}{2}$	$\cos 2x$
$\downarrow +$	$\frac{1}{2} \sin 2x$
$\frac{1}{2}$	$\downarrow -$
$0$	$-\frac{1}{4} \cos 2x$

$$\frac{x}{4} - \frac{1}{4} \sin 4x - \frac{8}{3} \cos^3 x \dots$$

$n=3$  odd

5)  $\int \sin^3 x \, dx$

$$\begin{aligned} \sin^2 x + \cos^2 x &= 1 \\ \sin^2 x &= 1 - \cos^2 x \end{aligned}$$

$$\int \sin^2 x \sin x \, dx$$

$$u = \cos x$$

$$\int (1 - \cos^2 x) \sin x \, dx$$

$$du = -\sin x \, dx$$

$$-\int (1 - u^2) \, du$$

$$\int (u^2 - 1) \, du = \frac{u^3}{3} - u + C$$

$$= \frac{\cos^3 x}{3} - \cos x + C$$

22)  $\int_0^{\frac{\pi}{2}} \sin^2 \theta \cos^3 \theta \, d\theta$

Annotations: 2 (sin), 3 (cos), odd, opp

$$\int_0^{\frac{\pi}{2}} \sin^2 \theta \cos^2 \theta \cos \theta \, d\theta$$

Annotations:  $\cos^2 \theta$  in a cloud,  $u^2$  pointing to  $\sin^2 \theta$

Q. How to integrate  $\int \sin^m x \cos^n x \, dx$ ?

Look for odd #

$$u = \sin 2\theta$$

$$du = 2 \cos 2\theta \, d\theta$$

$$du = \cos 2\theta \, d\theta$$

$$\int_0^{\frac{\pi}{2}} \sin^2 \theta (1 - \sin^2 \theta) \cos 2\theta \, d\theta$$

$$\int_0^{\frac{\pi}{2}} \sin^2 \theta (1 - \sin^2 \theta) \cos \theta d\theta$$

$$\int_0^1 u^2 (1 - u^2) \frac{du}{2}$$

$$\frac{du}{2} = \cos \theta d\theta$$

when  $\theta = 0 \Rightarrow$   
 $u = \sin 0 = 0$

when  $\theta = \frac{\pi}{2} \Rightarrow$   
 $u = \sin \pi = 0$

Exp  $\int \sin^2 x \cos^3 x dx$

$\downarrow$  (odd)       $\downarrow$  (odd)

$$\int \sin^2 x \sin x \cos^3 x dx$$

$\downarrow$   
 $u^3$

$$u = \cos x$$

$$du = -\sin x dx$$

$$-du = \sin x dx$$

$$\int (1 - \cos^2 x) \cos^3 x \sin x dx$$

$$= \int (1 - u^2) u^3 du$$

$$\int (u^2 - 1) u^3 du = \int (u^5 - u^3) du$$

$$= \frac{u^6}{6} - \frac{u^4}{4} + C$$

-- 6

$$= \frac{\cos^6 x}{6} - \frac{\cos^4 x}{4} + C$$

Exp  $\int 16 \sin^2 x \cos^2 x dx$

n and m  
 \ /  
 even  
 ↓  
 Use

$$\int 16 \left( \frac{1 - \cos 2x}{2} \right) \left( \frac{1 + \cos 2x}{2} \right) dx$$

$$\checkmark \cos^2 x = \frac{1 + \cos 2x}{2}$$

$$\checkmark \sin^2 x = \frac{1 - \cos 2x}{2}$$

$$4 \int (1 - \cos 2x)(1 + \cos 2x) dx$$

$$4 \int (1 - \cos^2 2x) dx$$

$$x^2 - y^2 = (x-y)(x+y)$$

$x = 1$   
 $y = \cos 2x$

$$\int 4 dx - 4 \int \cos^2 2x dx$$

$$\checkmark \cos^2 x = \frac{1 + \cos 2x}{2}$$

$$\checkmark \cos^2(2x) = \frac{1 + \cos 4x}{2}$$

$$4x - \frac{4}{2} \int \left( \frac{1 + \cos 4x}{2} \right) dx$$

$$4x - 2 \int (1 + \cos 4x) dx$$

$$4x - 2 \left( x + \frac{\sin 4x}{4} \right) + C$$

$$4x - 2x - \frac{1}{2} \sin 4x + C$$

$$2x - \frac{1}{2} \sin 4x + C$$

$$\int 16 \sin^2 x \cos^2 x dx$$

$$\sin 2x = 2 \sin x \cos x$$

$$\sin^2 2x = 4 \sin^2 x \cos^2 x$$

$$4 \int 4 \sin^2 x \cos^2 x dx$$

$$4 \int \sin^2 2x dx$$

$$\sin^2 2x = \frac{1 - \cos 4x}{2}$$

$$\frac{4}{2} \int \frac{1 - \cos 4x}{2} dx$$

$$2 \int (1 - \cos 4x) dx$$

$$2 \left( x - \frac{\sin 4x}{4} \right) + C$$

$$2x - \frac{1}{2} \sin 4x + C$$

$$\int \sin mx \sin nx \, dx = \int \frac{1}{2} [\cos(m-n)x - \cos(m+n)x] \, dx$$

$$\bullet \int \cos mx \cos nx \, dx = \int \frac{1}{2} [\cos(m-n)x + \cos(m+n)x] \, dx$$

$$\int \sin mx \cos nx \, dx = \int \frac{1}{2} [\sin(m-n)x + \sin(m+n)x] \, dx$$

(3)

---

Exp (51)

$$\int \sin^{\overset{m=3}{3}x} \cos^{\overset{n=2}{2}x} \, dx = \int \frac{1}{2} [\sin(3-2)x + \sin(3+2)x] \, dx$$
$$= \int \frac{1}{2} [\sin x + \sin 5x] \, dx$$
$$= \frac{1}{2} \left[ -\cos x - \frac{\cos 5x}{5} \right] + C$$
$$= -\frac{1}{2} \cos x - \frac{1}{10} \cos 5x + C$$


Exp  $\int \cos 3x \cos 4x dx = \frac{1}{2} \int [\cos(3-4)x + \cos(3+4)x] dx$

$$= \frac{1}{2} \int [\cos(-x) + \cos(7x)] dx$$

$$= \frac{1}{2} \int [\cos x + \cos 7x] dx$$

$$= \frac{1}{2} \left( \sin x + \frac{\sin 7x}{7} \right) + C$$

$$= \frac{\sin x}{2} + \frac{\sin 7x}{14} + C$$



## Estimating Square Roots

$$\cos^2 x = 1 - \sin^2 x$$

Exp  $\int_0^{\pi} \sqrt{1 - \sin^2 x} dx$


$$\sqrt{\cos^2 x} = \cancel{\cos x}$$

$$= |\cos x|$$

$$\int_0^{\pi} \sqrt{\cos^2 x} dx$$

$$\int_0^{\pi} |\cos x| dx$$

$$= \int_0^{\frac{\pi}{2}} \cos x dx$$



$$\int_0^{\pi} \cos x dx = 0$$




$$= 2 \int_0^{\frac{\pi}{2}} \cos x \, dx$$

$$= 2 \sin x \Big|_0^{\frac{\pi}{2}}$$

$$= 2 \left[ \sin \frac{\pi}{2} - \sin 0 \right]$$

$$= 2 [1 - 0] = 2$$

$$\int_0^{\pi} \cos x \, dx = 0$$



or

$$\int_0^{\pi} |\cos x| \, dx = \int_0^{\frac{\pi}{2}} \cos x \, dx + \int_{\frac{\pi}{2}}^{\pi} -\cos x \, dx$$

$$|\cos x| = \begin{cases} \cos x & \text{if } 0 \leq x \leq \frac{\pi}{2} \\ -\cos x & \text{if } \frac{\pi}{2} \leq x \leq \pi \end{cases}$$

$$= \sin x \Big|_0^{\frac{\pi}{2}} - \sin x \Big|_{\frac{\pi}{2}}^{\pi}$$

$$= \sin \frac{\pi}{2} - \sin 0 - \left( \sin \pi - \sin \frac{\pi}{2} \right)$$

$$= 1 - 0 - (0 - 1)$$

$$= 1 + 1$$

$$= 2$$

Exp

$$\int \sec^4 x \, dx = \int \underbrace{\sec^2 x}_{1 + \tan^2 x} \sec^2 x \, dx$$

$$= \int (1 + \tan^2 x) \sec^2 x \, dx$$

$$= \int (1 + u^2) \, du$$

$$= u + \frac{u^3}{3} + C$$

$$= \tan x + \frac{1}{3} \tan^3 x + C$$

$$u = \tan x$$

$$du = \sec^2 x \, dx$$

$$\int \sin^m x \cos^n x \, dx$$

✓ if  $m$  odd  $\Rightarrow \sin^m x = \sin^{2k+1} x = \sin^{2k} x \sin x$   
 and use  $\sin^2 x = 1 - \cos^2 x$

✓ if  $n$  odd  $\Rightarrow \cos^n x = \cos^{2k+1} x = \cos^{2k} x \cos x$   
 and use  $\cos^2 x = 1 - \sin^2 x$

$$\rightarrow \dots \cos^2 x = \frac{1 + \cos 2x}{2}$$

---

if  $m$  and  $n$  both even  $\implies$  use

$$\cos^2 x = \frac{1 + \cos 2x}{2}$$

$$\sin^2 x = \frac{1 - \cos 2x}{2}$$

---