

MVT : f cont. on $[a, b]$
 f diff on (a, b)


$\Rightarrow \exists$ at least one number $c \in (a, b)$
 s.t $f'(c) = \frac{f(b) - f(a)}{b - a}$

Exp $f(x) = \begin{cases} -x^2 + 3x + a & , x = 0 \\ mx + b & , 0 < x < 1 \\ mx + b & , 1 \leq x \leq 2 \end{cases}$

Assume f satisfies MVT on $[0, 2]$

Find constants a, m, b

* f cont. on $[0, 2]$ $\Rightarrow f$ cont. at 0 ✓
 $\Rightarrow f$ cont. at 1 ✓✓



$$3 = \lim_{x \rightarrow 0^+} (-x^2 + 3x + a)$$



$$3 = a$$

f cont. at $x=1 \Rightarrow f(1) = \lim_{x \rightarrow 1} f(x)$

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^-} f(x)$$

$$\lim_{x \rightarrow 1^+} (mx + b) = \lim_{x \rightarrow 1^-} (-x^2 + 3x + a)$$

$$m + b = -(1)^2 + 3(1) + a$$

$$m + b = -1 + 3 + 3$$

$$m + b = 5 \Rightarrow m = 1 \Rightarrow 1 + b = 5$$

$$b = 4$$

f diff on $(0, 2)$

$$f'(x) = \begin{cases} 0 & x = 0 \\ -2x + 3 & 0 < x < 1 \\ m & 1 < x < 2 \end{cases}$$

f cont on $[-1, 0]$ } Bolzano قانون
 $f(-1) f(0) < 0$

\exists at least one root $c \in [a, b]$
 such that $f(c) = 0$


To show c is unique $f = x^3 + 3x + 1$

$$f'(x) = 3x^2 + 3$$

$$f' \quad + \quad + \quad + \quad + \quad +$$

f is always increasing

f cross x -axis only one time at $x=c$



Bolzano

f cont. on $[a, b]$

$$a = \frac{1}{2}$$


$$b = 3$$

$$[a, b] = \left[\frac{1}{2}, 3\right]$$

$$\left\{ \begin{array}{l} f \text{ cont. on } \left[\frac{1}{2}, 3\right] \\ f\left(\frac{1}{2}\right) f(3) < 0 \end{array} \right. \Rightarrow$$

$$\begin{array}{l} f\left(\frac{1}{2}\right) < 0 \\ f(3) > 0 \end{array}$$

\exists root.



Exp

$$f(x) = x\sqrt{1-x^2}$$

① Domain $\Rightarrow D(f)$

$$1 - x^2 \geq 0$$

$$1 \geq x^2$$

$$\sqrt{1} \geq \sqrt{x^2}$$

$$1 \geq |x|$$

$$|x| \leq 1$$

$$\boxed{-1 \leq x \leq 1}$$

$$D(f) = \underline{\underline{[-1, 1]}}$$

② CP's

$$f(x) = x\sqrt{1-x^2}$$

$$= x(1-x^2)^{\frac{1}{2}}$$

$$= \frac{1}{2}(-1/x) + \sqrt{1-x^2}$$

$$\begin{aligned}
 f'(x) &= x \cdot \frac{1}{2} (1-x^2)^{-\frac{1}{2}} (-2x) + \frac{\sqrt{1-x^2}}{1-x^2} \\
 &= \frac{-x^2}{\sqrt{1-x^2}} + \frac{\sqrt{1-x^2} \cdot \sqrt{1-x^2}}{\sqrt{1-x^2}} \\
 &= \frac{-x^2 + 1 - x^2}{\sqrt{1-x^2}}
 \end{aligned}$$

$$f' = \frac{1-2x^2}{\sqrt{1-x^2}} = 0$$

$$1-2x^2=0$$

$$\Rightarrow 2x^2=1$$

$$x^2 = \frac{1}{2}$$

$$\text{in domain} \leftarrow x = \pm \frac{1}{\sqrt{2}}$$

$$x = \pm \frac{1}{\sqrt{2}} \in [-1, 1]$$

$x = \pm \frac{1}{\sqrt{2}}$ critical point

$$\left(\frac{1}{\sqrt{2}}, f\left(\frac{1}{\sqrt{2}}\right)\right) = \left(\frac{1}{\sqrt{2}}, \frac{1}{2}\right)$$

$$\left(-\frac{1}{\sqrt{2}}, f\left(-\frac{1}{\sqrt{2}}\right)\right) = \left(-\frac{1}{\sqrt{2}}, -\frac{1}{2}\right)$$

$$f(x) = \sqrt{1-x^2}$$


$$f(x) = x \sqrt{1-x^2}$$

$$f\left(\frac{1}{\sqrt{2}}\right) = \frac{1}{\sqrt{2}} \sqrt{1-\frac{1}{2}} = \frac{1}{\sqrt{2}} \sqrt{\frac{1}{2}} = \frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}} = \frac{1}{2}$$

$$f\left(-\frac{1}{\sqrt{2}}\right) = -\frac{1}{\sqrt{2}} \sqrt{1-\frac{1}{2}} = -\frac{1}{\sqrt{2}} \sqrt{\frac{1}{2}} = -\frac{1}{\sqrt{2}} \frac{1}{\sqrt{2}} = -\frac{1}{2}$$

③ intervals of \uparrow and \downarrow for f

$$f' = \frac{1-2x^2}{\sqrt{1-x^2}}$$



$f \uparrow$ on $\left[-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$

$f \downarrow$ on $\left[-1, -\frac{1}{\sqrt{2}}\right)$ and $\left[\frac{1}{\sqrt{2}}, 1\right]$

④ EV's

check CP's : $\left(\frac{1}{\sqrt{2}}, \frac{1}{2}\right)$, $\left(-\frac{1}{\sqrt{2}}, -\frac{1}{2}\right)$

check Endpoints $(-1, 0)$, $(1, 0)$

$$f(x) = x \sqrt{1-x^2} \quad \text{on } [-1, 1]$$

f has L. Max of $f\left(\frac{1}{\sqrt{2}}\right) = \frac{1}{2}$

$$f(x) = x\sqrt{1-x^2} \text{ on } [-1, 1]$$

$$f(-1) = -1\sqrt{1-(-1)^2} = (-1)(0) = 0$$

$$f(1) = (1)\sqrt{1-(1)^2} = (1)(0) = 0$$

$$f\left(\frac{1}{\sqrt{2}}\right) = \frac{1}{2}$$


f has L. Min of

$$f\left(-\frac{1}{\sqrt{2}}\right) = -\frac{1}{2}$$

(5) sketch f

$$f(x) = x\sqrt{1-x^2}$$

keypoint
(0,0)



$$(6) R(f) = \left[-\frac{1}{2}, \frac{1}{2}\right]$$