



COMP 233 Discrete Mathematics

---

# Chapter 5

## Sequences and Mathematical Induction



# Outline

---

- Sequences:
  - Explicit Formulas;
  - Summation Notation;
  - Sequences in Computer Programming;
- Proof by Mathematical Induction (I and II)
  - Proving sum of integers and geometric sequences
  - Proving a Divisibility Property and Inequality
  - Proving a Property of a Sequence

# Sequences

**Idea:** Think of a sequence as a set of elements written in a row:


$$\begin{array}{ll} a_1, a_2, a_3, \dots, a_n & \text{finite sequence} \\ a_1, a_2, a_3, \dots, a_n, \dots & \text{infinite sequence} \end{array}$$

Each individual element  $a_k$  is called a term.

The  $k$  in  $a_k$  is called a subscript or index

## Observe patterns

Determine the number of points in the 4<sup>th</sup> and 5<sup>th</sup> figure



Determine the next 2 terms of the sequence

4, 8, 16, 32, 64

Induce the formula that could be used to determine any term in the sequence

# Finding Terms of Sequences Given by Explicit Formulas

Define sequences  $a_1, a_2, a_3, \dots$  and  $b_2, b_3, b_4, \dots$  by the following explicit formulas:

$$a_k = \frac{k}{k+1} \text{ for some integers } k \geq 1$$

$$b_i = \frac{i-1}{i} \text{ for some integers } i \geq 2$$

Compute the first five terms of both sequences.

Compute the first six terms of the sequence  $c_0, c_1, c_2, \dots$  defined as follows:  $c_j = (-1)^j$  for all integers  $j \geq 0$ .



# Finding Explicit Formula to Fit Given Initial Terms

Find an explicit formula for a sequence that has the following initial terms:

$$1, -\frac{1}{4}, \frac{1}{9}, -\frac{1}{16}, \frac{1}{25}, -\frac{1}{36}, \dots$$

$$a_k = \frac{-1^{k+1}}{k^2} \quad \text{for all integers } k \geq 1.$$

OR

$$a_k = \frac{-1^k}{(k+1)^2} \quad \text{for all integers } k \geq 0.$$



# Exercises

---

**Example:** Find an explicit formula for a sequence that has the following initial terms:

$$\frac{1}{3}, -\frac{2}{4}, \frac{3}{5}, -\frac{4}{6}, \frac{5}{7}, -\frac{6}{8}, \dots$$

**Solutions:** The sequence satisfies the formulas

for all integers  $n \geq 0$ ,

$$a_n = (-1)^n \frac{n+1}{n+3}$$

for all integers  $n \geq 1$ ,

$$a_n = (-1)^{n-1} \frac{n}{n+2}$$



# Summation Notation

Suppose  $a_1, a_2, a_3, \dots, a_n$  are real numbers. The “summation from  $i$  equals 1 to  $n$  of  $a$ -sub- $i$ ” is

$$\sum_{i=1}^n a_i = a_1 + a_2 + a_3 + \cdots + a_n.$$

## • Definition

If  $m$  and  $n$  are integers and  $m \leq n$ , the symbol  $\sum_{k=m}^n a_k$ , read the **summation from  $k$  equals  $m$  to  $n$  of  $a$ -sub- $k$** , is the sum of all the terms  $a_m, a_{m+1}, a_{m+2}, \dots, a_n$ . We say that  $a_m + a_{m+1} + a_{m+2} + \dots + a_n$  is the **expanded form** of the sum, and we write

$$\sum_{k=m}^n a_k = a_m + a_{m+1} + a_{m+2} + \cdots + a_n.$$

We call  $k$  the **index** of the summation,  $m$  the **lower limit** of the summation, and  $n$  the **upper limit** of the summation.



# Exercises

---

**Ex:** Use summation notation to write the following sum:

$$\frac{1}{3} - \frac{2}{4} + \frac{3}{5} - \frac{4}{6} + \frac{5}{7} - \frac{6}{8}$$

**Solution:** By the example on the previous slide, we can write:

$$\frac{1}{3} - \frac{2}{4} + \frac{3}{5} - \frac{4}{6} + \frac{5}{7} - \frac{6}{8} = \sum_{n=0}^5 (-1)^n \left( \frac{n+1}{n+3} \right).$$

or:

$$\frac{1}{3} - \frac{2}{4} + \frac{3}{5} - \frac{4}{6} + \frac{5}{7} - \frac{6}{8} = \sum_{n=1}^6 (-1)^{n+1} \left( \frac{n}{n+2} \right).$$



# Exercises

Let  $a_1 = -2$ ,  $a_2 = -1$ ,  $a_3 = 0$ ,  $a_4 = 1$ , and  $a_5 = 2$ .  
Compute the following:

a.  $\sum_{k=1}^5 a_k$

b.  $\sum_{k=2}^2 a_k$

c.  $\sum_{k=1}^2 a_{2k}$

## Example 5.1.4 Computing Summations

Let  $a_1 = -2$ ,  $a_2 = -1$ ,  $a_3 = 0$ ,  $a_4 = 1$ , and  $a_5 = 2$ . Compute the following:

a.  $\sum_{k=1}^5 a_k$

b.  $\sum_{k=2}^2 a_k$

c.  $\sum_{k=1}^2 a_{2k}$

### Solution

a.  $\sum_{k=1}^5 a_k = a_1 + a_2 + a_3 + a_4 + a_5 = (-2) + (-1) + 0 + 1 + 2 = 0$

b.  $\sum_{k=2}^2 a_k = a_2 = -1$

c.  $\sum_{k=1}^2 a_{2k} = a_{2 \cdot 1} + a_{2 \cdot 2} = a_2 + a_4 = -1 + 1 = 0$



# Summation to Expanded Form

Write the following summation in expanded form:

$$\sum_{i=0}^n \frac{(-1)^i}{i+1}$$

$$\begin{aligned}\sum_{i=0}^n \frac{(-1)^i}{i+1} &= \frac{(-1)^0}{0+1} + \frac{(-1)^1}{1+1} + \frac{(-1)^2}{2+1} + \frac{(-1)^3}{3+1} + \dots + \frac{(-1)^n}{n+1} \\ &= \frac{1}{1} + \frac{(-1)}{2} + \frac{1}{3} + \frac{(-1)}{4} + \dots + \frac{(-1)^n}{n+1} \\ &= 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{(-1)^n}{n+1}\end{aligned}$$



# Expanded Form to Summation

Express the following using summation notation:

$$\frac{1}{n} + \frac{2}{n+1} + \frac{3}{n+2} + \dots + \frac{n+1}{2n}$$

$$\frac{1}{n} + \frac{2}{n+1} + \frac{3}{n+2} + \dots + \frac{n+1}{2n} = \sum_{k=0}^n \frac{k+1}{n+k}$$

# Separating Off a Final Term and Adding On a Final Term $n$

Rewrite  $\sum_{i=1}^{n+1} \frac{1}{i^2}$  by separating off the final term.

$$\sum_{i=1}^{n+1} \frac{1}{i^2} = \sum_{i=1}^n \frac{1}{i^2} + \frac{1}{(n+1)^2}$$

Write  $\sum_{k=0}^n 2^k + 2^{n+1}$  as a single summation.


$$\sum_{k=0}^n 2^k + 2^{n+1} = \sum_{k=0}^{n+1} 2^k$$



# Telescoping Sum

$$\sum_{k=1}^n \left( \frac{1}{k} - \frac{1}{k+1} \right)$$

$$= \left( \frac{1}{1} - \frac{1}{2} \right) + \left( \frac{1}{2} - \frac{1}{3} \right) + \left( \frac{1}{3} - \frac{1}{4} \right) + \cdots + \left( \frac{1}{n-1} - \frac{1}{n} \right) + \left( \frac{1}{n} - \frac{1}{n+1} \right)$$

$$= 1 - \frac{1}{n+1}.$$




# Product Notation

## • Definition

If  $m$  and  $n$  are integers and  $m \leq n$ , the symbol  $\prod_{k=m}^n a_k$ , read the **product from  $k$  equals  $m$  to  $n$  of  $a$ -sub- $k$** , is the product of all the terms  $a_m, a_{m+1}, a_{m+2}, \dots, a_n$ .

We write

$$\prod_{k=m}^n a_k = a_m \cdot a_{m+1} \cdot a_{m+2} \cdots a_n.$$

A recursive definition for the product notation is the following: If  $m$  is any integer, then

$$\prod_{k=m}^m a_k = a_m \quad \text{and} \quad \prod_{k=m}^n a_k = \left( \prod_{k=m}^{n-1} a_k \right) \cdot a_n \quad \text{for all integers } n > m.$$



# Computing Products

---

- Compute the following products:

a.  $\prod_{k=1}^5 k$

- $= 1*2*3*4*5=120$

b.  $\prod_{k=1}^1 \frac{k}{k+1}$

- $= 1/2$



# Properties of Summations

## Theorem 5.1.1

If  $a_m, a_{m+1}, a_{m+2}, \dots$  and  $b_m, b_{m+1}, b_{m+2}, \dots$  are sequences of real numbers and  $c$  is any real number, then the following equations hold for any integer  $n \geq m$ :

$$1. \sum_{k=m}^n a_k + \sum_{k=m}^n b_k = \sum_{k=m}^n (a_k + b_k)$$

$$2. c \cdot \sum_{k=m}^n a_k = \sum_{k=m}^n c \cdot a_k \quad \text{generalized distributive law}$$

$$3. \left( \prod_{k=m}^n a_k \right) \cdot \left( \prod_{k=m}^n b_k \right) = \prod_{k=m}^n (a_k \cdot b_k).$$



Let  $a_k = k + 1$  and  $b_k = k - 1$  for all integers  $k$ . Write each of the following expressions as a single summation or product:

a.  $\sum_{k=m}^n a_k + 2 \cdot \sum_{k=m}^n b_k$       b.  $\left( \prod_{k=m}^n a_k \right) \cdot \left( \prod_{k=m}^n b_k \right)$

**Solution**

a.  $\sum_{k=m}^n a_k + 2 \cdot \sum_{k=m}^n b_k = \sum_{k=m}^n (k + 1) + 2 \cdot \sum_{k=m}^n (k - 1)$  by substitution

$$= \sum_{k=m}^n (k + 1) + \sum_{k=m}^n 2 \cdot (k - 1)$$

by Theorem 5.1.1 (2)

$$= \sum_{k=m}^n ((k + 1) + 2 \cdot (k - 1))$$

by Theorem 5.1.1 (1)

$$= \sum_{k=m}^n (3k - 1)$$

by algebraic simplification

b.  $\left( \prod_{k=m}^n a_k \right) \cdot \left( \prod_{k=m}^n b_k \right) = \left( \prod_{k=m}^n (k + 1) \right) \cdot \left( \prod_{k=m}^n (k - 1) \right)$  by substitution

$$= \prod_{k=m}^n (k + 1) \cdot (k - 1)$$

by Theorem 5.1.1 (3)

$$= \prod_{k=m}^n (k^2 - 1)$$

by algebraic simplification ■



# Change of Variable

---

**Example:** Transform  $\sum_{k=1}^n k^n$  by making the change of variable  $j = k - 1$ .

When  $k = 1$ , then  $j = 1 - 1 = 0$

When  $k = n$ , then  $j = n - 1$

$j = k - 1 \Rightarrow k = j + 1$  Thus  $k^n = (j + 1)^n$

So:  $\sum_{k=1}^n k^n = \sum_{j=0}^{n-1} (j+1)^n$

# Exercises

Transform the following summation by making the specified change of variable.

$$\sum_{k=0}^6 \frac{1}{k+1} \quad \text{Change variable } j = k+1$$

**For** (k=0; k<=6; k++)  
Sum = Sum + 1/(k+1)

$$\sum_{j=1}^7 \frac{1}{j} = \sum_{k=1}^7 \frac{1}{k}.$$

$$\sum_{k=0}^6 \frac{1}{k+1} = \sum_{k=1}^7 \frac{1}{k}$$

**For** (k=1; k<=7; k++)  
Sum = Sum + 1/(k)

# Exercises

Transform the following summation by making the specified change of variable.

$$\sum_{k=1}^{n+1} \frac{k}{n+k}$$

*Change of variable:  $j = k - 1$*

$$\sum_{j=0}^n \frac{j+1}{n+(j+1)} = \sum_{k=0}^n \frac{k+1}{n+(k+1)}$$

$$\sum_{k=1}^{n+1} \frac{k}{n+k} = \sum_{k=0}^n \frac{k+1}{n+(k+1)}$$

**For** (k=1; k<=n+1; k++)  
Sum = Sum + k/(n+k)


**For** (k=0; k<=n; k++)  
Sum = Sum + (k+1)/(n+k+1)



# Sequences in Computer Programming


- What is the difference

1. <b>for</b> $i := 1$ <b>to</b> $n$	2. <b>for</b> $j := 0$ <b>to</b> $n - 1$	3. <b>for</b> $k := 2$ <b>to</b> $n + 1$
<b>print</b> $a[i]$	<b>print</b> $a[j + 1]$	<b>print</b> $a[k - 1]$
<b>next</b> $i$	<b>next</b> $j$	<b>next</b> $k$



- Computing the sum

$s := a[1]$	$s := 0$
<b>for</b> $k := 2$ <b>to</b> $n$	<b>for</b> $k := 1$ <b>to</b> $n$
$s := s + a[k]$	$s := s + a[k]$
<b>next</b> $k$	<b>next</b> $k$



---

$$\sum_{k=1}^{n+1} \frac{k}{n+k} = \sum_{k=0}^n \frac{k+1}{n+(k+1)}$$

**Sum=0**

**For (k=0; k<=n; k++)**

**Sum = Sum + (k+1)/(n+k+1)**



# Factorial !

---

- **Definition**

For each positive integer  $n$ , the quantity  **$n$  factorial** denoted  $n!$ , is defined to be the product of all the integers from 1 to  $n$ :

$$n! = n \cdot (n - 1) \cdots 3 \cdot 2 \cdot 1.$$

**Zero factorial**, denoted  $0!$ , is defined to be 1:

$$0! = 1.$$

### Example 5.1.16 Computing with Factorials

Simplify the following expressions:

a.  $\frac{8!}{7!}$     b.  $\frac{5!}{2! \cdot 3!}$     c.  $\frac{1}{2! \cdot 4!} + \frac{1}{3! \cdot 3!}$     d.  $\frac{(n+1)!}{n!}$     e.  $\frac{n!}{(n-3)!}$

**Solution**

a.  $\frac{8!}{7!} = \frac{8 \cdot 7!}{7!} = 8$

b.  $\frac{5!}{2! \cdot 3!} = \frac{5 \cdot 4 \cdot \cancel{3!}}{2! \cdot \cancel{3!}} = \frac{5 \cdot 4}{2 \cdot 1} = 10$

c.  $\frac{1}{2! \cdot 4!} + \frac{1}{3! \cdot 3!} = \frac{1}{2! \cdot 4!} \cdot \frac{3}{3} + \frac{1}{3! \cdot 3!} \cdot \frac{4}{4}$  *by multiplying each numerator and denominator by just what is necessary to obtain a common denominator*  
 $= \frac{3}{3 \cdot 2! \cdot 4!} + \frac{4}{3! \cdot 4 \cdot 3!}$  *by rearranging factors*  
 $= \frac{3}{3! \cdot 4!} + \frac{4}{3! \cdot 4!}$  *because  $3 \cdot 2! = 3!$  and  $4 \cdot 3! = 4!$*   
 $= \frac{7}{3! \cdot 4!}$  *by the rule for adding fractions with a common denominator*  
 $= \frac{7}{144}$

d.  $\frac{(n+1)!}{n!} = \frac{(n+1) \cdot \cancel{n!}}{\cancel{n!}} = n+1$

e.  $\frac{n!}{(n-3)!} = \frac{n \cdot (n-1) \cdot (n-2) \cdot \cancel{(n-3)!}}{\cancel{(n-3)!}} = n \cdot (n-1) \cdot (n-2)$   
 $= n^3 - 3n^2 + 2n$





# $n$ choose $r$

## • Definition

Let  $n$  and  $r$  be integers with  $0 \leq r \leq n$ . The symbol

$$\binom{n}{r}$$

is read “ $n$  choose  $r$ ” and represents the number of subsets of size  $r$  that can be chosen from a set with  $n$  elements.

Observe that the definition implies that  $\binom{n}{r}$  will always be an integer because it is a number of subsets. In Section 9.5 we will explore many uses of  $n$  choose  $r$  for solving problems involving counting, and we will prove the following computational formula:

## • Formula for Computing $\binom{n}{r}$

For all integers  $n$  and  $r$  with  $0 \leq r \leq n$ ,

$$\binom{n}{r} = \frac{n!}{r!(n-r)!}$$



### Example 5.1.17 Computing $\binom{n}{r}$ by Hand

Use the formula for computing  $\binom{n}{r}$  to evaluate the following expressions:

a.  $\binom{8}{5}$

b.  $\binom{4}{0}$

c.  $\binom{n+1}{n}$

#### Solution

$$\begin{aligned} \text{a. } \binom{8}{5} &= \frac{8!}{5!(8-5)!} \\ &= \frac{8 \cdot 7 \cdot \cancel{6} \cdot \cancel{5} \cdot \cancel{4} \cdot \cancel{3} \cdot \cancel{2} \cdot 1}{(\cancel{5} \cdot \cancel{4} \cdot \cancel{3} \cdot \cancel{2} \cdot 1) \cdot (\cancel{3} \cdot \cancel{2} \cdot 1)} \\ &= 56. \end{aligned}$$

always cancel common factors  
before multiplying

$$\text{b. } \binom{4}{4} = \frac{4!}{4!(4-4)!} = \frac{4!}{4!0!} = \frac{\cancel{4} \cdot \cancel{3} \cdot \cancel{2} \cdot 1}{(\cancel{4} \cdot \cancel{3} \cdot \cancel{2} \cdot 1)(1)} = 1$$

The fact that  $0! = 1$  makes this formula computable. It gives the correct value because a set of size 4 has exactly one subset of size 4, namely itself.

$$\text{c. } \binom{n+1}{n} = \frac{(n+1)!}{n!((n+1)-n)!} = \frac{(n+1)!}{n!1!} = \frac{(n+1) \cdot n!}{n!} = n+1 \quad \blacksquare$$



5.2

---

# Mathematical Induction



# Mathematical Induction: A Way to Prove Such Formulas and Other Things

- Given an integer variable  $n$ , we can consider a variety of properties  $P(n)$  that might be true or false for various values of  $n$ . For instance, we could consider

$$P(n): 1 + 3 + 5 + 7 + \dots + (2n - 1) = n^2$$

$$P(n): 4^n - 1 \text{ is divisible by } 3$$

$$P(n): n \text{ cents can be obtained using } 3\text{¢ and } 5\text{¢ coins.}$$

- A **proof by mathematical induction**: shows that a given property  $P(n)$  is true for all integers greater than or equal to some initial integer.



---

## Principle of Mathematical Induction

Let  $P(n)$  be a property that is defined for integers  $n$ , and let  $a$  be a fixed integer. Suppose the following two statements are true:

1.  $P(a)$  is true.
2. For all integers  $k \geq a$ , if  $P(k)$  is true then  $P(k + 1)$  is true.

Then the statement

for all integers  $n \geq a$ ,  $P(n)$

is true.



# Outline of Proof by Mathematical Induction

## Method of Proof by Mathematical Induction

Consider a statement of the form, “For all integers  $n \geq a$ , a property  $P(n)$  is true.”

To prove such a statement, perform the following two steps:

**Step 1 (basis step):** Show that  $P(a)$  is true.

**Step 2 (inductive step):** Show that for all integers  $k \geq a$ , if  $P(k)$  is true then  $P(k + 1)$  is true. To perform this step,

**suppose** that  $P(k)$  is true, where  $k$  is any particular but arbitrarily chosen integer with  $k \geq a$ .

*[This supposition is called the **inductive hypothesis**.]*

Then

**show** that  $P(k + 1)$  is true.



# Mathematical Induction: Example

**Example:** Prove that for all integers  $n \geq 1$ ,

$$1 + 3 + 5 + 7 + \dots + (2n - 1) = n^2.$$

**Proof:** Let the property  $P(n)$  be the equation

$$1 + 3 + 5 + 7 + \dots + (2n - 1) = n^2 \quad \leftarrow \text{The property } P(n)$$


Show that the property is true for  $n = 1$ :

Basis Step

When  $n = 1$ , the property is the equation  $1 = 1^2$ .

But the left-hand side (LHS) of this equation is 1, and the right-hand side (RHS) is  $1^2$ , which equals 1 also.


So the property is true for  $n = 1$ .



## Inductive Step for the proof that for all integers $n \geq 1$ , $1 + 3 + 5 + 7 + \dots + (2n - 1) = n^2$ .

- Show that  $\forall$  integers  $k \geq 1$ , if  $p(k)$  is true then it is true for  $p(k+1)$ :
- Let  $k$  be any integer with  $k \geq 1$ , and suppose that the property is true for  $n = k$ . In other words, **suppose** that
- $1 + 3 + 5 + 7 + \dots + (2k - 1) = k^2$ . Inductive Hypothesis
- We must show that the property is true for  $n = k + 1$ .
- $P(k+1) = (k + 1)^2$ ,
- or, equivalently, we must **show** that
- $1 + 3 + 5 + 7 + \dots + (2(k + 1) - 1) = (k + 1)^2$ .





**Inductive hypothesis:**  $1 + 3 + 5 + 7 + \dots + (2k - 1) = k^2$ .

**Show:**  $1 + 3 + 5 + 7 + \dots + (2k + 1) = (k + 1)^2$ .

But the LHS of the equation to be shown is

$$1 + 3 + 5 + 7 + \dots + (2(k + 1) - 1)$$

$$= 1 + 3 + 5 + 7 + \dots + (2k - 1) + (2(k + 1) - 1)$$

*by making the next-to-last-term explicit*

$$= k^2 + (2k + 1) \quad \text{by substitution from the inductive hypothesis}$$

$$= (k + 1)^2 \quad \text{by algebra,}$$

which equals the RHS of the equation to be shown.

So, the property is true for  $n = k + 1$ .

*Therefore the property  $P(n)$  is true.*



# Proving sum of integers and geometric sequence

**Formula for the sum of the first  $n$  integers:** For all integers  $n \geq 1$ ,

$$1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}.$$

**Formula for the sum of the terms of a geometric sequence:** For all real numbers  $r \neq 1$  and all integers  $n \geq 0$ ,

$$1 + r + r^2 + r^3 + \dots + r^n = \frac{r^{n+1} - 1}{r - 1}.$$

# Example

## Theorem 5.2.2 Sum of the First $n$ Integers

For all integers  $n \geq 1$ ,

$$1 + 2 + \cdots + n = \frac{n(n+1)}{2}.$$

**Proof (by mathematical induction):**

Let the property  $P(n)$  be the equation

$$1 + 2 + 3 + \cdots + n = \frac{n(n+1)}{2}. \quad \leftarrow P(n)$$

**Show that  $P(1)$  is true:**

To establish  $P(1)$ , we must show that

$$1 = \frac{1(1+1)}{2} \quad \leftarrow P(1)$$

But the left-hand side of this equation is 1 and the right-hand side is

$$\frac{1(1+1)}{2} = \frac{2}{2} = 1$$

also. Hence  $P(1)$  is true.

**Show that for all integers  $k \geq 1$ , if  $P(k)$  is true then  $P(k+1)$  is also true:**

[Suppose that  $P(k)$  is true for a particular but arbitrarily chosen integer  $k \geq 1$ .

That is:] Suppose that  $k$  is any integer with  $k \geq 1$  such that

$$1 + 2 + 3 + \cdots + k = \frac{k(k+1)}{2} \quad \leftarrow P(k) \text{ inductive hypothesis}$$

[We must show that  $P(k+1)$  is true. That is:] We must show that

$$1 + 2 + 3 + \cdots + (k+1) = \frac{(k+1)[(k+1)+1]}{2},$$

or, equivalently, that

$$1 + 2 + 3 + \cdots + (k+1) = \frac{(k+1)(k+2)}{2}. \quad \leftarrow P(k+1)$$

# Exercises

The left-hand side of  $P(k + 1)$  is

$$1 + 2 + 3 + \cdots + (k + 1)$$

$$= 1 + 2 + 3 + \cdots + k + (k + 1) \quad \text{by making the next-to-last term explicit}$$

$$= \frac{k(k + 1)}{2} + (k + 1) \quad \text{by substitution from the inductive hypothesis}$$

$$= \frac{k(k + 1)}{2} + \frac{2(k + 1)}{2}$$

$$= \frac{k^2 + k}{2} + \frac{2k + 2}{2}$$

$$= \frac{k^2 + 3k + 1}{2} \quad \text{by algebra.}$$

And the right-hand side of  $P(k + 1)$  is

$$\frac{(k + 1)(k + 2)}{2} = \frac{k^2 + 3k + 1}{2}.$$

Thus the two sides of  $P(k + 1)$  are equal to the same quantity and so they are equal to each other. Therefore the equation  $P(k + 1)$  is true *[as was to be shown]*.

*[Since we have proved both the basis step and the inductive step, we conclude that the theorem is true.]*



## • Definition Closed Form

If a sum with a variable number of terms is shown to be equal to a formula that does not contain either an ellipsis or a summation symbol, we say that it is written **in closed form**.

- Evaluate  $2 + 4 + 6 + \cdots + 500$ .
- Evaluate  $5 + 6 + 7 + 8 + \cdots + 50$ .
- For an integer  $h \geq 2$ , write  $1 + 2 + 3 + \cdots + (h - 1)$  in closed form.

### Solution

- $$\begin{aligned} 2 + 4 + 6 + \cdots + 500 &= 2 \cdot (1 + 2 + 3 + \cdots + 250) \\ &= 2 \cdot \left( \frac{250 \cdot 251}{2} \right) && \text{by applying the formula for the sum} \\ &= 62,750. && \text{of the first } n \text{ integers with } n = 250 \end{aligned}$$
- $$\begin{aligned} 5 + 6 + 7 + 8 + \cdots + 50 &= (1 + 2 + 3 + \cdots + 50) - (1 + 2 + 3 + 4) \\ &= \frac{50 \cdot 51}{2} - 10 && \text{by applying the formula for the sum} \\ &= 1,265 && \text{of the first } n \text{ integers with } n = 50 \end{aligned}$$
- $$\begin{aligned} 1 + 2 + 3 + \cdots + (h - 1) &= \frac{(h - 1) \cdot [(h - 1) + 1]}{2} && \text{by applying the formula for the sum} \\ &= \frac{(h - 1) \cdot h}{2} && \text{of the first } n \text{ integers with} \\ & && n = h - 1 \\ & && \text{since } (h - 1) + 1 = h. \quad \blacksquare \end{aligned}$$

# Proving Sum of Geometric Sequences

$$\sum_{i=0}^n r^i = \frac{r^{n+1} - 1}{r - 1}.$$

$$\sum_{i=0}^0 r^i = \frac{r^{0+1} - 1}{r - 1} \leftarrow P(0) = \frac{r - 1}{r - 1} = 1$$

$$\sum_{i=0}^k r^i = \frac{r^{k+1} - 1}{r - 1} \leftarrow P(k) \text{ inductive hypothesis}$$

$$\sum_{i=0}^{k+1} r^i = \frac{r^{k+2} - 1}{r - 1}. \leftarrow P(k+1)$$

$$\begin{aligned} &= \sum_{i=0}^k r^i + r^{k+1} \\ &= \frac{r^{k+1} - 1}{r - 1} + r^{k+1} \\ &= \frac{r^{k+1} - 1}{r - 1} + \frac{r^{k+1}(r - 1)}{r - 1} \\ &= \frac{(r^{k+1} - 1) + r^{k+1}(r - 1)}{r - 1} \\ &= \frac{r^{k+1} - 1 + r^{k+2} - r^{k+1}}{r - 1} \\ &= \frac{r^{k+2} - 1}{r - 1} \end{aligned}$$

### Theorem 5.2.3 Sum of a Geometric Sequence

For any real number  $r$  except 1, and any integer  $n \geq 0$ ,

$$\sum_{i=0}^n r^i = \frac{r^{n+1} - 1}{r - 1}.$$

#### Proof (by mathematical induction):

Suppose  $r$  is a particular but arbitrarily chosen real number that is not equal to 1, and let the property  $P(n)$  be the equation

$$\sum_{i=0}^n r^i = \frac{r^{i+1} - 1}{r - 1} \quad \leftarrow P(n)$$

We must show that  $P(n)$  is true for all integers  $n \geq 0$ . We do this by mathematical induction on  $n$ .

*Show that  $P(0)$  is true:*

To establish  $P(0)$ , we must show that


$$\sum_{i=0}^0 r^i = \frac{r^{0+1} - 1}{r - 1} \quad \leftarrow P(0)$$

The left-hand side of this equation is  $r^0 = 1$  and the right-hand side is

$$\frac{r^{0+1} - 1}{r - 1} = \frac{r - 1}{r - 1} = 1$$

also because  $r^1 = r$  and  $r \neq 1$ . Hence  $P(0)$  is true.





*Show that for all integers  $k \geq 0$ , if  $P(k)$  is true then  $P(k + 1)$  is also true:*

*[Suppose that  $P(k)$  is true for a particular but arbitrarily chosen integer  $k \geq 0$ . That is:]*

Let  $k$  be any integer with  $k \geq 0$ , and suppose that

$$\sum_{i=0}^k r^i = \frac{r^{k+1} - 1}{r - 1} \quad \leftarrow \begin{array}{l} P(k) \\ \text{inductive hypothesis} \end{array}$$

[We must show that  $P(k + 1)$  is true. That is:] We must show that

$$\sum_{i=0}^{k+1} r^i = \frac{r^{(k+1)+1} - 1}{r - 1},$$

or, equivalently, that

$$\sum_{i=0}^{k+1} r^i = \frac{r^{k+2} - 1}{r - 1}. \quad \leftarrow P(k + 1)$$

[We will show that the left-hand side of  $P(k + 1)$  equals the right-hand side.]

The left-hand side of  $P(k + 1)$  is

$$\sum_{i=0}^{k+1} r^i = \sum_{i=0}^k r^i + r^{k+1}$$

by writing the  $(k + 1)$ st term separately from the first  $k$  terms

$$= \frac{r^{k+1} - 1}{r - 1} + r^{k+1}$$

by substitution from the inductive hypothesis

$$= \frac{r^{k+1} - 1}{r - 1} + \frac{r^{k+1}(r - 1)}{r - 1}$$

by multiplying the numerator and denominator of the second term by  $(r - 1)$  to obtain a common denominator

$$= \frac{(r^{k+1} - 1) + r^{k+1}(r - 1)}{r - 1}$$

by adding fractions

$$= \frac{r^{k+1} - 1 + r^{k+2} - r^{k+1}}{r - 1}$$

by multiplying out and using the fact that  $r^{k+1} \cdot r = r^{k+1} \cdot r^1 = r^{k+2}$


$$= \frac{r^{k+2} - 1}{r - 1}$$

by canceling the  $r^{k+1}$ 's.

which is the right-hand side of  $P(k + 1)$  [as was to be shown.]

[Since we have proved the basis step and the inductive step, we conclude that the theorem

is true.]



a.  $1 + 3 + 3^2 + \dots + 3^{m-2} = 3^{(m-2)+1} - 1/3 - 1$

by applying the formula for the sum of a geometric sequence with  $r = 3$  and

$$n = m - 2$$
$$= \frac{3^{m-1} - 1}{2}$$

b.

$$3^2 + 3^3 + 3^4 + \dots + 3^m = 3^2 \cdot (1 + 3 + 3^2 + \dots + 3^{m-2})$$

by factoring out  $3^2$

$$= 9 \times \frac{3^{m-1} - 1}{2} \text{ by part (a).}$$



# Mathematics in Programming

Example : Finding the sum of a integers

**Same Question:** Prove that these programs prints the same results in case  $n \geq 1$

```
For (i=1, i≤n; i++)  
    S=S+i;  
Print ("%d", S);
```

```
S=(n(n+1))/2  
Print ("%d",S);
```



# Mathematics in Programming

## Example : Finding the sum of a geometric series

Prove that these codes will return the same output.

```
int n, r, sum=0;
int i;
scanf("%d",&n);
scanf("%d",&r);

if(r != 1) {
    for(i=0 ; i<=n ; i++) {
        sum = sum + pow(r,i);
    }
    printf("%d\n", sum);
}
```

```
int n, r, sum=0;
scanf("%d",&n);
scanf("%d",&r);

if(r != 1) {
    sum=((pow(r,n+1))-1)/(r-1);
    printf("%d\n", sum);
}
```



5.3

---

# Mathematical Induction II

## Proving Divisibility



# Mathematics in Programming

## Proving Divisibility Property

What will the output of this program be for any input  $n$ ?

```
int n;  
scanf("%d",&n);  
  
if(n >= 0) {  
    if( (pow(2,(2*n)) - 1) %3 == 0)  
        printf("this property is true");  
    else  
        printf("this property isn't true");  
}
```

\\ does  $2^{2n} - 1 \mid 3??$  \\

# Proving a Divisibility Property

For all integers  $n \geq 0$ ,  $2^{2n} - 1$  is divisible by 3.

$$3 \mid 2^{2n} - 1 \quad \leftarrow P(n)$$

**Basis Step:** Show that  $P(0)$  is true.

$$P(0): 2^{2 \cdot 0} - 1 = 2^0 - 1 = 1 - 1 = 0 \text{ as } 3 \mid 0, \text{ thus } P(0) \text{ is true.}$$

**Inductive Step:** Show that for all integers  $k \geq 0$ , if  $P(k)$  is true then  $P(k + 1)$  is also true:

Suppose:  $2^{2k} - 1$  is divisible by 3.  $\leftarrow P(k)$  inductive hypothesis

$$2^{2k} - 1 = 3r \text{ for some integer } r.$$

We want to prove  $2^{2(k+1)} - 1$  is divisible by 3.  $\leftarrow P(k+1)$

$$\begin{aligned} 2^{2(k+1)} - 1 &= 2^{2k+2} - 1 && \text{by the laws of exponents} \\ &= 2^{2k} \cdot 2^2 - 1 = 2^{2k} \cdot 4 - 1 \\ &= 2^{2k}(3 + 1) - 1 = 2^{2k} \cdot 3 + (2^{2k} - 1) = 2^{2k} \cdot 3 + 3r \\ &= 3(2^{2k} + r) && \text{Which is integer} \end{aligned}$$

so, by definition of divisibility,  $2^{2(k+1)} - 1$  is divisible by 3





Outline a proof by math induction for the statement:

For all integers  $n \geq 0$ ,  $5^n - 1$  is divisible by 4.

Proof by mathematical induction:

Let the property  $P(n)$  be the sentence

$5^n - 1$  is divisible by 4. ← *the property  $P(n)$*

Show that the property is true for  $n = 0$ :

We must show that  $5^0 - 1$  is divisible by 4.

But  $5^0 - 1 = 1 - 1 = 0$ , and 0 is divisible by 4 because  $0 = 4 \cdot 0$ .


Show that for all integers  $k \geq 0$ , if the property is true for  $n = k$ , then it is true for  $n = k + 1$ :

Let  $k$  be an integer with  $k \geq 0$ , and **suppose** that  
*[the property is true for  $n = k$ .*

$5^k - 1$  is divisible by 4. ← *inductive hypothesis*

We must **show** that  $P(k + 1)$  is true.

$5^{k+1} - 1$  is divisible by 4.



## Scratch Work for proving that

For all integers  $n \geq 0$ ,  $5^n - 1$  is divisible by 4.

---

$$\begin{aligned}5^{k+1} - 1 &= 5^k \cdot 5 - 1 \\ &= 5^k \cdot (4 + 1) - 1 \\ &= 5^k \cdot 4 + 5^k \cdot 1 - 1 \\ &= 5^k \cdot 4 + (5^k - 1)\end{aligned}$$

**Note:** Each of these terms is divisible by 4.

**So:**  $5^{k+1} - 1 = 5^k \cdot 4 + 4 \cdot r$  (where  $r$  is an integer)

$$= 4 \cdot (5^k + r)$$

$(5^k + r)$  is an integer because it is a sum of products of integers, and so, by definition of divisibility  $5^{k+1} - 1$  is divisible by 4.

# Proving Inequality

For all integers  $n \geq 3$ ,  $2n + 1 < 2^n$

Let  $P(n)$  be  $2n+1 < 2^n$

**Basis Step:** Show that  $P(3)$  is true.  $P(3)$ :  $2 \cdot 3 + 1 < 2^3$  which is true.

**Inductive Step:** Show that for all integers  $k \geq 3$ , if  $P(k)$  is true then  $P(k + 1)$  is also true:

Suppose:  $2k + 1 < 2^k$  is true  $\leftarrow P(k)$  inductive hypothesis

$$2(k+1) + 1 < 2^{k+1} \leftarrow P(k+1)$$

$$2k+3 = (2k+1) + 2 \quad \text{by algebra}$$

$$< 2^k + 2^k \quad \text{as } 2k + 1 < 2^k \text{ by the hypothesis} \\ \text{and because } 2 < 2^k \quad (k \geq 2)$$

$$\therefore 2k + 3 < 2 \cdot 2^k = 2^{k+1}$$

*[This is what we needed to show.]*



# Exercise

---

For each positive integer  $n$ , let  $P(n)$  be the property

$$2^n < (n + 1)!$$



# Proving a Property of a Sequence

Define a sequence  $a_1, a_2, a_3 \dots$  as follows:

$$a_1 = 2$$

$$a_k = 5a_{k-1} \quad \text{for all integers } k \geq 2.$$

$$a_1 = 2$$

$$a_2 = 5a_{2-1} = 5a_1 = 5 \cdot 2 = 10$$

$$a_3 = 5a_{3-1} = 5a_2 = 5 \cdot 10 = 50$$

$$a_4 = 5a_{4-1} = 5a_3 = 5 \cdot 50 = 250$$

Property  $\rightarrow$  The terms of the sequence satisfy the equation

$$a_n = 2 \cdot 5^{n-1}$$

# Proving a Property of a Sequence

Prove this property:

$$a_n = 2 \cdot 5^{n-1} \text{ for all integers } n \geq 1$$

**Basis Step:** Show that  $P(1)$  is true.  $a_1 = 2 \cdot 5^{1-1} = 2 \cdot 5^0 = 2$

**Inductive Step:** Show that for all integers  $k \geq 1$ , if  $P(k)$  is true then  $P(k+1)$  is also true:

Suppose:  $a_k = 2 \cdot 5^{k-1}$

←  $P(k)$  inductive hypothesis

$$a_{k+1} = 2 \cdot 5^k$$

←  $P(k+1)$

$$= 5a_{(k+1)-1}$$

by definition of  $a_1, a_2, a_3 \dots$

$$= 5a_k$$

$$= 5 \cdot (2 \cdot 5^{k-1})$$

by the hypothesis

$$= 2 \cdot (5 \cdot 5^{k-1})$$

$$= 2 \cdot 5^k$$

*[This is what we needed to show.]*



# Important Formulas

---

**Formula for the sum of the first  $n$  integers:** For all integers  $n \geq 1$ ,

$$1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}.$$

**Formula for the sum of the terms of a geometric sequence:** For all real numbers  $r \neq 1$  and all integers  $n \geq 0$ ,

$$1 + r + r^2 + r^3 + \dots + r^n = \frac{r^{n+1} - 1}{r - 1}.$$

# Exercises

$$\text{a. } 1 + 2 + 3 + \dots + 100 = \frac{100(100 + 1)}{2} = 50(101) = 5050$$

$$\text{b. } 1 + 2 + 3 + \dots + k = \frac{k(k + 1)}{2}$$

$$\text{c. } 1 + 2 + 3 + \dots + (k - 1) = \frac{(k - 1)((k - 1) + 1)}{2} = \frac{(k - 1)k}{2}$$

$$\begin{aligned} \text{d. } 4 + 5 + 6 + \dots + (k - 1) &= (1 + 2 + 3 + \dots + (k - 1)) - (1 + 2 + 3) \\ &= \frac{k(k - 1)}{2} - (1 + 2 + 3) = \frac{k(k - 1)}{2} - 6 \end{aligned}$$

$$\begin{aligned} \text{e. } 3 + 3^2 + 3^3 + \dots + 3^k &= (1 + 3 + 3^2 + 3^3 + \dots + 3^k) - 1 = \frac{3^{k+1} - 1}{3 - 1} - 1 \\ &= \frac{3^{k+1} - 1}{2} - 1 = \frac{3^{k+1} - 1}{2} - \frac{2}{2} = \frac{3^{k+1} - 3}{2} \end{aligned}$$

$$\begin{aligned} \text{f. } 3 + 3^2 + 3^3 + \dots + 3^k &= 3(1 + 3 + 3^2 + \dots + 3^{k-1}) \\ &= 3 \left( \frac{3^{(k-1)+1} - 1}{3 - 1} \right) = \frac{3(3^k - 1)}{2} \end{aligned}$$